

Nota

Clutch size and hatchling morphology of *Erythrolamprus poecilogyrus schotti* (Serpentes: Dipsadidae: Xenodontinae) from northeastern Brazil

Tamaño de puesta y morfología de neonatos de *Erythrolamprus poecilogyrus schotti* (Serpentes: Dipsadidae: Xenodontinae) en el noroeste de Brasil

Élida Francisco da Silva  ^a, Mateus Lima Dutra ^a

Alerandro Soares da Silva ^a, Jéssica Monique da Silva Amaral ^a

Vanessa do Nascimento Barbosa ^a, Frederico Gustavo Rodrigues França ^a

^a Universidade Federal da Paraíba, Brazil

Recibido: junio 30, 2024

Aceptado: abril 29, 2025

Publicado en línea: agosto 5, 2025

<https://doi.org/10.21068/2539200X.1267>



Abstract

We report on the clutch size and neonate morphology of *Erythrolamprus poecilogyrus schotti*, an oviparous snake species. A clutch of seven eggs was found in a rural area of Paraíba, Brazil. Neonates had an average total length of 181.29 mm and an average weight of 3.0 g, with females being larger sizes than males—a pattern consistent with the species' known reproductive traits. Defensive behaviors observed in the neonates included cloacal discharge and substrate-burrowing. These findings offer new insights into the species' natural history and highlight its early-life defensive strategies and developmental characteristics.

Keywords: reproductive ecology, embryonic development, sexual dimorphism, neonate behavior, Atlantic Forest.

Resumen

Se reporta el tamaño de la puesta y la morfología de neonatos de *Erythrolamprus poecilogyrus schotti*, una especie de serpiente ovípara. Se encontró una puesta de siete huevos en una zona rural de Paraíba, Brasil. Los neonatos presentaron una longitud total promedio de 181,29 mm y un peso promedio de 3,0 g, con las hembras siendo de mayor tamaño que los machos, un patrón consistente con los rasgos reproductivos conocidos de la especie. Entre los comportamientos defensivos observados en los neonatos se incluyen la descarga cloacal y el ocultamiento en el sustrato. Estos hallazgos aportan nuevos datos sobre la historia natural de la especie y destacan sus estrategias defensivas y características de desarrollo en etapas tempranas.

Palabras clave: ecología reproductiva, desarrollo embrionario, dimorfismo sexual, bosque atlántico, comportamiento de neonatos.

Introduction

Information on the timing and duration of egg-laying, clutch size, egg dimensions, incubation period, and offspring morphology is essential for studies on snake population ecology and for understanding species' life history (Qu et al., 2011; Amaral et al., 2020). However, such data are difficult to obtain due to the need to locate gravid females in the wild and/or find eggs and monitor their development through hatching. Despite these challenges, there has been a recent increase in studies documenting early reproductive traits in oviparous Brazilian snakes. These studies span various families and genera (Pizzatto & Marques, 2002; Scartozzoni et al., 2005; Morais et al., 2018; Muscat et al., 2019; Amaral et al., 2020; Guedes & Guedes, 2020; Amaral et al., 2021; Marques et al., 2021; Amaral et al., 2022; Barbosa et al., 2023).

The genus *Erythrolamprus* Boie, 1826 (family Dipsadidae, subfamily Xenodontinae) comprises 55 species distributed throughout South America and the Antilles. It represents one of the most morphologically diverse radiations of New World snakes (Dixon, 1989; Ascenso et al., 2019; Uetz et al., 2024). *Erythrolamprus poecilogyrus* (Wied-Newied, 1824) occurs from southeastern Venezuela and eastern Guyana to Pampas and Chaco regions of Argentina (Fernandes, 2006; Uetz et al., 2024). Due to its considerable variation in coloration, meristic patterns, and morphology, Dixon &

Markezich (1992) recognized four subspecies: *E. p. caesius* (Cope, 1862), found in the Pantanal and Chaco; *E. p. sublineatus* (Cope, 1860), in the Pampas; *E. p. poecilogyrus* (Wied-Newied, 1825), mainly in the Central and Southern Atlantic Forest; and *E. p. schotti* (Schlegel, 1837) in the Amazon, cerrado, caatinga, and the Northeastern Atlantic Forest.

All subspecies of *Erythrolamprus poecilogyrus* are terrestrial, active both during the day and at night, and primarily feed on amphibians, with occasional consumption of fish, lizards, and invertebrates (Prieto et al., 2012; Corrêa et al., 2016; Marques et al., 2017; Oliveira et al., 2021). Although the species is oviparous, information on its reproductive biology remains limited. Pizzatto et al. (2008) and Quintella et al. (2017) reported that follicular cycles in *E. p. poecilogyrus* and *E. p. schotti* can be continuous in southeastern Brazil, whereas in *E. p. sublineatus* from southern Brazil, reproduction is strictly seasonal and occurs during the dry season. Clutch size varies among subspecies: *E. p. poecilogyrus* may lay 2-9 or 6-15 eggs depending on the population (Pinto & Fernandes, 2004; Alencar & Nascimento, 2014); *E. p. caesius* has been recorded with 8-9 eggs (Campbell & Murphy, 1984); *E. p. sublineatus* with 2-9 eggs; and *E. p. schotti* with 3-17 eggs (Vitt & Vangilder, 1984; Mesquita et al., 2013). Herein, we present data on clutch size and hatchling morphology of *E. p. schotti* from a region of the Northeastern Atlantic Forest.

Materials and methods

On August 8, 2023, a clutch of seven eggs was found in a rural area in the municipality of Mamanguape, Paraíba, northeastern Brazil (6°39.44'S, 35°08'55'W; elevation: 30 m). According to the Köppen climate classification, the region has a hot and humid climate (type Aw), with a dry season in summer and a rainy season in winter. The eggs were partially protected inside a hole in the ground and were collected under SISBIO License 74327-1, then transported to the Laboratório de Ecologia Animal at the Universidade Federal da Paraíba. Each egg was measured for length and width using a digital caliper (precision: 0.01 mm) and weighed with a precision digital scale (0.1 g). The eggs were incubated at 20-22 °C with approximately 60% humidity, semi-buried in vermiculite (Figure 1A) in a rounded glass container (185 mm height x 130 mm diameter).

Immediately after birth, all hatchlings were measured using a digital caliper (precision: 0.01 mm), weighed using a spring scale (0.1 g), and sexed using a probe. The recorded measurements included snout-vent length (SVL), tail length (TL), body width (BW), body height (BH), head width (HW), head height (HH), head length (HL), distance between the nostrils (NW), eye diameter (ED) and mass (M). One individual was preserved in 10% formalin and deposited in the Herpetological Collection of the Universidade Federal da Paraíba (CHUFPB-32179). The remaining hatchlings were released at the Guaribas Biological Reserve in the municipality of Mamanguape.

Results

The eggs averaged 26.1 mm in length (SD = 1.2), 15.1 mm in width (SD = 0.5), and 3.9 g in mass (SD = 0.4). After 37 days of incubation, the eggs began to pip, and all snakes successfully hatched. On 13 September 2023, at 21:41, five offspring began hatching (Figure 1B). On 14 September 2023, at 08:39, the first snake fully emerged from its egg after 10 hours and 58 minutes (Figure 1C). The

second snake emerged at 10:44, after 13 hours and 3 minutes. The third and fourth snakes emerged simultaneously at 11:49, after 14 hours and 8 minutes. The fifth snake emerged at 17:03, after 19 hours and 22 minutes.

The last two eggs began to pipe only on 14 September 2023, at 09:40 and 12:03, respectively. The sixth snake emerged at 04:30 on 15 September, after 18 hours and 50 minutes. The seventh snake emerged at 14:59 on 14 September, taking only 2 hours and 56 minutes to fully emerge from the egg (Table 1).

The hatchlings' mean total length was 181.29 mm (SD = 7.4), with an average mass of 3.0 g (SD = 0.2). Of the seven snakes, five were females and two were males (Table 2). All individuals displayed the typical coloration of juvenile *E. p. schotti*, characterized by a brown head with black spots, a light cream dorsum with black spots extending to the tail, and a light cream venter with sparse black edging or black bands (Figure 1D). After hatching, all the snakes exhibited escape behavior, burying themselves in the vermiculite substrate.

Discussion

Compared with data available in the literature, the clutch size of *Erythrolamprus poecilogyrus schotti* is consistent with that reported for other subspecies of *E. poecilogyrus*, which ranges from 2 to 17 eggs per clutch (Vitt & Vangilder, 1983; Pinto & Fernandes, 2004; Alencar & Nascimento, 2014). The mean snout-vent length (SVL) of the newborns was slightly higher than that reported by Vitt & Vangilder (1983), who documented an SVL of 136.3 mm for three hatchlings. Additionally, females were slightly larger than males, a pattern also observed in *E. poecilogyrus*, as distinct male-female size ratios have been reported for *E. p. poecilogyrus*, *E. p. sublineatus* and *E. p. schotti* (Alencar & Nascimento, 2014; Quintela et al., 2017).

Although *E. p. schotti* does not exhibit seasonal reproduction (Pizzatto et al., 2008), slight variations between populations in different locations may

occur. The clutch was found during the wet season in the northeastern Atlantic Forest, a period that may favor the development of eggs and hatchling survival (França et al., 2023). The high humidity characteristic of this region likely promotes adequate egg hydration, thereby optimizing embryonic development. Moreover, environments with elevated humidity may enhance phenotypic and immunological traits in some species (Ji & Du, 2001; Brown & Shine, 2006; Brown & Shine, 2018).

Snakes of the genus *Erythrolamprus* typically exhibit non-aggressive defensive behavior, generally avoiding biting even when handled—*E. aesculapii* being a notable exception—and fleeing when threatened (Marques & Sazima, 2004; Marques et al., 2019). All *E. p. schotti* newborns displayed escape behaviors: when handled, they released cloacal discharge and rotated their bodies but did not strike or attempt to bite. These observations suggest an innate preference for evasion over offensive defense mechanisms in *E. p. schotti* (Greene, 1988).

Acknowledgements

FGRF thanks the Universidade Federal da Paraíba (UFPB) for support through Edital PROPESQ/PRPG/UFPB No. 03/2020-PVP13459-2020). VNB and EFS thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the master's and PhD scholarships (grant numbers 88887.935257/2024-00 and 88887.713778/2022-00, respectively). We also thank the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBIO) for the collection permit (SISBIO nº 74327-1).

Table 1. Hatching times and birth order of *Erythrolamprus poecilogyrus schotti* hatchlings.

ID	CR	Date CR	HT	Date HT	CR to HT
1	12:03	2023-09-14	14:59	2023-09-14	2h and 56 min
2	09:40	2023-09-14	04:30	2023-09-15	18h and 50 min
3	21:41	2023-09-13	17:03	2023-09-14	19h and 22 min
4	21:41	2023-09-13	08:39	2023-09-14	10h and 58 min
5	21:41	2023-09-13	11:49	2023-09-14	14h and 8 min
6	21:41	2023-09-13	10:44	2023-09-14	13h and 3 min
7	21:41	2023-09-13	11:49	2023-09-14	14h and 8 min


Notes. ID = egg identification; CR = time when the eggshell was cracked (pierced by the hatchling); Date CR = date when the eggshell was cracked; HT = time of hatching (emergence from the egg); Date HT = date of hatching; CR to HT = total hatching duration (time from shell cracking to complete emergence).

Table 2. Morphometry (mm) and mass (g) of *Erythrolamprus poecilogyrus schotti* hatchlings.

ID	SEX	SLV	TL	BW	BH	HL	HH	HW	NW	ED	M
1	M	144	32	4.15	5.14	12.41	4.20	6.04	2.10	3.37	2.5
2	F	152	36	4.38	5.26	11.20	3.82	5.45	2.11	2.19	3.2
3	F	148	34	4.05	4.70	12.51	3.77	5.65	2.09	2.36	3.0
4	M	142	28	4.09	4.75	11.40	4.44	6.29	2.39	2.75	3.0
5	F	155	36	5.00	5.32	12.70	3.54	5.71	2.00	2.21	3.1
6	F	146	31	4.95	5.15	11.64	4.20	5.89	2.31	2.43	3.2
7	F	153	32	4.83	5.47	12.11	3.94	6.50	1.96	2.33	3.0
Mean		148.57	32.71	4.49	5.11	11.99	3.98	5.93	2.13	2.37	3.0
SD		4.89	2.87	0.42	0.28	0.58	0.30	0.37	0.15	0.18	0.23

Notes. SVL = snout-vent length; TL = tail length; BW = body width; BH = body height; HW = head width; HH = head height; HL = head length; NW = inter-nasal distance; ED = eye diameter; DN = distance between the navel and the cloaca; M = mass; Sd = Standard deviation.

Figure 1. Eggs, pipping and hatchling of *Erythrolamprus poecilogyrus schotti*.



Notes. A) The seven eggs during incubation; B) Egg pipping. The red arrow indicates hatchling's head emerging; C) A newborn fully emerged from the egg; D) A neonate of *E. p. schotti* showing typical post-hatching coloration.

References

- Alencar, L. R. V., & Nascimento, L. B. (2014). Natural history data of a common snake suggest interpopulational variation and conservatism in life history traits: the case of *Erythrolamprus poecilogyrus*. *The Herpetological Journal*, 24(2), 79-85.
- Amaral, J. M. S., Araújo, P., Barbosa, V. N., & França, F. G. R. (2020). First record of egg development time and hatchling morphology of *Amerotyphlops paucisquamis* (Dixon & Hendricks, 1979) (Scolophoridae: Typhlopidae) from northeast Brazil. *Revista de Ciências Ambientais*, 14(3), 79-83.
- Amaral, J. M. S., Barbosa, V. do N., Sampaio, I. L. R., Morais, M. da S. R. de, & França, F. G. R. (2021). Growth rates and prey-handling behavior of hatchling *Helicops angulatus* (Linnaeus, 1758) (Serpentes; Hydropsini) in the atlantic forest of northeastern Brazil. *Acta Ambiental Catarinense*, 18(1), 97-104.
- Amaral, J. M. S., Barbosa, V. N., & França, F. G. R. (2022). Clutch size, hatching time, and hatchling morphology of Boettger's Sipo *Chironius flavolineatus* Jan 1863 (Serpentes: Colubridae: Colubrini) in the northeast Atlantic Forest. *North-Western Journal of Zoology*, 18(2), 237-239.
- Ascenso, A. C., Costa, J. C., & Prudente, A. L. (2019). Taxonomic revision of the *Erythrolamprus reginae* species group, with description of a new species from Guiana Shield (Serpentes: Xenodontinae). *Zootaxa*, 4586(1), 65-97. <https://doi.org/10.11646/zootaxa.4586.1.3>
- Barbosa, V. N., Silva, A. S., Silva, A. S., Silva, É. F., Linnik, J., Dutra, M. L., & França, F. G. R. (2023). Clutch size and hatchling morphology of Parrot snake *Leptophis liocercus* in the north-east Atlantic Forest of Brazil. *Herpetological Bulletin*, 166, 38-39. <https://doi.org/10.33256/hb166.3839>
- Brown, G. P., & Shine, R. (2006). Effects of nest temperature and moisture on phenotypic traits of hatchling snakes (*Tropidonophis mairii*, Colubridae) from tropical Australia. *Biological Journal of the Linnean Society*, 89, 159-168.
- Brown, G. P., & Shine, R. (2018). Immune configuration in hatchling snakes is affected by incubation moisture, and is linked to subsequent growth and survival in the field. *Journal of Experimental Zoology - A*, 329, 222-229.
- Campbell, J. A., & Murphy, J. E. (1984). Reproduction in five species of Paraguayan colubrids. *Transactions of the Kansas Academy of Science*, 87(1/2), 63-65. <https://doi.org/10.2307/3627766>
- Corrêa, D. N., Quintela, F. M., & Loebmann, D. (2016). Feeding ecology of *Erythrolamprus jaegeri jaegeri* (Günther, 1858) and *Erythrolamprus poecilogyrus sublineatus* (Cope, 1860) in the coastal zone of Subtropical Brazil (Serpentes, Dipsadidae). *Anais da Academia Brasileira de Ciências*, 88(1), 293-308. <https://doi.org/10.1590/0001-3765201520140570>
- Dixon, J. R. (1989). A key and checklist to the Neotropical snake genus *Liophis* with country lists and maps. *Smithsonian Herpetological Information Service*, 79, 1-40. <https://doi.org/10.5479/si.23317515.79.1>
- Dixon, J. R., & Marquezich, A. L. (1992). Taxonomy and geographic variation of *Liophis poecilogyrus* (Wied) from South America (Serpentes: Colubridae). *Texas Journal of Science*, 44, 131-131.
- Fernandes, D. S. (2006). *Revisão sistemática de Liophis poecilogyrus (Wied-Neuwied, 1825) (Serpentes: Colubridae)* [Doctoral dissertation, Universidade Federal do Rio de Janeiro].
- França, F. G. R., Vasconcellos, A., Alves, R. R. N., & Pereira-Filho, G. A. (2023). An introduction to the knowledge of animal diversity and conservation in the most threatened forests of Brazil. In G. A. Pereira-Filho, F. G. R. França, R. R. N. Alves, & A. Vasconcellos (Eds.), *Animal*

- biodiversity and conservation in Brazil's Northern Atlantic Forest (pp. 1-5). Springer.
- Greene, H. W. (1988). Antipredator mechanisms in reptiles. In C. Gans, & R. B. Huey (Eds.), *Biology of the Reptilian* (pp. 1-152). Alan R. Liss.
- Guedes, T. B., & Guedes, A. (2020). Notes on court and copula, fertility, nest, eggs and hatchlings of the Caatinga's black snake *Boiruna sertaneja* Zaher, 1996 (Serpentes: Dipsadidae) from northeastern Brazil. *Anais da Academia Brasileira de Ciências*, 92, e20190588. <https://doi.org/10.1590/0001-3765202020190588>
- Ji, X., & Du, W. G. (2001). The effects of thermal and hydric environments on hatching success, embryonic use of energy and hatchling traits in a colubrid snake, *Elaphe carinata*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 129(2-3), 461-471. [https://doi.org/10.1016/s1095-6433\(01\)00271-9](https://doi.org/10.1016/s1095-6433(01)00271-9)
- Marques, O. A. V., Eterovic, A., & Sazima, I. (2019). *Serpentes da Mata Atlântica: Guia ilustrado para a Serra do Mar*. Ponto A.
- Marques, O. A. V., & Sazima, I. (2004). História natural dos répteis da Estação Ecológica Juréia-Itatins. In O. A. V. Marques, & W. Dulepa (Eds.), *Estação Ecológica Juréia-Itatins: ambientes físicos, flora e fauna* (pp. 257-277). Holos.
- Marques, O. A. V., Sazima, I., & Guedes, T. B. (2017). *Serpentes da Caatinga: guia ilustrado*. Editora Ponto A.
- Marques, R., da Silva, W. P., & Coelho, F. E. A. (2021). Clutch size, morphology, hatching time and offspring behavior of *Pseudoboa nigra* (Dipsadidae: Pseudoboini) from Caatinga. *North-Western Journal of Zoology*, 17(1), 122-125.
- Mesquita, P. C. M. D., Passos, D. C., Borges-Nojosa, D. M., & Cechin, S. Z. (2013). Ecologia e história natural das serpentes de uma área de Caatinga no nordeste brasileiro. *Papéis Avulsos de Zoologia*, 53(8), 99-113. <https://doi.org/10.1590/S0031-12412013000800001>
- Morais, M. S. R., França, R. C., Delfim, F. R., & França, F. G. R. (2018). Eggs and hatchling morphometry of *Spilotes sulphureus* (Wagler in Spix, 1824) (Serpentes: Colubridae: Colubrinae: Colubroidea: Caenophidia) from northeastern Brazil. *Herpetology Notes*, 11, 441-444.
- Muscat, E., Rotenberg, E., & Machado, I. F. (2019). Oviposition observation of *Tropidodryas serra* (Serpentes: Dipsadidae). *Neotropical Biology and Conservation*, 14(1), 137-141. <https://doi.org/10.3897/neotropical.14.e35410>
- Oliveira, M. C., Duarte, R. G., Silva, E. G., Teles, D. A., Filho, J. A. A., & Almeida, W. O. (2021). First record of *Tropidurus hispidus* (Squamata: Tropiduridae) predation by *Erythrolamprus poecilogyrus* (Colubridae). *Herpetology Notes*, 14, 133-134.
- Pinto, R. R., & Fernandes, R. (2004). Reproductive biology and diet of *Liophis poecilogyrus poecilogyrus* (Serpentes, Colubridae) from southeastern Brazil. *Phyllomedusa: Journal of Herpetology*, 3(1), 9-14. <https://doi.org/10.11606/issn.2316-9079.v3i1p9-14>
- Pizzatto, L., & Marques, O. A. V. (2002). Reproductive biology of the false coral snake *Oxyrhopus guibei* (Colubridae) from southeastern Brazil. *Amphibia-Reptilia*, 23, 495-504.
- Pizzatto, L., Jordão, R. S., & Marques, O. A. V. (2008). Overview of Reproductive Strategies in Xenodontini (Serpentes: Colubridae: Xenodontinae) with New Data for *Xenodon Neuwiedii* and *Waglerophis Merremii*. *Journal of Herpetology*, 42(1), 153-162. <https://doi.org/10.1670/06-150R2.1>
- Prieto, Y. A., Giraudo, A. R., & López, M. S. (2012). Diet and sexual dimorphism of *Liophis poecilogyrus* (Serpentes, Dipsadidae) from the wetland regions of northeast Argentina. *Journal*

- of *Herpetology*, 46(3), 402-406.
<https://doi.org/10.1670/10-228>
- Qu, Y.-F., Li, H., Gao, J.-F., & Ji, X. (2011). Geographical variation in reproductive traits and trade-offs between size and number of eggs in the king ratsnake, *Elaphe carinata*: Female reproduction in a snake. *Biological Journal of the Linnean Society*, 104(3), 701-709.
<https://doi.org/10.1111/j.1095-8312.2011.01749.x>
- Quintela, F. M., Marques, W. C., & Loebmann, D. (2017). Reproductive biology of the Green Ground Snake *Erythrolamprus poecilogyrus sublineatus* (Serpentes: Dipsadidae) in Subtropical Brazil. *Anais Da Academia Brasileira de Ciências*, 89(3), 2189-2197.
<https://doi.org/10.1590/0001-3765201720160805>
- Scartozzoni, R. R., Almeida-Santos, S. M., & Salomão, M. G. (2005): Aspectos da reprodução da cobra bicuda *Oxybelis fulgidus* (Serpentes: Colubridae). *Publicações Avulsas do Instituto Pau Brasil*, 8(9), 85-90.
- Uetz, P., Freed, P., Aguilar, R., Reyes, F., Kuder, J., & Hošek, J. (Eds.). (2024). *The Reptile Database*.
<http://www.reptile-database.org>
- Vitt, L. J. (1983). Ecology of an anuran-eating guild of terrestrial tropical snakes. *Herpetologica*, 39(1), 52-66.
- Vitt, L. J., & Vangilder, L. D. (1983). Ecology of a snake community in northeastern Brazil. *Amphibia-Reptilia*, 4(2), 273-296.