


BIOTA COLOMBIANA

ISSN 0124-5376

Volumen 5 - Número 1, Junio de 2004



UNIVERSIDAD
NACIONAL
DE COLOMBIA

Instituto de Ciencias Naturales



invamar



BIOTA COLOMBIANA

Special Issue: List of Neotropical Ants


Número monográfico: *Lista de las hormigas neotropicales*

Fernando Fernández
Sebastián Sendoya

Volumen 5 - Número 1 (monográfico), Junio de 2004



Instituto de Ciencias Naturales



Synonymic list of Neotropical ants (Hymenoptera: Formicidae)

Fernando Fernández¹ and Sebastián Sendoya²

¹Profesor Asociado, Instituto de Ciencias Naturales, Facultad de Ciencias, Universidad Nacional de Colombia, AA 7495, Bogotá D.C, Colombia. *ffernandezca@unal.edu.co*

² Programa de Becas ABC, Sistema de Información en Biodiversidad y Proyecto Atlas de la Biodiversidad de Colombia, Instituto Alexander von Humboldt. *sfsendoyae@unal.edu.co*

Key words: Formicidae, Ants, Taxa list, Neotropical Region, Synopsis

Introduction

Ants are conspicuous and dominant all over the globe. Their diversity and abundance both peak in the tropical regions of the world and gradually decline towards temperate latitudes. Nonetheless, certain species such as *Formica* can be locally abundant in some temperate countries. In the tropical and subtropical regions numerous species have been described, but many more remain to be discovered. Multiple studies have shown that ants represent a high percentage of the biomass and individual count in canopy forests. Moreover, it has been established that ants seldom organize in dominant and subdominant species assemblages, a configuration that influences the structure and dynamics of not only native forests, but also agricultural systems.

About 11500 ant species of ants have been described worldwide. Of these, 3100 (30%) are present in the Neotropical Region, making it one of the richest regions worldwide in terms of ant species.

Under such perspective, the present work aims to present a synonym lists of the subfamilies, tribes, genera and species of ants present in the Neotropical Region, the geographical area confined between Mexico and Chile and Argentina (including the Caribbean). In this work we present a quick-reference alphabetical checklist of the valid names of ants available until August 2004. Readers can find additional taxonomic information in the excellent catalogue by Bolton (1995), as well as in the classic abridged catalogue by Kempf (1972), subsequently updated by Brandão (1991).

In the following paragraphs we offer a brief revision of some aspects of the biology of ants, rather as an excuse to mention some recent and interesting advances on the study of ants.

Ant Phylogeny

All ants belong to the family Formicidae, in the superfamily Vespoidea, within the order Hymenoptera. The most widely accepted phylogenetic schemes for the superfamily Vespoidea place the ants as a sister group to Vespidae + Scoliidae (Brother & Carpenter 1993; Brothers 1999). Numerous studies have demonstrated the monophyletic nature of ants (Bolton 1994, 2003; Fernández 2003). Among the most widely accepted characters used to define ants as a group are the presence of a metapleural gland in females and workers, with its reduction or secondary loss in some groups, and its complete absence of in males. Additionally, Bolton (2003) lists some apomorphies such as eusocial behavior and perennial colonies, prognathous head in females, presence of a infrabuccal pocket between the labium and the hypopharynx, geniculate antenna, second abdominal segment forming a petiole, deciduous wings in females and transverse veins 3rs-m and 2m-cu always absent.

Although the monophyletic nature of the ants remains an unquestioned issue, the relationships within the group are rather controversial. Fernández & Palacio (2003) reviewed the numerous phylogenetic proposals (mainly at the subfamily level), including examples as early as those by Brown in the 1950's up to works published in the year 2003. The work by Baroni Urbani *et al.* (1992) is probably the most thorough pertaining the internal phylogeny of ants, yet several groupings proposed in that work had weak support. Since then, even though some advances have been made (e.g. Grimaldi *et al.* 1997 and Ward & Brady 2003), the major problems remain unsolved. There is a robust body of evidence that supports the monophyly of the ants of the section doryline (Bolton 1990b; Brady 2003), but outside of

these families there is no clarity on either the limits of the groups or their relationships. Recently, Astruc *et al.* (2004) used genetic data in an attempt to resolve the problem, although their results were not quite decisive; indeed they cast doubt on the monophyletic nature of the subfamily Myrmicinae. Putative convergent evolution of morphological characters (and behavior) imposes great difficulty on the study of the internal relationships of ants; unfortunately genetic data has not solved this issue yet.

Ant Classification

We owe the first rigorous classification of the ants to William Brown Jr. As phylogenetic methods became readily available at the end of the 1950's, great efforts have been made to "polish" the taxonomy of the group; Brown (2000), Bolton (1995, 2003) and Agosti & Johnson (2003) have provided integral perspectives on this problem.

Bolton's (1995) proposal is considered the standard classification scheme; it summarizes the different advances and perspectives from the last two decades. More recently, Bolton (2003) proposed a taxonomic arrangement that in some instances significantly departs from the traditional views. Since the works by Brown in the 1950's, there has been a tendency to decrease the number of ant subfamilies, although this number slightly increased in the 1990's. Moreover, in 2003, Bolton presented a synopsis that, again, increased the number of subfamilies to 25 (of these, 21 are extant and 4 are known only from the fossil record). In the Box 1 of this work, we present a list of all the accepted subfamilies to date.

Furthermore, the situation of the classification within each subfamily is far from being stable. Although some subfamilies are quite small and do not represent any taxonomic trouble in terms of their internal classification (Leptanilloidinae, Leptanillinae, Apomyrmicinae...), some other groups are more problematic. Those ants that conform the doryline section (Aenictinae, Dorylinae, Ecitoninae...) are reasonably well studied and their delimitations and internal composition are well established (Bolton 1990a b; Brady 2003). However, one of the groups in greater need of a taxonomic solution is the "Ponerinae". This group was until recently considered a subfamily, but Bolton (2003) called its members the "poneromorphs" as an indication of the paraphyletic nature of the group. The phylogeny and internal classification of important families such as Formicinae and Myrmicinae represent cumbersome challenges for both current and future generations of myrmecologists.

In Boxes 2 and 3, we present a synopsis of the subfamilies, tribes and genera of ants of the world and the Neotropical

Region; we follow the arrangements proposed by Bolton (2003). Naturally, in some cases, the classification schemes proposed by this author should be understood as provisional. There are still many unanswered questions and in the near future someone, under the light of new evidence, might propose completely different arrangements. That is how taxonomic work goes.

Some key publications on ants

Due to their abundance and social behavior, the ants have called, in great extent, the attention of the scientific community. According the database FORMIS, there are about 30000 publications dealing directly with any aspect of ant biology. The basic literature source for any beginner in the ant world is the book by Hölldobler & Wilson (1990), which comprehensively reviews the biology of ants. The synopsis on the systematics presented in this book is somewhat out of date, but it could be supplemented with more recent references. Bourke & Franks (1995) presented a review on the evolution of ant behavior. Fernández (2003) edited a book on the Neotropical ants, which includes updated keys for the genera of the Region.

Agosti *et al.* (2000) edited a book on the capture methods and study techniques of ants dwelling in the leaf litter; some chapters go beyond and make special remarks on the usefulness of studying ants in the leaf litter as ecological and monitoring tools.

The most comprehensive catalogue of the ants in the Neotropical Region is that by Kempf (1972), which Brandão (1991) updated by subsequently. Bolton's (1995, 2003) worldwide catalogue is the most comprehensive and incorporates the most of the taxonomic changes that occurred recently in the family.

Subgenera and subspecies in ants

In the realm of ant taxonomy, there is a marked tendency to only use binomial names (i.e. genera and species only), as it is the case with the vast majority of insects, except for butterflies, on which subspecies and races are commonplace. Not until after the so-called "classic age" of myrmecology (1889-1992), ant names were depurated of supraspecific names. Before this time, ant names could be composed of up to five words, including genus, subgenus, species, subspecies, variety and even race! But since Creighton in 1930's, and William Brown in the 1950's, ant names follow the binomial system and supraspecific names either became synonyms or "good" species names. It is better to use groups of species and species complexes to divide genera in potential monophyletic units.

In the present list we not only discard the use of subgenera names, but we cite them as either real genera or likely synonyms. However, subspecies names are included since in some cases their status is not clear and in some cases they might represent valid species.

Ants in the Web

The Internet has become a powerful tool for students and researchers working on biodiversity. Certainly, ants do not escape to this trend. In fact, numerous websites such as Formis, a database with free access to near 30000 references, embody great examples on how useful web sources could be as means for diffusing knowledge. Donat Agosti has devoted significant efforts and energy to make the knowledge on ant systematics available to everyone. Agosti and Johnson have also made available several electronic tools, such as a list of publications on Hymenoptera, as well as a site with publications on pdf files dealing with the taxonomy of ants. Looking

into the future, it seems that soon we will be able to access not only primary publications, but also drawings, pictures, maps and collection data of type specimens.

The diversity of the ants in the Neotropical Region

According to Agosti (2004), there are more than 11500 valid species that belong to 21 extant subfamilies. In the Neotropical Region, little more than 3100 species, 12 subfamilies and 95 genera have been described to date. In the checklist presented herein, we included taxa whose descriptions are still in press; we omitted unpublished names and only included them for the statistics of the species and genera.

Brazil and Colombia are the countries bearing the highest species richness. This can be partially explained by size (Brazil) and complexity of the ecosystems (Colombia). However, countries such as Peru or Bolivia have been poorly studied. It can also be said that those countries with elevated figures have either been monitored better (Costa Rica) or have resident myrmecologists (Brazil, Argentina, Colombia, Venezuela).

Lista sinónímica de las hormigas Neotropicales (Hymenoptera: Formicidae)

Fernando Fernández y Sebastián Sendoya

Palabras Clave: Formicidae, Hormigas, Lista de taxones, Neotrópico, Sinopsis

Introducción

Desde casi cualquier perspectiva, se puede decir que las hormigas son insectos conspicuos y dominantes en el Mundo. Siempre se dice que las hormigas son muy abundantes en los ecosistemas tropicales, disminuyendo hacia las latitudes templadas; sin embargo en países de climas extremos pueden ser localmente abundantes (como las grandes colonias de *Formica*) y decisivas. Por supuesto, los trópicos y subtrópicos son el lugar preferido por las hormigas, con mayores números de especies, y con muchas por descubrir y describir. Muchos estudios han demostrado que estos insectos llegan a altos porcentajes en biomasa y número de individuos en las copas de los árboles de los bosques. También se ha establecido que forman mosaicos de especies dominantes y subdominantes, mosaicos que influencian la estructura y dinámica no sólo de los bosques nativos, sino de sistemas agrícolas.

Actualmente se reconocen alrededor de 11500 especies de hormigas descritas en todo el Mundo, de las cuales unas 3100 corresponden a la Región Neotropical. Con casi un 30% de las hormigas del Mundo, el Neotrópico es la región más rica en hormigas.

Esta contribución presenta listas sinónímicas de las subfamilias, tribus, géneros y especies de hormigas de la Región Neotropical, esto es, las hormigas que habitan América desde México hasta Chile y Argentina, incluyendo el Caribe. El principal objetivo es ofrecer una lista de chequeo rápida, en orden alfabético, de los nombres válidos disponibles de hormigas hasta agosto de 2004. El lector puede encontrar amplia información taxonómica en el excelente catálogo de Bolton (1995), además del clásico "catálogo abreviado" de Kempf (1972) actualizado por Brandão (1991).

Se ofrecen, en los apartados de abajo, algunas breves palabras en algunos aspectos de la biología de hormigas, especialmente como excusa para reseñar publicaciones relevantes en esos tópicos.

Filogenia de las Hormigas

Las hormigas comprenden una familia, Formicidae, dentro de la superfamilia Vespoidea, en el orden Hymenoptera. Los esquemas filogenéticos más aceptados

en Vespoidae sitúan a las hormigas como el grupo hermano de Vespidae + Scoliidae (Brother & Carpenter 1993; Brothers 1999). Numerosos estudios demuestran la monofilia de las hormigas (Bolton 1994, 2003; Fernández 2003). Un atributo universal es la presencia de glándula metapleural en hembras y obreras (reducido o perdido secundariamente en algunos pocos grupos, ausente en machos). Bolton (2003) lista estas otras apomorfias: Insectos eusociales con colonias perennes, cabeza prognata en hembras, saco infrabucal presente entre el labio y la hipofaringe, antena geniculada, segundo segmento abdominal formando un peciolo, alas de las hembras deciduas, venas transversa 3rs-m y 2m-cu siempre ausentes.

Si bien la monofilia de las hormigas no parece tener reparos o incertidumbres, otro problema es el de sus relaciones internas. Fernández & Palacio (2003) revisan las numerosas propuestas de filogenia de las grandes categorías (principalmente subfamilias) desde los primeros escritos de Brown (hacia 1950) hasta el 2003. Baroni Urbani et al. (1992) realizaron uno de los estudios más completos en filogenia interna de las hormigas; sin embargo varias de sus agrupaciones son débiles y la consistencia general de sus árboles es baja. Desde entonces se han realizado algunos ajustes (Grimaldi et al. 1997; Ward & Brady 2003) pero sin resolver estos grandes problemas. Hay evidencias robustas de la monofilia de las hormigas llamadas de la sección dorylina (Bolton 1990b; Brady 2003), pero por fuera de estas subfamilias no hay claridad sobre sus límites y relaciones. Recientemente Astruc et al. (2004) publican la primera gran aproximación a la filogenia de las hormigas, pero sus resultados son poco decisivos, e incluso el uso de genes pone una sombra de duda sobre la monofilia de Myrmicinae.

La posible evolución convergente de muchos rasgos morfológicos (y de comportamiento) dificultan notoriamente cualquier intento de resolver la filogenia interna de las hormigas. A esto no escapan tampoco los genes.

Clasificación de las Hormigas

Nuevamente debemos a William Brown Jr. los primeros intentos de ofrecer una clasificación coherente de las hormigas. A partir de la década de los 50 se hacen cada vez mejores esfuerzos por "limpiar" la taxonomía del grupo y ofrecer agrupaciones acordes con los estudios filogenéticos en las hormigas. Brown (2000), Bolton (1995, 2003) y Agosti & Johnson (2003) ofrecen una visión más integral de los aportes en taxonomía de las hormigas.

La propuesta de Bolton en 1995 puede tomarse como la estándar en el arreglo de la clasificación de las hormigas en las últimas dos décadas del siglo XX. Recientemente

Bolton (2003) ha ofrecido un nuevo arreglo que, en algunos casos, se aleja bastante de las clasificaciones "tradicionales". A partir del trabajo de Brown una tendencia fue la de disminuir el número de subfamilias, número que aumentó ligeramente en los 90. La sinopsis de Bolton (2003) vuelve a aumentar notoriamente el número de subfamilias a 25, 21 de ellas vivientes y las restantes cuatro conocidas de sólo fósiles. El Cuadro 1 ofrece la lista de todas las subfamilias aceptadas hasta la fecha.

La situación dentro de cada subfamilia aún dista de ser estable. Algunas subfamilias son muy pequeñas y en principio no hay mayores conflictos en sus arreglos internos (Leptanilloidinae, Leptanillinae, Apomyrminae...). Aquellas que conforma la sección dorylina (Aenictinae, Dorylinae, Ecitoninae...) están razonablemente estudiadas en sus límites y composición interna (Bolton 1990ab; Brady 2003). Uno de los grupos más importantes por resolver es la composición de "Ponerinae", grupo hasta hace poco considerado una subfamilia pero que Bolton (2003) denomina "poneromorfos" como una agrupación parafilética. La filogenia y clasificación interna de subfamilias tan importantes como Formicinae y, sobre todo, Myrmicinae, son enormes retos para los actuales y futuros mirmecólogos.

Los cuadros 2 y 3 ofrecen sinopsis de las subfamilias, tribus y géneros de hormigas en el Mundo y en la Región Neotropical, siguiendo el arreglo de Bolton (2003). Naturalmente, la propuesta de este autor debe tomarse como provisional en algunos casos. Todavía hay muchas preguntas por resolver, y en algunos pocos años alguien podrá ofrecer nuevos arreglos.

Algunas publicaciones claves sobre hormigas

Debido a su abundancia y comportamiento social, las hormigas han despertado un enorme interés en la ciencia. Según la base de datos FORMIS, hay alrededor de 30000 publicaciones directas sobre cualquier aspecto sobre hormigas. El libro base para cualquiera interesado en hormigas es el de Hölldobler & Wilson (1990), el cual es una gran síntesis en biología de este grupo. La sinopsis sistemática y claves presentan desfases que se pueden obviar con referencias más recientes. Bourke & Franks (1995) presentan una revisión sobre la evolución del comportamiento en estos himenópteros. Fernández (2003) edita un libro sobre hormigas neotropicales, el cual incluye claves actualizadas para los géneros de la región.

Agosti et al. (2000) editan un libro clave sobre métodos de captura y estudio de hormigas de hojarasca; algunos capítulos van más allá de las hormigas de ese sustrato y tienen gran importancia para quien piense en usar hormigas en monitoreos y estudios ecológicos.

Para la Región Neotropical el catálogo “clásico” es el de Kempf (1972) actualizado por Brandão en 1991. Bolton (1995, 2003) es indispensable para el catálogo del Mundo y para los varios cambios taxonómicos en la familia.

Subgéneros y subespecies en hormigas

Existe la tendencia a usar sólo nombres binominales para hormigas (esto es género y especie), así como para la mayoría de insectos con la notable excepción de las mariposas. Las hormigas han sido uno de los grupos más “castigados” con el uso y abuso de los nombres inferiores a especie. En los años de la edad clásica de la mirmecología (1889 a 1922) se proponían hasta cinco nombres para una hormiga, con el género, subgénero, especie, subespecie, variedad y hasta raza! Desde Creighton en 1930 y sobretodo William Brown en los años 50, las hormigas se nombran sólo con género y especie. Esto significa que subespecies y demás artificios han pasado a sinonimia menor o a buenas especies. Es mejor usar grupos de especies y complejos de especies para dividir géneros en unidades potencialmente monofiléticas.

En la presente lista no se usarán los subgéneros, y se citarán sólo como sinónimos reales o probables en donde corresponda. Las subespecies se incluyen debido a que en muchos casos su estatus no está claro, y varios de estos nombres pueden ser especies válidas.

Hormigas en la Web

La internet se ha vuelto una herramienta poderosa para los estudiosos de la Biodiversidad. A esto no escapan las hormigas; de hecho, estos insectos se han convertido en

un ejemplo de las bondades de la web para difundir conocimientos. FORMIS es un excelente ejemplo: una base de datos de libre acceso y uso para casi 30000 referencias. Donat Agosti ha dedicado muchos esfuerzos y energías por hacer democrática la información sobre sistemática de hormigas. Agosti y Johnson han puesto varias ayudas electrónicas, como la lista de publicaciones en sistemática de Hymenoptera, publicaciones para descargar en pdf en taxonomía de hormigas (donde no se apliquen restricciones de copyright). Mirando hacia el futuro cercano, podremos tener acceso no sólo a las publicaciones primarias, sino dibujos, fotos, mapas y datos de los tipos.

La diversidad de hormigas en la Región Neotropical

De acuerdo a Agosti (2004) hay algo más de 11500 especies de hormigas descritas en 21 subfamilias vivientes. Para la Región Neotropical, se han descrito un poco más de 3100 especies en 12 subfamilias y 95 géneros. Este número incluye especies cuyas descripciones están en prensa o próximas a publicarse. La lista, por supuesto, no incluye nombres no publicados, por lo que es inferior a 3100; sin embargo, en la sinopsis de géneros y en las estadísticas estas especies si se incluyen.

Brasil y Colombia son los países más ricos en especies. Esto puede relacionarse con el tamaño (Brasil) o complejidad de ecosistemas (Colombia). Sin embargo, hay países como Perú o Bolivia muy pobemente estudiados. También puede decirse que los países con mejores cifras han sido mejor inventariados (como Costa Rica) o tienen mirmecólogos residentes (Brasil, Argentina, Colombia, Venezuela).

Box 1. Extant and extinct subfamilies of ants.

Cuadro 1. Subfamilias vivientes y fósiles de hormigas del Mundo.

Formicomorph subfamilies 1 Aneuretinae 2 Dolichoderinae 3 Formicinae Myrmeciomorph subfamilies 4 Myrmeciinae 5 Pseudomyrmecinae	Dorylomorph subfamilies 6 Aenictinae 7 Aenictogitoninae 8 Dorylinae 9 Cerapachyinae 10 Ecitoninae 11 Leptanilloidinae
-------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------

Leptanilimorph subfamilies	Myrmicomorph subfamilies
12 Apomyrminae	20 Agroecomyrmecinae
13 Leptanillinae	21 Myrmicinae
Poneromorph subfamilies	Extinct subfamilies
14 Amblyoponinae	22 Armaniinae
15 Ectatomminae	23 Sphecomyrmecinae
16 Heteroponerinae	24 Brownimicinae
17 Paraponerinae	25 Formiciinae
18 Ponerinae	
19 Proceratiinae	

Box 2. List of subfamilies and tribes of the ants of the World. Groups present in the Neotropical Region are displayed in **boldface**.

Cuadro 2. Lista de subfamilias y tribus de las hormigas del mundo. Aquellos grupos presentes en el Neotrópico se muestran en **negrita**.

Aenictinae:	Aenictini
Aenictogitoninae:	Aenictogitonini
Agroecomyrmecinae:	Agroecomyrmecini
Amblyoponinae:	Amblyoponini
Aneuretinae:	Aneuretini
Apomyrminae:	Apomyrmini
Cerapachyinae:	Acantostichini, Cerapachyini, Cylindromyrmecini
Dorylinae:	Dorylini
Dolichoderinae:	Dolichoderini
Ectitoninae:	Cheliomyrmecini, Ectonini
Ectatomminae:	Ectatommini, Typhlomyrmecini
Formiciniae:	Camponotini, Formicini, Gesomyrmecini, Giantiopini, Lasiini, Melophorini, Myrmecorhynchini, Notostigmatini, Oecophyllini, Plagiolepidini, Mymoteratini
Heteroponerinae:	Heteroponerini
Leptanillinae:	Anomalomyrmini, Leptanillini
Leptanilloidinae:	Leptanilloidini
Myrmeciinae:	Myrmeciini, Prionomyrmecini
Myrmicinae:	Adelomyrmecini, Ankylomyrmini, Attini, Basicerotini, Blepharidattini, Cataulacini, Cephalotini, Crematogastrini, Dacetini, Formicoxenini, Lenomyrmecini, Liomyrmecini, Melissotarsini, Meranoplini, Metaponini, Myrmecinini, Myrmicariini, Myrmicini, Paratopulini, Phalacromyrmecini, Pheidolini, Solenopsidini, Stegomyrmecini, Stenammini, Tetramoriini
Paraponerinae:	Paraponerini

Ponerinae:	Platythyreini, Ponerini, Thaumatomymecini
Proceratiinae:	Probolomyrmecini, Proceratiini
Pseudomyrmecinae:	Pseudomyrmecini
Armaniinae*	Armaniini*
Brownimecinae*	Brownimeciini*
Formiciinae*	Formiciini*
Sphecomyrminae*	Haidomyrmecini*, Sphecomyrmini*

Box 3. Synopsis of the subfamilies, tribes and genera of extant ants in the neotropics. ad: Dominican Amber.

Cuadro 3. Sinopsis de subfamilias, Tribes y géneros de las hormigas existentes en el neotropico. ad: Ambar dominicano.

Taxon Taxón	Species Number / Número de especies		Neotropical distribution / Distribución en el neotrópico	References / Referencias
	Neotropics /Neotrópico	Colombia / Colombia		
Family Formicidae Latreille, 1809			Neotropical	Brown 1973 2000; Wheeler & Wheeler 1976; Kempf 1972; Brandão 1991; Baroni Urbani <i>et al.</i> 1992; Bolton 1994 1995ab 2003; Longino & Hanson 1995; Fernández 2003c; Kusnezov 1963 1978; Jaffé <i>et al.</i> 1993
Subfamily Dolichoderinae Forel, 1878	221	38	Neotropical	Shattuck 1992ab 1994 1995; Brandão <i>et al.</i> 1998; Chiotis <i>et al.</i> 2000 ; Bolton 2003
Tribe Dolichoderini Forel, 1878	221	38	Neotropical	Shattuck 1992ab 1994 1995; Brandão <i>et al.</i> 1998; Bolton 2003
<i>Anillidris</i> Santschi, 1936	1	0	ar	Shattuck 1992a 1994
<i>Azteca</i> Forel, 1878	72	15	Neotropical	Emery 1893; Shattuck 1992a 1994; Longino 1989 1991 1996; Ayala <i>et al.</i> 1996
<i>Bothriomyrmex</i> Emery, 1869	1	0	cr	Shattuck 1992a 1994; Longino Ants of Costa Rica Web Page
<i>Dolichoderus</i> Lund, 1831	60	15	Neotropical	Shattuck 1992a 1994; MacKay 1993
<i>Dorymyrmex</i> Mayr, 1866	41	3	Neotropical	Shattuck 1992a 1994; Kusnezov 1952c; Snelling 1973 1995
<i>Forelius</i> Emery, 1888	17	0	ar pr ur bo br me us am	Shattuck 1992a 1994; Cuezzo 2000
<i>Leptomyrmex</i> Mayr, 1862	1	0	ad	Shattuck 1992a 1994
<i>Linepithema</i> Mayr, 1866	14	4	Neotropical	Shattuck 1992a 1994
<i>Tapinoma</i> Foerster, 1850	11	1	Neotropical	Shattuck 1992a 1994
<i>Technomyrmex</i> Mayr, 1872	3	0	pn ad	Shattuck 1992a 1994; Brandão <i>et al.</i> 1998
Subfamily Formicinae Latreille, 1809	407	60	Neotropical	Agosti 1991; Bolton 2003

Taxon Taxón	Species Number / Número de especies		Neotropical distribution / Distribución en el neotrópico	References / Referencias
	Neotropics / Neotrópico	Colombia / Colombia		
Tribe Camponotini Forel, 1878	227	41	Neotropical	Bolton 2003
<i>Camponotus</i> Mayr, 1861	227	41	Neotropical	Kusnezov 1952b; Hashmi 1973; MacKay 1997 2005; MacKay & Mackay 1997a; Snelling & Torres 1998; Brady <i>et al.</i> 2000; Fernández 2002a
Tribe Gigantiopini Ashmead, 1905	1	1	pe br gi gf su vn co	Bolton 2003
<i>Gigantiops</i> Roger, 1863	1	1	pe br gi gf su vn co	Kempf & Lenko 1968
Tribe Formicini Latreille, 1809	10	0	us	Agosti 1994; Bolton 2003
<i>Formica</i> Linnaeus, 1758	9	0	us	Wheeler W.M. 1913
<i>Polyergus</i> Latreille, 1804	1	0	us	
Tribe Lasiini Ashmead, 1905	101	7	Neotropical	Agosti 1991; Bolton 2003
<i>Acanthomyops</i> Mayr, 1862	2	0	me us	
<i>Acropyga</i> Roger, 1862	27	3	Neotropical	LaPolla 2004
<i>Anoplolepis</i> Santschi, 1914	2	0	ch me	
<i>Lasiophanes</i> Emery, 1895	5	0	ar ch	Kusnezov 1952a
<i>Lasius</i> Fabricius, 1804	3	0	us	Wilson 1955
<i>Myrmecocystus</i> Wesmael, 1838	17	0	me us	Snelling 1976 1982
<i>Myrmelachista</i> Roger, 1863	45	4	Neotropical	
Tribe Plagiolepidini Forel, 1886	68	11	Neotropical	Agosti 1991; Bolton 2003
<i>Brachymyrmex</i> Mayr, 1868	35	7	Neotropical	Santschi 1923
<i>Plagiolepis</i> Emery 1894	1	0	ber	
<i>Paratrechina</i> Motschoulsky, 1863	30	4	Neotropical	Trager 1984
<i>Plagiolepis</i> Mayr, 1861	2	0	me us	
Subfamily Myrmeciinae Emery, 1877	2	0	ar	Baroni Urbani 2000; Bolton 2003; Ward & Brady 2003
Insertae sedis:				
<i>Ameghinoia*</i> Viana & Haedo Rossi 1957	1	0	ar	Ward & Brady 2003
<i>Polanskiella*</i> Rossi de García, 1983	1	0		Ward & Brady 2003
Subfamily Pseudomyrmecinae M.R. Smith, 1952	139	32	Neotropical	Ward 1985 1989 1990 1991 1992 1993 1996 1999 2002; Bolton 2003
Tribe Pseudomyrmecini M.R. Smith, 1952	139	32	Neotropical	Ward 1990
<i>Myrcidris</i> Ward, 1990	2	0	br su	Ward 1990
<i>Pseudomyrmex</i> Lund, 1831	137	32	Neotropical	Ward 1989 1991 1992 1993 1996 1999
Subfamily Cerapachyinae Forel, 1893	39	6	Neotropical	Brown 1975; Bolton 1990a 2003
Tribe Acanthostichini Emery, 1901	20	2	Neotropical	Bolton 1990a 2003
<i>Acanthostichus</i> Mayr, 1887	20	2	Neotropical	Bolton 1990a; MacKay 1996 2003
Tribe Cerapachyini Forel, 1893	6	0	Neotropical	Bolton 1990a 2003
<i>Cerapachys</i> F. Smith, 1857	5	0	Neotropical	Bolton 1990a
<i>Sphinctomyrmex</i> Mayr, 1866	1	0	br	Bolton 1990a
Tribe Cylindromyrmecini Emery, 1901	13	4	Neotropical	Bolton 1990a 2003
<i>Cylindromyrmex</i> Mayr, 1870	13	4	Neotropical	Bolton 1990a; Andrade 1998 2001
Subfamily Ecitoninae Forel, 1893	144	18	Neotropical	Borgmeier 1955; Watkins 1976; Gotwald 1982 1995; Palacio 1999; Bolton 1990b 2003
Tribe Cheliomyrmecini Wheeler, 1921	4	2	Neotropical	Bolton 1990b 2003
<i>Cheliomyrmex</i> Mayr, 1870	4	2	Neotropical	Watkins 1976; Palacio 1999

Taxon Taxón	Species Number / Número de especies		Neotropical distribution / Distribución en el neotrópico	References / Referencias
	Neotropics /Neotrópico	Colombia / Colombia		
Tribe Ecitonini Forel, 1893	140	16	Neotropical	Bolton 1990b 2003
<i>Ectyon</i> Latreille, 1804	12	5	Neotropical	Watkins 1976; Palacio 1999
<i>Labidus</i> Jurine, 1807	8	3	Neotropical	Watkins 1976; Palacio 1999
<i>Neivamyrmex</i> Borgmeier, 1940	118	7	Neotropical	Watkins 1976; Palacio 1999
<i>Nomamyrmex</i> Borgmeier, 1936	2	1	Neotropical	Watkins 1976; Palacio 1999
Subfamily Leptanilloidinae Bolton, 1992	8	2	Neotropical	Bolton 1990ab 2003; Brandão <i>et al.</i> 1999
Tribe Leptanilloidini Bolton, 1992	8	2	Neotropical	Brandão <i>et al.</i> 1999
<i>Asphinctanilloides</i> Brandão, Diniz, Agosti & Delabie, 1999	3	0	br	Brandão <i>et al.</i> 1999
<i>Leptanilloides</i> Mann, 1923	5	2	bo ec co cr	Brandão <i>et al.</i> 1999; Longino 2004
Subfamily Amblyoponinae Forel, 1893	22	7	Neotropical	Brown 1960; Bolton 2003
Tribe Amblyoponini Forel 1893	22	7	Neotropical	Brown 1960; Bolton 2003
<i>Amblyopone</i> Erichson, 1842	16	5	Neotropical	Brown 1960; Lattke 1991; Lacau & Delabie 2002
<i>Prionopelta</i> Mayr, 1866	5	2	Neotropical	Brown 1960
<i>Insertae sedis:</i>				
<i>Paraprisopelta</i> Kusnezov, 1955	1	0	ar	Brown 1960
Subfamily Ponerinae Lepeletier, 1835	216	44	Neotropical	Bolton 1990b 2003; Ward 1994
Tribe Ponerini Lepeletier, 1835	202	42	Neotropical	Bolton & Brown 2002; Bolton 2003
<i>Anochetus</i> Mayr, 1861	30	7	Neotropical	Brown 1978
<i>Belonopelta</i> Mayr, 1870	2	1	co ho me	Baroni Urbani 1975
<i>Centromyrmex</i> Mayr, 1866	3	0	ar bo br cr	Kempf 1967
<i>Cryptopone</i> Emery, 1893	1	0	ni gu	
<i>Dinoponera</i> Roger, 1861	6	0	ar pr ur bo pe br	Kempf 1971
<i>Hypoponera</i> Santschi, 1938	36	3	Neotropical	
<i>Leptogenys</i> Roger, 1861	38	4	Neotropical	Lattke 2005
<i>Odontomachus</i> Latreille, 1804	25	8	Neotropical	Brown 1976
<i>Pachycondyla</i> F. Smith, 1858	51	19	Neotropical	Wild 2002
<i>Simopelta</i> Mann, 1922	10	0	br	Gotwald & Brown 1966
Tribe Platythyreini Emery, 1901	8	2	Neotropical	Brown 1975; Bolton 2003
<i>Platythyrea</i> Roger, 1863	8	2	Neotropical	Brown 1952 1975; Kugler 1977
Tribe Thaumatomymecini Emery, 1901	6	0	br gi ho am	Bolton 2003
<i>Taumatomymex</i> Mayr, 1887	6	0	br gi ho am	Kempf 1975; Longino 1988
Subfamily Ectatomminae Emery, 1895	102	24	Neotropical	Brown 1958; Lattke 1994; Keller 2000; Bolton 2003
Tribe Ectatommini Emery 1895	95	23	Neotropical	Brown 1958; Lattke 1994; Keller 2000; Bolton 2003
<i>Ectatomma</i> F. Smith, 1858	13	13	Neotropical	Kugler & Brown 1982
<i>Gnamptogenys</i> Roger, 1863	82	10	Neotropical	Brown 1958 1992; Lattke 1990a 1995 2002 2003
Tribe Typhlomyrmecini Emery, 1911	7	1	Neotropical	Brown 1965
<i>Typhlomyrmex</i> Mayr, 1862	7	1	Neotropical	Brown 1965

Taxon Taxón	Species Number / Número de especies		Neotropical distribution / Distribución en el neotrópico	References /Referencias
	Neotropics /Neotrópico	Colombia / Colombia		
Subfamily Heteroponerinae Bolton 2003	17	4	Neotropical	Bolton 2003
Tribe Heteroponerini Bolton 2003	17	4	Neotropical	Bolton 2003
<i>Acanthoponera</i> Mayr, 1862	4	1	Neotropical	Brown 1958
<i>Heteroponera</i> Mayr, 1887	13	3	Neotropical	Kempf 1962
Subfamily Paraponerinae Paraponerinae Emery, 1901	2	1	Neotropical	Brown 1958; Bolton 2003
Tribe Paraponerini Emery, 1901	2	1	Neotropical	Brown 1958; Lattke 1994; Keller 2000; Bolton 2003
<i>Paraponera</i> F. Smith, 1858	2	1	Neotropical	Brown 1958
Subfamily Proceratiinae Emery, 1895	18	2	Neotropical	Lattke 1994; Keller 2000; Bolton 2003
Tribe Proceratiini Emery, 1895	14	0	Neotropical	Lattke 1994; Keller 2000; Bolton 2003
<i>Discothyrea</i> Roger, 1863	8	0	Neotropical	Borgmeier 1949
<i>Proceratium</i> Roger, 1863	6	0	Neotropical	Brown 1979; Ward 1988; Baroni Urbani & de Andrade 2003
Tribe Probolomyrmecini, Perrault 2000	4	2	ar bo pe vn co pn cr	Lattke 1994; Keller 2000; Perrault 2000; Bolton 2003
<i>Probolomyrmex</i> Mayr, 1901	4	2	ar bo pe vn co pn cr	Agosti 1995; O'Keeffe & Agosti 1997
Subfamily Agroecomyrmecinae Carpenter, 1930	1	0	es	Bolton 2003
Tribe Agroecomyrmecini Carpenter, 1930	1	0	es	Brown & Kempf 1967; Bolton 2003
<i>Tatuidris</i> Brown & Kempf, 1968	1	0	es	Brown & Kempf, 1967
Subfamily Myrmicinae Lepeletier, 1835	1540	315	Neotropical	Bolton 2003
Tribe Adelomyrmecinini Fernandez, 2004	20	6	Neotropical	Fernández 2004b
<i>Adelomyrmex</i> Emery, 1897	18	6	Neotropical	Fernández 2003b 2004b
<i>Cryptomyrmex</i> Fernández, 2004	2	0	br pa	Fernández 2004b
Tribe Attini F. Smith, 1858	216	32	Neotropical	Weber 1972 1982; Schultz & Meier 1994; Wetterer et al 1998; Müller et al. 1998 2001; Bolton 2003
<i>Acromyrmex</i> Mayr, 1865	25	6	Neotropical	Santschi 1925; Goncalves 1961; Fowler 1988
<i>Apterostigma</i> Mayr, 1865	48	4	Neotropical	Lattke 1997 1999
<i>Atta</i> Fabricius, 1804	10	4	Neotropical	Borgmeier 1959; MacKay & MacKay 1986
<i>Cyphomyrmex</i> Mayr, 1862	38	10	Neotropical	Kempf 1964b 1966; MacKay & Baena 1993; Snelling & Longino 2002
<i>Mycetagoicus</i> Brandão & Mayhe-Nunes, 2001	3	0	br	Brandão y Mayhé-Nunes, 2001
<i>Mycetarotes</i> Emery, 1913	2	0	br	Kempf 1961; Mayhe-Nunes 1995
<i>Mycetophylax</i> Emery, 1913	7	1	Suramerica	
<i>Mycetosoritis</i> Wheeler, 1907	4	0	ar pr br	
<i>Mycocerpus</i> Forel, 1893	5	1	Neotropical	MacKay 1998
<i>Myrmicocrypta</i> F. Smith, 1860	24	0	Neotropical	Kempf 1963; MacKay 1998
<i>Pseudoatta</i> Gallardo, 1916	1	0	ar	
<i>Sericomyrmex</i> Mayr, 1865	19	1	Neotropical	
<i>Trachymyrmex</i> Forel, 1893	30	5	Neotropical	MacKay & MacKay 1997b
Tribe Basicerotini Brown, 1949	35	8	Neotropical	Brown & Kempf 1960; Bolton 2003

Taxon Taxón	Species Number / Número de especies		Neotropical distribution / Distribución en el neotrópico	References / Referencias
	Neotropics /Neotrópico	Colombia / Colombia		
<i>Basiceros</i> Schulz, 1906	4	0	br gi gf su cr ni ho an	Brown & Kempf 1960; Brown 1974
<i>Eurhopalothrix</i> Brown & Kempf, 1961	10	0	Neotropical	Brown & Kempf 1960
<i>Octostruma</i> Forel, 1912	11	6	Neotropical	Brown & Kempf 1960; Palacio 1997
<i>Protalaridris</i> Brown, 1980	1	1	ec co	Brown 1980
<i>Rhopalothrix</i> Mayr, 1870	8	1	ar br gi co pn me am	Brown & Kempf 1960
<i>Talaridris</i> Weber, 1941	1	0	an	Brown & Kempf 1960
Tribe Blepharidattini Wheeler & Wheeler, 1991	12	1	Neotropical	Schultz & Meier 1995; Bolton 2003
<i>Blepharidatta</i> Wheeler, 1915	2	0	ar br	
<i>Wasmannia</i> Forel, 1893	10	1	Neotropical	Longino & Fernández 2004
Tribe Cephalotini M.R. Smith, 1949	173	48	Neotropical	Kempf 1951 1958 1964a 1973b; Andrade & Baroni Urbani 1999
<i>Cephalotes</i> Latreille, 1802	131	36	Neotropical	Andrade & Baroni Urbani 1999
<i>Procryptocerus</i> Emery, 1887	42	12	Neotropical	Longino & Snelling 2002
Tribe Crematogastrini Forel, 1893	103	28	Neotropical	Bolton 2003
<i>Crematogaster</i> Lund, 1831	103	28	Neotropical	Longino 2003a
Tribe Dacetini Forel, 1892	191	55	Neotropical	Baroni Urbani & de Andrade 1994; Bolton 1998 1999 2000
<i>Acanthognathus</i> Mayr, 1887	7	2	Neotropical	Brown & Kempf 1969; Bolton 2000
<i>Dacetum</i> Perty, 1833	1	0	bo pe br gi gf su	Bolton 2000
<i>Pyramica</i> Roger, 1862	99	27	Neotropical	Bolton 2000
<i>Strumigenys</i> F. Smith, 1860	84	26	Neotropical	Lattke & Goitia 1997; Bolton 2000
Tribe Formicoxenini Forel, 1893	39	7	Neotropical	Bolton 2003
<i>Cardiocondyla</i> Emery, 1869	3	0	br me us am an bh	MacKay 1995
<i>Nesomyrmex</i> Wheeler W.M., 1910	21	5	Neotropical	Kempf 1959
<i>Ochetomyrmex</i> Mayr, 1878	2	2	Suramerica	Fernández 2003a
<i>Temnothorax</i> Mayr 1861	9	0	gu me us	Baroni Urbani 1978; Fontenla 2000; Bolton 2003
<i>Xenomyrmex</i> Forel, 1885	4	0	pn gu me us am bh	Creighton 1957
Tribe Lenomyrmicini Bolton, 2003	5	3	ec co pn	Bolton 2003
<i>Lenomyrmex</i> Fernández & Palacio, 1999	5	3	ec co pn	Fernández & Palacio 1999; Fernández 2001
Tribe Myrmecinini Ashmead, 1905	2	0	gu us	Moffet 1986; Okido & Ogata 2000
<i>Myrmecina</i> Curtis, 1829	1	0	us	
<i>Perissomyrmex</i> M.R. Smith, 1947	1	0	gu	Longino & Hartley 1994
Tribe Myrmicini Lepeletier, 1835	47	4	Neotropical	Bolton 2003
<i>Hylomyrma</i> Forel, 1912	6	2	Neotropical	Kempf 1973a
<i>Pogonomyrmex</i> Mayr, 1868	41	2	Neotropical	Lattke 1990b; Fernández & Palacio 1998; Taber 1998
Tribe Phalacromyrmecini Phalacromyrmecini Dlussky & Fedoseeva, 1988	1	0	br	Kempf 1960; Bolton 1998 2003
<i>Phalacromyrmex</i> Kempf, 1960	1	0	br	Kempf 1960
Tribe Pheidolini Emery, 1877	473	88	Neotropical	Bolton 2003
<i>Aphaenogaster</i> Mayr, 1853	5	0	pn cr ho me co	
<i>Messor</i> Forel, 1890	4	0	me us	
<i>Phiedole</i> Westwood, 1839	464	88	Neotropical	Brown 1981; Wilson 2003
Tribe Solenopsidini Forel, 1893	181	31	Neotropical	Kusnezov 1957; Ettershank 1966; Bolton 1987 2003
<i>Allomerus</i> Mayr, 1878	2	1	bo br gi	
<i>Carebara</i> Westwood, 1840	30	11	ar pr pe br gi gf su pn me am an	Fernández 2004a

Taxon Taxón	Species Number / Número de especies		Neotropical distribution / Distribución en el neotrópico	References /Referencias
	Neotropics /Neotrópico	Colombia / Colombia		
<i>Megalomyrmex</i> Forel, 1885	23	9	Neotropical	Ettershank 1966; Bolton 1987; Brandão 1990 2003
<i>Monomorium</i> Mayr, 1855	17	3	Neotropical	Ettershank 1966; DuBois 1986; Bolton 1987; Heterick 2001; Fernández 2004c
<i>Oxyepoecus</i> Santschi, 1926	9	0	ar pr bo br	Kempf 1974; Alburquerque & Brandão 2003
<i>Solenopsis</i> Westwood, 1840	97	6	Neotropical	Creighton 1930; Ettershank 1966; Bolton 1987; Trager 1991; Taber 2000
<i>Tranopelta</i> Mayr, 1866	2	1	bo ec br gi gf su vn co pn cr an	Fernández 2003a
<i>Insertae sedis</i>	1	0		
<i>Oxyidris*</i> Wilson 1985	1	0	ad	
Tribe Stegomyrmecini Wheeler, 1922	2	0	bo pe pn	Diniz 1990
<i>Stegomyrmex</i> Emery, 1912	2	0	bo pe pn	Diniz 1990
Tribe Stenammini Ashmead, 1905	36	2	Neotropical	Bolton 2003
<i>Bariamyrmra</i> Lattke, 1990	1	0	vn	Lattke 1990b
<i>Lachnomyrmex</i> Wheeler, 1910	4	0	br pn gu me an	Smith M.R. 1944; Fernández & Baena 1997
<i>Rogeria</i> Emery, 1894	21	1	ar br gi gf su co cr ho am an	Kugler 1994
<i>Stenamma</i> Westwood, 1839	9	1	co cr ho me us	Smith M.R. 1962
<i>Insertae sedis:</i>				
<i>Hemomyrmex*</i> Wilson 1985	1	0	ad	
Tribe Tetramoriini Emery, 1895	4	2	Neotropical	Bolton 1976 1979
<i>Tetramorium</i> Mayr, 1855	4	2	Neotropical	Bolton 1979
<i>Insertae sedis</i> en Formicidae				
Subfamily* Paleosminthurinae Pierce & Gibron 1962	1	0	br	
<i>Cariridris*</i> Brandão & Martins-Nieto 1990	1	0	br	

Taxonomic List / Listado Taxonómico

List of ants species present from the south of the United States (U.S.) to Patagonia (ar and ch) through the Caribbean islands. The classification scheme supraspecies used is the proposed one by Bolton (2003). The geographic distribution of each one of the species is based on I catalogue of Kempf (1972), in the adendos to this I catalogue by Brandão (1991) and in the descriptions of species and later generic revisions. The bibliographical references include single those taxonomically excellent publications for each species, for information but detailed to consult I catalogue of Bolton 1995, (*): Species reported single like fossils.

Lista de especies de hormigas presentes en la región, desde el sur de los Estados Unidos (us) hasta la Patagonia (ar y ch) pasando por las islas del caribe. El esquema de clasificación supraespecífica utilizado es el propuesto por Bolton (2003). La distribución geográfica de cada una de las especies se basa en el catálogo de Kempf (1972), en los adendos a este catálogo por Brandão (1991) y en las descripciones de especies y revisiones genéricas posteriores. Las referencias bibliográficas incluyen solo aquellas publicaciones taxonómicamente relevantes para cada especie, para información mas detallada consultar el catálogo de Bolton 1995. (): Especies reportadas solo como fósiles.*

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
Subfamily Dolichoderinae		
Tribe Dolichoderini		
<i>Anillidris bruchi</i> Santschi, 1936	ar	Shattuck 1992a 1994
<i>Azteca adrepens</i> Forel, 1911	pr	Shattuck 1994
<i>Azteca aesopus</i> Forel, 1908	br	Shattuck 1994
<i>Azteca alfari</i> Emery, 1893	ar pr bo pe br gi gf su tt vn co pn cr ni be ho gu me am	Longino 1989 1991; Shattuck 1994
<i>Azteca alpha*</i> Wilson, 1985	br	Wilson 1985; Shattuck 1994
<i>Azteca angusticeps</i> Emery, 1893	vn	Emery 1893; Shattuck 1994
<i>Azteca aragua</i> Longino, 1991	bo br co pn cr	Longino 1991; Shattuck 1994
<i>Azteca aurita</i> Emery, 1893	bo pe	Emery 1893; Shattuck 1994
<i>Azteca australis</i> Wheeler W. M., 1942	br tt	Longino 1991; Shattuck 1994
<i>Azteca barbifex</i> Forel, 1906	br	Shattuck 1994
<i>Azteca bequaerti</i> Wheeler W. M. & Bequaert, 1929	br	Shattuck 1994
<i>Azteca brevicornis</i> (Mayr, 1878)	br	Emery 1893; Shattuck 1994
<i>Azteca brevicornis boliviiana</i> Wheeler W. M., 1942	bo	Emery 1893; Shattuck 1994
<i>Azteca brevis</i> Forel, 1899	cr	Shattuck 1994
<i>Azteca chartifex</i> Emery, 1896	br tt	Shattuck 1994
<i>Azteca chartifex cearensis</i> Forel, 1903	br	Shattuck 1994
<i>Azteca chartifex decipiens</i> Forel, 1906	br	Shattuck 1994
<i>Azteca chartifex lanians</i> Emery, 1912	br	Shattuck 1994
<i>Azteca chartifex laticeps</i> Forel, 1899	br gi gf su pn cr	Shattuck 1994
<i>Azteca chartifex multinida</i> Forel, 1899	co	Emery 1893; Shattuck 1994
<i>Azteca chartifex spiriti</i> Forel, 1912	br	Shattuck 1994
<i>Azteca chartifex stalactitica</i> Emery, 1896	br	Shattuck 1994
<i>Azteca christopherseni</i> Forel, 1912	pn	Shattuck 1994
<i>Azteca coeruleipennis</i> Emery, 1893	tt pn cr me	Longino 1991; Shattuck 1994
<i>Azteca constructor</i> Emery, 1896	ec gi gf su tt pn cr gu	Longino 1991; Shattuck 1994
<i>Azteca coussapoae</i> Forel, 1904	br	Emery 1893; Shattuck 1994
<i>Azteca crassicornis</i> Emery, 1893	br	Emery 1893; Shattuck 1994
<i>Azteca delpini</i> Emery, 1893	br gi gf su gu	Emery 1893; Shattuck 1994
<i>Azteca delpini antillana</i> Forel, 1899	co an	Shattuck 1994
<i>Azteca delpini trinidadensis</i> Forel, 1899	tt	Shattuck 1994
<i>Azteca depilis</i> Emery, 1893	br	Shattuck 1994
<i>Azteca duckei</i> Forel, 1906	br	Shattuck 1994
<i>Azteca duroiae</i> Forel, 1904	br	Shattuck 1994
<i>Azteca emeryi</i> Forel, 1904	br	Shattuck 1994
<i>Azteca eumeceus*</i> Wilson, 1985	am	Wilson 1985; Shattuck 1994
<i>Azteca fasciata</i> Emery, 1893	bo br	Emery 1893; Shattuck 1994
<i>Azteca fasciata laeta</i> Wheeler W.M., 1942	pn	Shattuck 1994
<i>Azteca fasciata nigricans</i> Forel, 1899	pn	Shattuck 1994
<i>Azteca fasciata similis</i> Mann, 1916	br	Shattuck 1994
<i>Azteca fiebrigii</i> Forel, 1909	pr	Shattuck 1994
<i>Azteca forelii</i> Emery, 1893	cr ho gu	Emery 1893; Shattuck 1994
<i>Azteca forelii championi</i> Forel, 1899	co	Shattuck 1994
<i>Azteca forelii eiseni</i> Pergande, 1896	me	Shattuck 1994
<i>Azteca forelii raptrix</i> Forel, 1912	co	Shattuck 1994
<i>Azteca forelii ursina</i> Forel, 1899	tt	Shattuck 1994
<i>Azteca forelii xysticola</i> Forel, 1899	co pn	Shattuck 1994
<i>Azteca foveiceps</i> Wheeler W.M. , 1921	gi	Shattuck 1994
<i>Azteca gnava</i> Forel, 1906	cr	Shattuck 1994
<i>Azteca gnava cayennensis</i> Forel, 1912	gi	Shattuck 1994
<i>Azteca gnava surubrensis</i> Forel, 1912	pn cr	Shattuck 1994
<i>Azteca godmani</i> Forel, 1899	pn	Shattuck 1994
<i>Azteca goeldii</i> Forel, 1906	br	Shattuck 1994
<i>Azteca huberti</i> Forel, 1906	br	Shattuck 1994
<i>Azteca hypophylla</i> Forel, 1899	co	Shattuck 1994
<i>Azteca iheringi</i> Forel, 1915	br	Shattuck 1994
<i>Azteca instabilis</i> Smith F., 1862	pe br gi gf su co pn cr be ho gu me	Emery 1893; Shattuck 1994

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Azteca instabilis major</i> Forel, 1899	gi gf su co pn	Shattuck 1994
<i>Azteca instabilis mexicana</i> Emery, 1896	ho me	Shattuck 1994
<i>Azteca isthmica</i> Wheeler W.M., 1942	br pn	Longino 1991; Shattuck 1994
<i>Azteca jelskii</i> Emery, 1893	gi gf su tt vn	Emery 1893; Shattuck 1994
<i>Azteca lallemandi</i> Forel, 1899	co	Shattuck 1994
<i>Azteca lanuginosa</i> Emery, 1893	br	Emery 1893; Shattuck 1994
<i>Azteca lanuginosa clariceps</i> Santschi, 1933	ar pr	Shattuck 1994
<i>Azteca lanuginosa pruinosa</i> Mann, 1916	br	Shattuck 1994
<i>Azteca lattkei</i> Longino, 1991	vn	Longino 1991; Shattuck 1994
<i>Azteca longiceps</i> Emery, 1893	pn cr gu	Emery 1893; Shattuck 1994
<i>Azteca longiceps cordincola</i> Forel, 1921	bo	Shattuck 1994
<i>Azteca longiceps juruensis</i> Forel, 1904	bo br	Shattuck 1994
<i>Azteca longiceps patruelis</i> Forel, 1908	me	Shattuck 1994
<i>Azteca longiceps sapii</i> Forel, 1912	br	Shattuck 1994
<i>Azteca lucida</i> Forel, 1899	gu	Longino 1989; Shattuck 1994
<i>Azteca luederwaldti</i> Forel, 1909	ar pr	Shattuck 1994
<i>Azteca mayrii</i> Emery, 1893	br	Emery 1893; Shattuck 1994
<i>Azteca merida</i> Longino, 1991	vn	Longino 1991; Shattuck 1994
<i>Azteca minor</i> Forel, 1904	br	Shattuck 1994
<i>Azteca muelleri</i> Emery, 1893	bo br es	Emery 1893; Longino 1991; Shattuck 1994
<i>Azteca muelleri pallida</i> Stitz, 1937	me	Shattuck 1994
<i>Azteca muelleri terminalis</i> Mann, 1916	br	Shattuck 1994
<i>Azteca olitrix</i> Forel, 1904	br	Shattuck 1994
<i>Azteca ovaticeps</i> Forel, 1904	bo pe br tt vn cr	Longino 1989 1991; Shattuck 1994
<i>Azteca paraensis</i> Forel, 1904	br	Shattuck 1994
<i>Azteca paraensis bondari</i> Borgmeier, 1937	br	Shattuck 1994
<i>Azteca petalocephala</i> Longino, 1991	bo	Longino 1991; Shattuck 1994
<i>Azteca pittieri</i> Forel, 1899	pn cr	Shattuck 1994
<i>Azteca pittieri emarginatisquamis</i> Forel, 1921	cr	Shattuck 1994
<i>Azteca polymorpha</i> Forel, 1899	co	Shattuck 1994
<i>Azteca polymorpha bahiana</i> Forel, 1912	br	Shattuck 1994
<i>Azteca prorsa</i> Wheeler W.M., 1942	gu	Shattuck 1994
<i>Azteca salti</i> Wheeler W.M., 1945	co	Longino 1991; Shattuck 1994
<i>Azteca schimperi</i> Emery, 1893	br gi gf su cr be	Shattuck 1994
<i>Azteca schumannii</i> Emery, 1893	vn co	Emery 1893; Shattuck 1994
<i>Azteca schumannii dubia</i> Mann, 1916	br	Shattuck 1994
<i>Azteca schumannii taediosa</i> Forel, 1904	br	Shattuck 1994
<i>Azteca sericea</i> (Mayr, 1866)	gi gf su cr gu me	Emery 1893; Shattuck 1994
<i>Azteca severini</i> Emery, 1896	br	Shattuck 1994
<i>Azteca stanleyuli</i> Forel, 1921	br	Shattuck 1994
<i>Azteca stigmatica</i> Emery, 1896	br	Shattuck 1994
<i>Azteca stollii</i> Forel, 1912	gu	Shattuck 1994
<i>Azteca subopaca</i> Forel, 1899	pn	Shattuck 1994
<i>Azteca tachigaliae</i> Forel, 1904	pe	Shattuck 1994
<i>Azteca theresiae</i> Forel, 1899	co	Shattuck 1994
<i>Azteca theresiae menceps</i> Forel, 1912	pn	Shattuck 1994
<i>Azteca tonduzi</i> Forel, 1899	cr	Shattuck 1994
<i>Azteca tonduzi columbica</i> Forel, 1912	co	Shattuck 1994
<i>Azteca trailii</i> Emery, 1893	br gi gf su	Emery 1893; Shattuck 1994
<i>Azteca trailii filicis</i> Forel, 1904	pe br	Shattuck 1994
<i>Azteca trailii tillandsiarum</i> Wheeler W.M., 1942	gi	Shattuck 1994
<i>Azteca trailii tococae</i> Forel, 1904	pe br	Shattuck 1994
<i>Azteca trianguliceps</i> Forel, 1912	br	Shattuck 1994
<i>Azteca trigona</i> Emery, 1893	br gi gf su co pn	Emery 1893; Shattuck 1994
<i>Azteca trigona gaigei</i> Forel, 1914	co	Shattuck 1994
<i>Azteca trigona mathildae</i> Forel, 1906	br	Shattuck 1994
<i>Azteca mediopsis</i> Forel, 1904	br gi gf su tt	Shattuck 1994
<i>Azteca trigona subdentata</i> Forel, 1904	br gi gf su	Shattuck 1994
<i>Azteca ulei</i> Forel, 1904	br	Shattuck 1994
<i>Azteca ulei cordiae</i> Forel, 1904	bo br gi gf su	Shattuck 1994

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Azteca ulei gagatina</i> Wheeler W.M., 1942	bo	Shattuck 1994
<i>Azteca ulei nigricornis</i> Forel, 1904	br	Shattuck 1994
<i>Azteca ulei rossi</i> Forel, 1909	me	Shattuck 1994
<i>Azteca velox</i> Forel, 1899	ec br gi gf su tt co pn cr ni gu	Shattuck 1994
<i>Azteca velox nigra</i> Forel, 1912	pn	Shattuck 1994
<i>Azteca velox nigriventris</i> Forel, 1899	br tt co pn cr	Shattuck 1994
<i>Azteca velox rectinota</i> Forel, 1908	cr	Shattuck 1994
<i>Azteca velox rochae</i> Forel, 1912	br	Shattuck 1994
<i>Azteca xanthochroa</i> Roger, 1863	tt cr be	Longino 1991; Shattuck 1994
<i>Dolichoderus abruptus</i> (Smith F., 1858)	bo pe br co	MacKay 1993; Shattuck 1994
<i>Dolichoderus andinus</i> (Kempf, 1962)	pe	MacKay 1993; Shattuck 1994
<i>Dolichoderus antiquus</i> * Carpenter, 1930	us	MacKay 1993; Shattuck 1994
<i>Dolichoderus attelaboides</i> (Fabricius, 1775)	bo pe br gi gf su tt	MacKay 1993; Shattuck 1994
<i>Dolichoderus baenae</i> MacKay, 1993	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus bidens</i> (Linnaeus, 1758)	br gi gf su tt	MacKay 1993; Shattuck 1994
<i>Dolichoderus bispinosus</i> (Olivier, 1792)	pe br gi gf su tt vn co	MacKay 1993; Shattuck 1994
	pn cr be ho gu me	MacKay 1993; Shattuck 1994
<i>Dolichoderus caribbaeus</i> * (Wilson, 1985)	an	MacKay 1993; Wilson 1985; Shattuck 1994
<i>Dolichoderus cogitans</i> Forel, 1912	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus curvifolius</i> (Lattke, 1987)	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus debilis</i> Emery, 1890	bo br gi gf su tt vn co pn gu	MacKay 1993; Shattuck 1994
<i>Dolichoderus decollatus</i> Smith, F. 1858	bo pe br gi gf su tt co	MacKay 1993; Shattuck 1994
<i>Dolichoderus dibolius</i> * Wilson, 1985	am	MacKay 1993; Wilson 1985; Shattuck 1994
<i>Dolichoderus diversus</i> Emery, 1894	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus doloniger</i> (Roger, 1862)	bo vn	MacKay 1993; Shattuck 1994
<i>Dolichoderus epetreius</i> (Lattke, 1987)	vn	MacKay 1993; Shattuck 1994
<i>Dolichoderus fernandezi</i> Mackay, 1993	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus gagates</i> Emery, 1890	bo br gi gf su vn	MacKay 1993; Shattuck 1994
<i>Dolichoderus germaini</i> Emery, 1894	pr br	MacKay 1993; Shattuck 1994
<i>Dolichoderus ghilianii</i> Emery, 1894	pe br	MacKay 1993; Shattuck 1994
<i>Dolichoderus haradai</i> MacKay, 1993	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus imitator</i> Emery, 1894	bo br	MacKay 1993; Shattuck 1994
<i>Dolichoderus inermis</i> MacKay, 1993	cr	MacKay 1993; Shattuck 1994
<i>Dolichoderus inopai</i> (Harada, 1987)	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus intermedius</i> MacKay, 1993	an	MacKay 1993; Shattuck 1994
<i>Dolichoderus lamellosus</i> (Mayr, 1870)	ar bo br gi gf su vn co cr	MacKay 1993; Shattuck 1994
<i>Dolichoderus laminatus</i> (Mayr, 1870)	br gi gf su vn co pn	MacKay 1993; Shattuck 1994
<i>Dolichoderus laurae</i> MacKay, 1993	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus lobicornis</i> (Kempf, 1959)	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus longicollis</i> MacKay, 1993	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus luederwaldti</i> Santschi, 1921	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus lugens</i> Emery, 1894	bo br gi gf su	MacKay 1993; Shattuck 1994
<i>Dolichoderus lujae</i> Santschi, 1923	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus lutosus</i> (Smith F., 1858)	pr br gi gf su tt vn co	MacKay 1993; Shattuck 1994
	pn cr ho me	MacKay 1993; Shattuck 1994
<i>Dolichoderus mariae</i> Forel, 1885	us	MacKay 1993; Shattuck 1994
<i>Dolichoderus mesonotalis</i> Forel, 1907	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus obliteratus</i> * (Scudder, 1877)	us	MacKay 1993; Shattuck 1994
<i>Dolichoderus omacanthus</i> (Kempf, 1972)	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus piceus</i> MacKay, 1993	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus plagiatus</i> (Mayr, 1870)	us	MacKay 1993; Shattuck 1994
<i>Dolichoderus primitivus</i> * (Wilson, 1985)	am	MacKay 1993; Wilson 1985; Shattuck 1994
<i>Dolichoderus prolaminatus</i> * (Wilson, 1985)	am	MacKay 1993; Wilson 1985; Shattuck 1994
<i>Dolichoderus pustulatus</i> Mayr, 1886	us	MacKay 1993; Shattuck 1994
<i>Dolichoderus quadridentigulatus</i> (Roger, 1862)	bo pe br vn	MacKay 1993; Shattuck 1994
<i>Dolichoderus rohweri</i> * Carpenter, 1930	us	MacKay 1993; Shattuck 1994
<i>Dolichoderus rosenbergi</i> Forel, 1911	ec	MacKay 1993; Shattuck 1994
<i>Dolichoderus rufescens</i> Mann, 1912	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus rugosus</i> Smith, F. 1858	bo pe ec br gi gf su	MacKay 1993; Shattuck 1994
<i>Dolichoderus schulzi</i> Emery, 1894	br	MacKay 1993; Shattuck 1994

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Dolichoderus septempinosus</i> Emery, 1894	bo pe ec br gi gf su vn	MacKay 1993; Shattuck 1994
<i>Dolichoderus setosus</i> (Kempf, 1959)	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus shattucki</i> MacKay, 1993	pn	MacKay 1993; Shattuck 1994
<i>Dolichoderus smithi</i> MacKay, 1993	bo	MacKay 1993; Shattuck 1994
<i>Dolichoderus spinicollis</i> (Latreille, 1817)	br vn	MacKay 1993; Shattuck 1994
<i>Dolichoderus spurius</i> Forel, 1903	gi gf su tt	MacKay 1993; Shattuck 1994
<i>Dolichoderus superaculus</i> (Lattke, 1987)	co	MacKay 1993; Shattuck 1994
<i>Dolichoderus taschenbergi</i> (Mayr, 1866)	us	MacKay 1993; Shattuck 1994
<i>Dolichoderus tristis</i> Mann, 1916	pe br	MacKay 1993; Shattuck 1994
<i>Dolichoderus validus</i> (Kempf, 1959)	pn cr	MacKay 1993; Shattuck 1994
<i>Dolichoderus varians</i> Mann, 1916	br	MacKay 1993; Shattuck 1994
<i>Dolichoderus voraginosus</i> MacKay, 1993	br	MacKay 1993; Shattuck 1994
<i>Dorymyrmex agallardoi</i> Snelling, 1975	ch	Snelling 1975; Shattuck 1994
<i>Dorymyrmex antarcticus</i> Forel, 1904	ar ch	Shattuck 1994
<i>Dorymyrmex baeri</i> André, 1903	ar	Shattuck 1994
<i>Dorymyrmex bicolor</i> Wheeler W.M., 1906	ar me us	Shattuck 1994
<i>Dorymyrmex biconis</i> Forel, 1912	co	Shattuck 1994
<i>Dorymyrmex bituber</i> Santschi, 1916	ar	Shattuck 1994
<i>Dorymyrmex breviscapis</i> Forel, 1912	ar	Shattuck 1994
<i>Dorymyrmex breviscapis alvarezi</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex breviscapis elongatus</i> Santschi, 1919	ar	Shattuck 1994
<i>Dorymyrmex breviscapis speculiceps</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex bruchi</i> Forel, 1912	ar	Shattuck 1994
<i>Dorymyrmex bruchi ebeninus</i> Forel, 1914	ar	Shattuck 1994
<i>Dorymyrmex brunneus</i> Forel, 1908	ar pr bo br gi gf su	Shattuck 1994
<i>Dorymyrmex carettei</i> Forel, 1913	ar	Shattuck 1994
<i>Dorymyrmex caretteoides</i> Forel, 1914	ar	Shattuck 1994
<i>Dorymyrmex chilensis</i> Forel, 1911	ch	Shattuck 1994
<i>Dorymyrmex confusus</i> (Kusnezov, 1952)	ar	Shattuck 1994
<i>Dorymyrmex conicus</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex emmaericaellus</i> Kusnezov, 1951	bo	Shattuck 1994
<i>Dorymyrmex ensifer</i> Forel, 1912	ar	Shattuck 1994
<i>Dorymyrmex ensifer laevigatus</i> Gallardo, 1916	ar	Shattuck 1994
<i>Dorymyrmex ensifer weiseri</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex exsanguis</i> Forel, 1912	ar	Shattuck 1994
<i>Dorymyrmex exanguis anaemicus</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex exanguis carbonarius</i> Forel, 1913	ar	Shattuck 1994
<i>Dorymyrmex exanguis sordidus</i> Santschi, 1919	ar	Shattuck 1994
<i>Dorymyrmex flavescens</i> Mayr, 1866	ar	Shattuck 1994
<i>Dorymyrmex flavescens jactans</i> Santschi, 1916	ar	Shattuck 1994
<i>Dorymyrmex flavescens mandibularis</i> Santschi, 1925	ar	Shattuck 1994
<i>Dorymyrmex fusculus</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex goeldii</i> Forel, 1904	br	Shattuck 1994
<i>Dorymyrmex goeldii dubius</i> Forel, 1912	br	Shattuck 1994
<i>Dorymyrmex goeldii fumigatus</i> Forel, 1908	br	Shattuck 1994
<i>Dorymyrmex goestchi</i> Goetsch, 1933	ch	Shattuck 1994
<i>Dorymyrmex hunti</i> (Snelling, 1975)	ch	Shattuck 1994
<i>Dorymyrmex hypocritus</i> (Snelling, 1975)	ch	Shattuck 1994
<i>Dorymyrmex incomptus</i> (Snelling, 1975)	ch	Shattuck 1994
<i>Dorymyrmex insanus</i> (Buckley, 1866)	pr br vn co cr gu me am	Shattuck 1994
<i>Dorymyrmex jheringi</i> Forel, 1912	br	Shattuck 1994
<i>Dorymyrmex joergensi</i> Bruch, 1917	ar	Shattuck 1994
<i>Dorymyrmex minutus</i> Emery, 1895	ar ch	Shattuck 1994
<i>Dorymyrmex morenoi</i> Bruch, 1921	ar	Shattuck 1994
<i>Dorymyrmex morenoi patagon</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex pappodes</i> (Snelling, 1975)	ch	Shattuck 1994
<i>Dorymyrmex planidens</i> Mayr, 1868	ar ch	Shattuck 1994
<i>Dorymyrmex pogonius</i> (Snelling, 1975)	ch	Shattuck 1994
<i>Dorymyrmex pulchellus</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex pyramicus</i> Roger, 1863	ar pr ur pe br gi gf su co cr be gu me us am an	Shattuck 1994

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Dorymyrmex pyramicus alticonis</i> Forel, 1912	br	Shattuck 1994
<i>Dorymyrmex pyramicus garbei</i> Foel, 1911	ar ur	Shattuck 1994
<i>Dorymyrmex pyramicus guyanensis</i> Santschi, 1922	gf	Shattuck 1994
<i>Dorymyrmex pyramicus mesonotalis</i> Forel, 1912	pn	Shattuck 1994
<i>Dorymyrmex pyramicus nigritiventris</i> Santschi, 1922	ar	Shattuck 1994
<i>Dorymyrmex pyramicus paranensis</i> Santschi, 1922	pr	Shattuck 1994
<i>Dorymyrmex pyramicus peruvianus</i> Wheeler W.M., 1919	pe	Shattuck 1994
<i>Dorymyrmex pyramicus rubriceps</i> Forel, 1912	br	Shattuck 1994
<i>Dorymyrmex richteri</i> Forel, 1912	ar	Shattuck 1994
<i>Dorymyrmex santschii</i> Gallardo, 1917	ar	Shattuck 1994
<i>Dorymyrmex silvestrii</i> Gallardo, 1916	ar	Shattuck 1994
<i>Dorymyrmex spurius</i> Santschi, 1929	ar pr ur br	Shattuck 1994
<i>Dorymyrmex steigeri</i> Santschi, 1912	ar ur	Shattuck 1994
<i>Dorymyrmex tener</i> Mayr, 1868	ar ch	Shattuck 1994
<i>Dorymyrmex tener donisthorpei</i> (Santschi, 1936)	ch	Shattuck 1994
<i>Dorymyrmex thoracicus</i> Gallardo, 1916	ar	Shattuck 1994
<i>Dorymyrmex thoracicus gracilis</i> Santschi, 1920	ar	Shattuck 1994
<i>Dorymyrmex thoracicus tigris</i> Santschi, 1925	ar	Shattuck 1994
<i>Dorymyrmex wolffhuegeli</i> Forel, 1911	ar	Shattuck 1994
<i>Forelius albiventris</i> Forel, 1912	ar br	Shattuck 1994; Cuezzo 2000
<i>Forelius analis</i> Cuezzo, 2000	me us	Shattuck 1994; Cuezzo 2000
<i>Forelius andinus</i> Kuznezov, 1957	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius bahianus</i> Cuezzo, 2000	br	Shattuck 1994; Cuezzo 2000
<i>Forelius brasiliensis</i> Forel, 1908	ar pr ur bo br	Shattuck 1994; Cuezzo 2000
<i>Forelius breviscapus</i> Forel, 1914	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius breviscapus pusillus</i> Santschi, 1922	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius chalybaeus</i> Emery, 1906	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius fiebrigii</i> Forel, 1912	pr	Shattuck 1994; Cuezzo 2000
<i>Forelius golbachii</i> (Kusnezov, 1953)	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius grandis</i> Forel 1912	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius instabilis</i> Santschi, 1920	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius lilloi</i> Cuezzo, 2000	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius macrops</i> Kusnezov, 1957	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius maranhaoensis</i> Cuezzo, 2000	br	Shattuck 1994; Cuezzo 2000
<i>Forelius nigriventris</i> Forel, 1912	ar pr bo	Shattuck 1994; Cuezzo 2000
<i>Forelius rubriceps</i> Gallardo, 1916	ar	Shattuck 1994; Cuezzo 2000
<i>Forelius rufus</i> Gallardo, 1916	ar bo	Shattuck 1994; Cuezzo 2000
<i>Forelius mccooki</i> McCook, 1879	me us am	Shattuck 1994; Cuezzo 2000
<i>Leptomyrmex neotropicus*</i> Baroni Urbani, 1980	ad	Shattuck 1994
<i>Linepithema aspidocoptum</i> (Kempf, 1969)	br	Shattuck 1994
<i>Linepithema dispertitum</i> (Forel, 1885)	bo co pn gu	Shattuck 1994
<i>Linepithema fuscum</i> Mayr, 1866	pe	Shattuck 1994
<i>Linepithema humile</i> (Mayr, 1868)	ar ur br co	Shattuck 1994
<i>Linepithema humile angulatum</i> (Emery, 1894)	ar bo br	Shattuck 1994
<i>Linepithema humile breviscapum</i> (Santschi, 1929)	ar ur	Shattuck 1994
<i>Linepithema humile platense</i> (Forel, 1912)	ar	Shattuck 1994
<i>Linepithema humile scotti</i> (Santschi, 1919)	ar	Shattuck 1994
<i>Linepithema humiloides*</i> (Wilson, 1985)	ad	Wilson 1985; Shattuck 1994
<i>Linepithema impotens</i> (Santschi, 1923)	br	Shattuck 1994
<i>Linepithema inacatum</i> Bolton, 1995	ar	Shattuck 1994; Bolton 1995
<i>Linepithema iniquum</i> (Mayr, 1870)	br vn co am an	Shattuck 1994
<i>Linepithema iniquum bicolor</i> (Forel, 1912)	br an	Shattuck 1994
<i>Linepithema iniquum nigellum</i> (Emery, 1890)	gi gf su cr gu me am	Shattuck 1994
<i>Linepithema iniquum succineum</i> (Forel, 1908)	br	Shattuck 1994
<i>Linepithema keiteli</i> (Forel, 1907)	am	Shattuck 1994
<i>Linepithema keiteli flavescens</i> (Wheeler W.M. & Mann, 1914)	am	Shattuck 1994
<i>Linepithema keiteli subfasciatum</i> (Wheeler W.M. & Mann, 1914)	am	Shattuck 1994
<i>Linepithema leucomelas</i> (Emery, 1894)	br	Shattuck 1994

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Linepithema melleum</i> (Wheeler W.M., 1908)	am	Shattuck 1994
<i>Linepithema melleum dominicense</i> (Wheeler W.M., 1913)	an	Shattuck 1994
<i>Linepithema melleum fuscescens</i> (Wheeler W.M., 1908)	am	Shattuck 1994
<i>Linepithema oblongum</i> (Santschi, 1929)	ar ch	Shattuck 1994
<i>Linepithema piliferum</i> (Mayr, 1870)	co	Shattuck 1994
<i>Linepithema pordescens</i> (Wheeler W.M., 1942)	cr	Shattuck 1994
<i>Linepithema riograndense</i> (Borgmeier, 1928)	br	Shattuck 1994
<i>Tapinoma amazone</i> Wheeler W.M., 1934	br	Shattuck 1994
<i>Tapinoma antarcticum</i> Forel, 1904	ch	Shattuck 1994
<i>Tapinoma atriceps</i> Emery, 1888	br	Shattuck 1994
<i>Tapinoma atriceps breviscapum</i> Forel, 1908	br	Shattuck 1994
<i>Tapinoma gibbosum</i> Stitz, 1933	br	Shattuck 1994
<i>Tapinoma heyeri</i> Forel, 1902	ur	Shattuck 1994
<i>Tapinoma heyeri risii</i> Forel, 1912	ur	Shattuck 1994
<i>Tapinoma litorale cubaense</i> Wheeler W.M., 1913	am	Shattuck 1994
<i>Tapinoma melanocephalum</i> Fabricius, 1793	ec br gi gf su tt co cr be ho me an bh	Shattuck 1994
<i>Tapinoma melanocephalum coronatum</i> Forel, 1908	cr am	Shattuck 1994
<i>Tapinoma opacum</i> Wheeler W.M. & Mann 1914	am	Shattuck 1994
<i>Tapinoma panamense</i> Wheeler W.M., 1934	pn	Shattuck 1994
<i>Tapinoma ramulorum</i> Emery, 1896	cr	Shattuck 1994
<i>Tapinoma ramulorum annellatum</i> Wheeler, 1934	cr	Shattuck 1994
<i>Tapinoma ramulorum inrectum</i> Forel, 1908	pn cr gu me	Shattuck 1994
<i>Tapinoma ramulorum satullum</i> Wheeler W.M., 1934	me	Shattuck 1994
<i>Tapinoma ramulorum toltecum</i> Wheeler W.M., 1934	me	Shattuck 1994
<i>Tapinoma rasenum</i> Smith D.R. & Lavigne, 1973	am	Shattuck 1994
<i>Tapinoma troche*</i> Wilson, 1985	ad	Shattuck 1994
<i>Technomyrmex caritatis*</i> Brandao & Baroni Urbani, 1998	ad	Shattuck 1994
<i>Technomyrmex fulvus</i> (Wheeler W.M. 1943)	pn	Shattuck 1994
<i>Technomyrmex fulvus sublucidus</i> (Wheeler W.M. 1943)	pn	Shattuck 1994
<i>Technomyrmex hispaniolae*</i> (Wilson, 1985)	ad	Shattuck 1994
Subfamily Formicinae		
Tribe Camponotini		
<i>Camponotus abdominalis</i> (Fabricius, 1804)	ar pr ur bo ec br gi gf su tt vn co pn cr ni be ho gu me us	MacKay 2005
<i>Camponotus abscisus</i> Roger, 1863	br pn gu me	MacKay 2005
<i>Camponotus abunanus</i> Mann, 1916	br	MacKay 2005
<i>Camponotus ager</i> (Smith F., 1858)	br gi gf su tt vn co	MacKay 2005
<i>Camponotus aguerai</i> Kusnezov, 1952	ar	MacKay 2005
<i>Camponotus alacer</i> Forel, 1912	br	MacKay 2005
<i>Camponotus albicoxis</i> Forel, 1899	cr	MacKay 2005
<i>Camponotus alboannulatus</i> Mayr, 1887	bo br	MacKay 2005
<i>Camponotus altivagans</i> Wheeler W.M., 1936	am	MacKay 2005
<i>Camponotus amoris</i> Forel, 1904	br	MacKay 2005
<i>Camponotus andrei</i> Forel, 1885	me	MacKay 2005
<i>Camponotus apostolus</i> Forel, 1901	cr	MacKay 2005
<i>Camponotus apicalis</i> (Mann, 1916)	bo pe ec tt vn co	Fernández 2002a
<i>Camponotus arboreus</i> (Smith F., 1858)	br pn	MacKay 2005
<i>Camponotus atriceps</i> (Smith F., 1858)	ar pr ur bo ec br gi gf su tt vn co pn cr ni be ho gu me us	MacKay 2005
<i>Camponotus auricomus</i> Roger, 1862	gi gf su tt co me am an	MacKay 2005
<i>Camponotus balzani</i> Emery, 1894	bo br	MacKay 2005
<i>Camponotus banghaasi</i> Emery, 1903	pe	MacKay 2005
<i>Camponotus baretoi</i> Santschi, 1922	ar	MacKay 2005
<i>Camponotus basirectus</i> Wheeler W.M., 1923	br	MacKay 2005
<i>Camponotus beebei</i> Wheeler W.M., 1918	gi gf su tt	MacKay 2005
<i>Camponotus bermudezi</i> Aguayo, 1932	am	MacKay 2005
<i>Camponotus bidens</i> Mayr, 1870	br gi gf su tt co pn	MacKay 2005
<i>Camponotus biolleyi</i> Forel, 1902	cr	MacKay 2005
<i>Camponotus bispinosus</i> Mayr, 1870	co	MacKay 2005

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Camponotus blandus</i> (Smith F., 1858)	ar pr ur bo br gi gf su	MacKay 2005
<i>Camponotus bocki</i> Forel, 1907	bo	MacKay 2005
<i>Camponotus bonariensis</i> Mayr, 1868	ar ur br	MacKay 2005
<i>Camponotus borellii</i> Emery, 1894	ar	MacKay 2005
<i>Camponotus branneri</i> Mann, 1916	br	MacKay 2005
<i>Camponotus brevis</i> Forel, 1899	co pn cr	MacKay 2005
<i>Camponotus bruchi</i> Forel, 1912	ar bo	MacKay 2005
<i>Camponotus bugnioni</i> Forel, 1899	co	MacKay 2005
<i>Camponotus caccius</i> Emery, 1903	pe br	MacKay 2005
<i>Camponotus cameroni</i> Forel, 1892	ec	MacKay 2005
<i>Camponotus canescens</i> Mayr, 1870	bo br tt co pn cr	MacKay 2005
<i>Camponotus capperi</i> Forel, 1899	am	MacKay 2005
<i>Camponotus caracalla</i> Forel, 1912	br	MacKay 2005
<i>Camponotus cerberulus</i> Emery, 1920	me	MacKay 2005
<i>Camponotus championi</i> Forel, 1899	pn me	MacKay 2005
<i>Camponotus chartifex</i> (F. Smith, 1860)	br gi gf su co pn cr ni ho	Fernández 2002a
<i>Camponotus chazaliei</i> Forel, 1899	an	MacKay 2005
<i>Camponotus cheesmanae</i> Donisthorpe, 1932	co	MacKay 2005
<i>Camponotus chilensis</i> (Spinola, 1851)	ar ch pe	MacKay 2005
<i>Camponotus christophei</i> Wheeler W.M. & Mann, 1914	am	MacKay 2005
<i>Camponotus chubutensis</i> Forel, 1931	ar br	MacKay 2005
<i>Camponotus cillae</i> Forel, 1912	br	MacKay 2005
<i>Camponotus cingulatus</i> Mayr, 1862	pr ur bo br	MacKay 2005
<i>Camponotus clarithorax</i> Creighton, 1950	us	MacKay 2005
<i>Camponotus claviscapus</i> Forel, 1899	br tt vn pn ni be ho gu	MacKay 2005
<i>Camponotus cocosensis</i> Wheeler W.M., 1919	cr	MacKay 2005
<i>Camponotus columbicus</i> Forel, 1899	co	MacKay 2005
<i>Camponotus compositor</i> Santschi, 1922	br	MacKay 2005
<i>Camponotus conspicuus</i> (Smith F., 1858)	ar ur br pn cr gu am an bh	MacKay 2005
<i>Camponotus constructor</i> Forel, 1899	pn	MacKay 2005
<i>Camponotus conulus</i> Mayr, 1870	co	MacKay 2005
<i>Camponotus convexiclypeus</i> MacKay, 1997	pe	MacKay 1997
<i>Camponotus corticalis</i> Forel, 1899	am	MacKay 2005
<i>Camponotus coruscus</i> (Smith F., 1862)	su vn co pn cr gu me	MacKay 2005
<i>Camponotus crassicornis</i> Emery, 1920	bo br	MacKay 2005
<i>Camponotus cressoni</i> André, 1887	br me	MacKay 2005
<i>Camponotus cuneidorsus</i> Emery, 1920	pe	MacKay 2005
<i>Camponotus curviscapus</i> Emery, 1896	cr	MacKay 2005
<i>Camponotus dalmasi</i> Forel, 1899	co	MacKay 2005
<i>Camponotus decessor</i> Forel, 1908	br	MacKay 2005
<i>Camponotus deletangi</i> Santschi, 1920	bo	MacKay 2005
<i>Camponotus dimorphus</i> Emery, 1894	bo br	MacKay 2005
<i>Camponotus distinguendus</i> (Spinola, 1851)	ar ch pe	MacKay 2005
<i>Camponotus diversipalpus</i> Santschi, 1922	br	MacKay 2005
<i>Camponotus dumetorum</i> Wheeler W.M., 1910	us	MacKay 2005
<i>Camponotus egregius</i> (Smith F., 1858)	br	MacKay 2005
<i>Camponotus elevatus</i> Forel, 1899	cr	MacKay 2005
<i>Camponotus essigi</i> Smith M.R., 1923	us	MacKay 2005
<i>Camponotus eurynotus</i> Forel, 1907	bo br	MacKay 2005
<i>Camponotus excisus</i> Mayr, 1870	br tt co pn	MacKay 2005
<i>Camponotus fasciatellus</i> Dalla Torre, 1892	bo gi gf su co me	MacKay 2005
<i>Camponotus fastigatus</i> Roger, 1863	bo br gi gf su tt co	MacKay 2005
<i>Camponotus femoratus</i> (Fabricius, 1804)	pe br gi gf su tt vn	MacKay 2005
<i>Camponotus festinatus</i> Buckley, 1866	me us	MacKay 2005
<i>Camponotus flavescentis</i> (Fabricius, 1793)	gf	MacKay 2005
<i>Camponotus folicola</i> Forel, 1904	br	MacKay 2005
<i>Camponotus formiciformis</i> Forel, 1885	be gu me	MacKay 2005
<i>Camponotus frontalis</i> Pergande, 1896	me	MacKay 2005
<i>Camponotus fryi</i> Mann, 1916	br	MacKay 2005
<i>Camponotus sumidus</i> Roger, 1863	ar br vn me us am an bh	MacKay 2005

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Camponotus genatus</i> Santschi, 1922	br	MacKay 2005
<i>Camponotus geraensis</i> Emery, 1920	ar pr bo br	MacKay 2005
<i>Camponotus germaini</i> Emery, 1903	pr br	MacKay 2005
<i>Camponotus gilviventris</i> Roger, 1863	am	MacKay 2005
<i>Camponotus godmani</i> Forel, 1899	br tt pn me	MacKay 2005
<i>Camponotus goeldii</i> Forel, 1894	br	MacKay 2005
<i>Camponotus gundlachi</i> Mann, 1920	am	MacKay 2005
<i>Camponotus haematocephalus</i> Emery, 1903	pe	MacKay 2005
<i>Camponotus hannani</i> Forel, 1899	am	MacKay 2005
<i>Camponotus hedwigae</i> Forel, 1912	br	MacKay 2005
<i>Camponotus helleri</i> Emery, 1903	pe	MacKay 2005
<i>Camponotus hermanni</i> Emery, 1911	br	MacKay 2005
<i>Camponotus holzi</i> Forel, 1921	ec	MacKay 2005
<i>Camponotus hyatti</i> Emery, 1893	us	MacKay 2005
<i>Camponotus hybridus</i> Forel, 1901	ar	MacKay 2005
<i>Camponotus ignestii</i> Menozzi, 1935	gi	MacKay 2005
<i>Camponotus improprius</i> (Forel, 1879)	co	MacKay 2005
<i>Camponotus inca</i> Emery, 1903	pe	MacKay 2005
<i>Camponotus indicatus</i> Santschi, 1922	gf	MacKay 2005
<i>Camponotus inlex</i> Forel, 1901	br	MacKay 2005
<i>Camponotus innocens</i> Forel, 1909	pr	MacKay 2005
<i>Camponotus integellus</i> Forel, 1899	pe ec br vn co cr	MacKay 2005
<i>Camponotus juliae</i> Emery, 1903	cr	MacKay 2005
<i>Camponotus kaura</i> Snelling y Torres 1998	am	MacKay 2005
<i>Camponotus keiferi</i> Wheeler W.M., 1934	me	MacKay 2005
<i>Camponotus koseritzii</i> Emery, 1888	bo br	MacKay 2005
<i>Camponotus kraepelini</i> Forel, 1901	bo	MacKay 2005
<i>Camponotus laevigatus</i> (Smith F., 1858)	us	MacKay 2005
<i>Camponotus lancifer</i> Emery, 1894	br	MacKay 2005
<i>Camponotus landolti</i> Forel, 1879	br co pn cr me	MacKay 2005
<i>Camponotus larvigerus</i> Wheeler W.M. & Mann, 1914	am	MacKay 2005
<i>Camponotus latangulus</i> Roger, 1863	bo br gi gf su tt	MacKay 2005
<i>Camponotus lenkoi</i> Kempf, 1960	br	MacKay 2005
<i>Camponotus leptcephalus</i> Emery, 1923	br	MacKay 1997
<i>Camponotus lespesii</i> Forel, 1886	pr br co pn	MacKay 2005
<i>Camponotus leydigi</i> Forel, 1886	ar pr br gi gf su	MacKay 2005
<i>Camponotus lillii</i> Forel, 1908	br	MacKay 2005
<i>Camponotus lindigi</i> Mayr, 1870	tt vn co pn cr ni be ho an	MacKay 2005
<i>Camponotus linnaei</i> Forel, 1886	pe br co pn	MacKay 2005
<i>Camponotus longipilis</i> Emery, 1911	pe bo	MacKay 1997
<i>Camponotus lutzi</i> Forel, 1905	br	MacKay 2005
<i>Camponotus maccanni</i> Forel, 1879	me	MacKay 2005
<i>Camponotus macrochaeta</i> Emery, 1903	br	MacKay 2005
<i>Camponotus melanocephalus</i> Roger, 1863	vn	MacKay 2005
<i>Camponotus melanoticus</i> Emery, 1894	ar bo ec br me	MacKay 2005
<i>Camponotus melinus</i> MacKay, 1997	me	MacKay & MacKay 1997a
<i>Camponotus microstictus</i> Wheeler W.M., 1937	am	MacKay 2005
<i>Camponotus mina</i> Forel, 1879	me	MacKay 2005
<i>Camponotus minozzii</i> Emery, 1920	pe	MacKay 2005
<i>Camponotus mirabilis</i> Emery, 1903	pe bo br	MacKay 1997
<i>Camponotus mocsayi</i> Forel, 1902	pe br gi gf su	MacKay 2005
<i>Camponotus moelleri</i> Forel, 1912	br	MacKay 2005
<i>Camponotus montivagus</i> Forel, 1885	me gu	MacKay & MacKay 1997a
<i>Camponotus morosus</i> (Smith F., 1858)	ar ch	MacKay 2005
<i>Camponotus mus</i> Roger, 1863	ar pr ur bo br	MacKay 2005
<i>Camponotus naegelii</i> Forel, 1879	br	MacKay 2005
<i>Camponotus nepos</i> Forel, 1912	br	MacKay 2005
<i>Camponotus nidulans</i> (Smith F., 1860)	bo ec br vn co	Fernández 2002a
<i>Camponotus nitens</i> Mayr, 1870	br co	MacKay 2005
<i>Camponotus nitidior</i> (Santschi, 1921)	co pn cr ni ho	Fernández 2002a

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Camponotus novogranadensis</i> Mayr, 1870	pe br gi gf su tt co cr ho	MacKay 2005
<i>Camponotus obreptivus</i> Forel, 1899	cr	MacKay 2005
<i>Camponotus occultus</i> Wheeler W.M. & Mann, 1914	am	MacKay 2005
<i>Camponotus ogloblini</i> Santschi, 1934	ar	MacKay 2005
<i>Camponotus orthocephalus</i> Emery 1894	ar bo pe br cr	MacKay 2005
<i>Camponotus pachylepis</i> Emery, 1920	bo pe	MacKay 2005
<i>Camponotus pallescens</i> Mayr, 1887	br	MacKay 2005
<i>Camponotus panamensis</i> Fernández, 2002	pn	Fernández 2002a
<i>Camponotus patimae</i> Wheeler W.M., 1942	gi	MacKay 2005
<i>Camponotus pellarius</i> Wheeler W.M., 1914	me	MacKay 2005
<i>Camponotus personatus</i> Emery, 1894	ar br	MacKay 2005
<i>Camponotus pertusus</i> MacKay, 1997	tt	MacKay & MacKay 1997a
<i>Camponotus piceatus</i> (Norton, 1868)	me	MacKay 2005
<i>Camponotus picipes</i> (Olivier, 1792)	bo ec br gi gf su gu me am	MacKay 2005
<i>Camponotus pittieri</i> Forel, 1899	pe br cr an	MacKay 2005
<i>Camponotus planatus</i> Roger, 1863	co gu me us	MacKay 2005
<i>Camponotus planus</i> Smith F., 1877	ec	MacKay 2005
<i>Camponotus platytarsus</i> Roger, 1863	gi gf su me	MacKay 2005
<i>Camponotus plombyi</i> Wheeler W.M. & Mann, 1914	am	MacKay 2005
<i>Camponotus polymorphicus</i> MacKay, López & Fernández, 2002	co	MacKay 2005
<i>Camponotus postangulatus</i> Emery, 1925	pe	MacKay 1997
<i>Camponotus propinquellus</i> Emery, 1920	pe	MacKay 2005
<i>Camponotus propinquus</i> Mayr, 1887	br	MacKay 2005
<i>Camponotus pullatus</i> Mayr, 1866	me	MacKay 2005
<i>Camponotus punctatus</i> Forel, 1912	ar br	MacKay 2005
<i>Camponotus punctulatus</i> Mayr, 1868	ar pr ur bo pe br me	MacKay 2005
<i>Camponotus quercicola</i> Smith M.R., 1954	us	MacKay 2005
<i>Camponotus ramulorum</i> Wheeler W.M., 1905	am bh	MacKay 2005
<i>Camponotus rapax</i> (Fabricius, 1804)	br gi gf su	MacKay 2005
<i>Camponotus rectangularis</i> Emery, 1890	pe br gi gf su pn cr ni be ho gu me	MacKay 2005
<i>Camponotus rectithorax</i> Forel, 1895	gu	MacKay & MacKay 1997a
<i>Camponotus rector</i> Santschi, 1923	br	MacKay 2005
<i>Camponotus renggeri</i> Emery, 1894	ar ur bo pe br gi gf su	MacKay 2005
<i>Camponotus ruficornis</i> Emery, 1895	ch	MacKay 2005
<i>Camponotus rufipes</i> (Fabricius, 1775)	ar pr ur bo br gi gf su vn co	MacKay 2005
<i>Camponotus rufipilis</i> Aguayo, 1932	am	MacKay 2005
<i>Camponotus rusticus</i> Santschi, 1916	ar	MacKay 2005
<i>Camponotus salvini</i> Forel, 1899	pn cr	MacKay 2005
<i>Camponotus sanctaeidei</i> Dalla Torre, 1892	ar bo pe br pn cr	MacKay 2005
<i>Camponotus sansabeanus</i> (Buckley, 1866)	us	MacKay 2005
<i>Camponotus santosi</i> Forel, 1908	am bh	MacKay 2005
<i>Camponotus saussurei</i> Forel, 1879	an	MacKay 2005
<i>Camponotus schmalzi</i> Emery, 1894	br	MacKay 2005
<i>Camponotus scissus</i> Mayr, 1887	br	MacKay 2005
<i>Camponotus sculptor</i> Santschi, 1920	ar	MacKay 2005
<i>Camponotus semitestaceus</i> Snelling, 1970	us	MacKay 2005
<i>Camponotus senex</i> (Smith F., 1858)	pe ec br tt co pn cr me	MacKay 2005
<i>Camponotus sericatus</i> Mayr, 1887	br	MacKay 2005
<i>Camponotus sericeiventris</i> Guérin-Méneville, 1838	ar ur bo pe br gi gf su	MacKay 2005
<i>Camponotus setipes</i> Forel, 1904	gi gf su tt	MacKay 2005
<i>Camponotus sexguttatus</i> (Fabricius, 1793)	ar pr ur bo pe ec br gi gf su co pn ni am an	MacKay 2005
<i>Camponotus sharpi</i> Forel, 1893	gu me an	MacKay 2005
<i>Camponotus silvestrii</i> Emery, 1906	br	MacKay 2005
<i>Camponotus silvicola</i> Forel, 1902	ar pr co	MacKay 2005
<i>Camponotus simillimus</i> (Smith F., 1862)	ec br gi gf su co pn cr	MacKay 2005
<i>Camponotus socius</i> Roger, 1863	us	MacKay 2005
<i>Camponotus socorroensis</i> Wheeler W.M., 1934	me	MacKay 2005

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Camponotus sphaericus</i> Roger, 1863	am	MacKay 2005
<i>Camponotus sphenoccephalus</i> Emery, 1911	pe	MacKay 1997
<i>Camponotus sphenoidalis</i> Mayr, 1870	co	MacKay 2005
<i>Camponotus spinolae</i> Roger, 1863	ch	MacKay 2005
<i>Camponotus subcircularis</i> Emery, 1920	bo	MacKay 2005
<i>Camponotus subdepilis</i> Wheeler W.M., 1917	am	MacKay 2005
<i>Camponotus substitutus</i> Emery, 1894	ar ch pr ur bo br gi gf su vn co pn cr gu me	MacKay 2005
<i>Camponotus subtruncatus</i> Borgmeier, 1929	br	MacKay 2005
<i>Camponotus taino</i> Snelling & Torres, 1998	am	MacKay 2005
<i>Camponotus tecpanicus</i> Pergande, 1896	me	MacKay 2005
<i>Camponotus tenuiscapus</i> Roger, 1863	ar ur br	MacKay 2005
<i>Camponotus terbimaculatus</i> Emery 1920	br	MacKay 2005
<i>Camponotus termitarius</i> Emery, 1902	ar br	MacKay 2005
<i>Camponotus testaceus</i> Emery, 1894	br	MacKay 2005
<i>Camponotus thysanopus</i> Wheeler W.M., 1937	am	MacKay 2005
<i>Camponotus tonduzi</i> Forel, 1899	cr	MacKay 2005
<i>Camponotus torrei</i> Aguayo, 1932	am	MacKay 2005
<i>Camponotus traili</i> Mayr, 1878	bo br vn pn	Fernández 2002a
<i>Camponotus trapeziceps</i> Forel, 1908	br	MacKay 2005
<i>Camponotus trapezoideus</i> Mayr, 1870	br co	MacKay 2005
<i>Camponotus tripartitus</i> Mayr, 1887	br	MacKay 2005
<i>Camponotus ulcerosus</i> Wheeler W.M., 1910	us	MacKay 2005
<i>Camponotus ulei</i> Forel, 1904	pe	MacKay 2005
<i>Camponotus ulvarum</i> Forel, 1899	cr	MacKay 2005
<i>Camponotus urichi</i> Forel, 1899	br tt	MacKay 2005
<i>Camponotus ustus</i> Forel, 1879	co am an	MacKay 2005
<i>Camponotus verae</i> Forel, 1908	br	MacKay 2005
<i>Camponotus vicinus</i> Mayr, 1870	us	MacKay 2005
<i>Camponotus westermanni</i> Mayr, 1862	ar br	MacKay 2005
<i>Camponotus wheeleri</i> Mann, 1916	br	MacKay 2005
<i>Camponotus wytmani</i> Emery, 1920	bo	MacKay 2005
<i>Camponotus xanthogaster</i> Santschi, 1924	br	MacKay 2005
<i>Camponotus zenon</i> Forel, 1912	br	MacKay 2005
<i>Camponotus zoc</i> Forel, 1879	gi gf su tt vn co gu	MacKay 2005
Tribe Formicini		
<i>Formica argentea</i> Wheeler W.M., 1912	me	Wheeler 1913; Rojas-Fernández, 1996
<i>Formica francoeuri</i> Bolton, 1995	me	Wheeler 1913; Rojas-Fernández 1996
<i>Formica gnava</i> Buckley, 1866	me	Wheeler 1913; Rojas-Fernández 1996
<i>Formica integroides</i> Wheeler W.M., 1913	me	Wheeler 1913; Rojas-Fernández 1996
<i>Formica lasioides</i> Emery, 1893	me	Wheeler 1913; Rojas-Fernández 1996
<i>Formica moki</i> Wheeler W.M., 1906	me	Wheeler 1913; Rojas-Fernández 1996
<i>Formica perpilosa</i> Wheeler W.M., 1913	me	Wheeler 1913; Rojas-Fernández 1996
<i>Formica subpolita</i> Mayr, 1886	me	Wheeler 1913; Rojas-Fernández 1996
<i>Formica xerophila</i> Smith M.R., 1939	me	Wheeler 1913; Rojas-Fernández 1996
<i>Polyergus breviceps</i> Emery, 1893	me	Wheeler 1913; Rojas-Fernández 1996
Tribe Gigantiopini		
<i>Gigantiops destructor</i> (Fabricius, 1804)	pe br gi gf su vn co	Kempf & Lenko 1968
Tribe Lasiini		
<i>Acanthomyops californicus</i> Wheeler W.M., 1917	me	Rojas-Fernández 1996
<i>Acanthomyops mexicanus</i> Wheeler W.M., 1914	me	Rojas-Fernández 1996
<i>Acropyga ayanganna</i> LaPolla, 2004	gi	LaPolla 2004
<i>Acropyga decadens</i> (Mayr, 1887)	ec br	LaPolla 2004
<i>Acropyga donisthorpei</i> Weber, 1944	gi pe bo	LaPolla 2004
<i>Acropyga dubitata</i> Wheeler W.M. & Mann, 1914	am	LaPolla 2004
<i>Acropyga exsanguis</i> Wheeler W.M., 1909	me be cr gu pn co gi su ar	LaPolla 2004
<i>Acropyga fuhrmanni</i> Forel, 1914	tt cr co gi ec pe bo br	LaPolla 2004
<i>Acropyga goeldii</i> Forel, 1893	am pn tt co vn su gi br pa	LaPolla 2004
<i>Acropyga guianensis</i> Weber, 1944	gi	LaPolla 2004
<i>Acropyga hirsutula</i> LaPolla, 2004	ec	LaPolla 2004

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Acropyga keira</i> LaPolla, 2004	cr	LaPolla 2004
<i>Acropyga palaga</i> LaPolla, 2004	cr	LaPolla 2004
<i>Acropyga panamensis</i> Weber, 1944	cr pn	LaPolla 2004
<i>Acropyga parvidens</i> Wheeler W.M. & Mann, 1914	am	LaPolla 2004
<i>Acropyga romeo</i> LaPolla, 2004	gi	LaPolla 2004
<i>Acropyga smithii</i> Forel, 1893	an be ho pn tt co pe br	LaPolla 2004
<i>Acropyga tricuspidis</i> LaPolla, 2004	br	LaPolla 2004
<i>Anoplolepis gracilipes</i> Smith F., 1857	ch me	Kempf 1972
<i>Anoplolepis longipes</i> (Jerdon, 1851)	ch me	Kempf 1972
<i>Lasiophanes hoffmanni</i> (Forel, 1903)	ch	Kusnezov 1952a; Snelling & Hunt 1976
<i>Lasiophanes nigritiventris</i> (Spinola, 1851)	ar ch	Kusnezov 1952a; Snelling & Hunt 1976
<i>Lasiophanes picinus</i> (Roger, 1863)	ar ch	Kusnezov 1952a; Snelling & Hunt 1976
<i>Lasiophanes picinus bruchi</i> (Forel, 1915)	ar ch	Kusnezov 1952a
<i>Lasiophanes uxorius</i> Emery, 1922	ch	Kusnezov 1952a; Snelling & Hunt 1976
<i>Lasiophanes valdiviensis</i> Forel, 1904	ch	Kusnezov 1952a; Snelling & Hunt 1976
<i>Lasius alienus</i> Foerster, 1850	me	Rojas-Fernández 1996
<i>Lasius niger</i> (Linnaeus, 1758)	me	Rojas-Fernández 1996
<i>Lasius sitiens</i> Wilson, 1955	me	Rojas-Fernández 1996
<i>Myrmecocystus depilis</i> (Forel, 1901)	me	Rojas-Fernández 1996
<i>Myrmecocystus flaviceps</i> Wheeler W.M., 1912	us	Rojas-Fernández 1996
<i>Myrmecocystus intonsus</i> Snelling, 1976	me	Rojas-Fernández 1996
<i>Myrmecocystus kennedyi</i> Snelling, 1969	me	Rojas-Fernández 1996
<i>Myrmecocystus melliger</i> Forel, 1886	me	Rojas-Fernández 1996
<i>Myrmecocystus mendax</i> Wheeler W.M., 1908	me	Rojas-Fernández 1996
<i>Myrmecocystus mexicanus</i> Wesmael, 1838	me	Rojas-Fernández 1996
<i>Myrmecocystus mimicus</i> Wheeler W.M., 1908	us	Rojas-Fernández 1996
<i>Myrmecocystus navajo</i> Wheeler W.M., 1908	us	Rojas-Fernández 1996
<i>Myrmecocystus perimeces</i> Snelling, 1976	me	Rojas-Fernández 1996
<i>Myrmecocystus placodops</i> Forel, 1908	me	Rojas-Fernández 1996
<i>Myrmecocystus romainei</i> Cole, 1936	me	Rojas-Fernández 1996
<i>Myrmecocystus semirufus</i> Emery, 1893	me	Rojas-Fernández 1996
<i>Myrmecocystus tenuinodis</i> Snelling, 1976	me	Rojas-Fernández 1996
<i>Myrmecocystus testaceus</i> Emery, 1893	me	Rojas-Fernández 1996
<i>Myrmecocystus wheeleri</i> Snelling, 1971	me	Rojas-Fernández 1996
<i>Myrmecocystus yuma</i> Wheeler W.M., 1912	me	Rojas-Fernández 1996
Tribe Plagiolepidini		
<i>Brachymyrmex admotus</i> Mayr, 1887	ar br	Santschi 1923
<i>Brachymyrmex antennatus</i> Santschi, 1929	br	Santschi 1923
<i>Brachymyrmex australis</i> Forel, 1901	br	Santschi 1923
<i>Brachymyrmex australis curtus</i> Santschi, 1922	ar ur	Santschi 1923
<i>Brachymyrmex brevicornis</i> Emery, 1906	ar pr br gi gf su	Santschi 1923
<i>Brachymyrmex bruchi</i> Forel, 1912	ar	Santschi 1923
<i>Brachymyrmex bruchi rufipes</i> Forel, 1912	ar	Santschi 1923
<i>Brachymyrmex cavernicola</i> Wheeler W.M., 1938	me	Santschi 1923
<i>Brachymyrmex coactus</i> Mayr, 1887	ar br	Santschi 1923
<i>Brachymyrmex coactus nictitans</i> Emery, 1906	cr	Santschi 1923
<i>Brachymyrmex coactus robustus</i> Santschi, 1923	br	Santschi 1923
<i>Brachymyrmex constrictus</i> Santschi, 1923	bo	Santschi 1923
<i>Brachymyrmex constrictus bonariensis</i> Santschi, 1933	ar	Santschi 1923
<i>Brachymyrmex cordemoyi</i> Forel, 1895	ar pr br gi gf su vn co gu an	Santschi 1923
<i>Brachymyrmex cordemoyi distinctus</i> Santschi, 1923	ar	Santschi 1923
<i>Brachymyrmex degener</i> Emery, 1906	br	Santschi 1923
<i>Brachymyrmex degener niger</i> Forel, 1912	br	Santschi 1923
<i>Brachymyrmex depilis</i> Emery, 1893	us	Santschi 1923
<i>Brachymyrmex donisthorpei</i> Santschi, 1939	co	Santschi 1923
<i>Brachymyrmex fiebrigi</i> Forel, 1908	ar pr	Santschi 1923
<i>Brachymyrmex fiebrigi fumidus</i> Santschi, 1923	ar	Santschi 1923
<i>Brachymyrmex fiebrigi funicularis</i> Santschi, 1922	ar	Santschi 1923
<i>Brachymyrmex flavidulus</i> (Roger, 1863)	am	Santschi 1923
<i>Brachymyrmex gagates</i> Wheeler W.M., 1934	me	Santschi 1923

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Brachymyrmex gaucho</i> Santschi, 1917	ar	Santschi 1923
<i>Brachymyrmex giardi</i> Emery, 1895	ch	Santschi 1923
<i>Brachymyrmex giardi cordobensis</i> Santschi, 1929	ar	Santschi 1923
<i>Brachymyrmex giardi nitidus</i> Santschi, 1922	ch	Santschi 1923
<i>Brachymyrmex goeldii</i> Forel, 1912	br	Santschi 1923
<i>Brachymyrmex heeri</i> Forel, 1874	ar bo br gi gf su tt vn co cr am an bh	Santschi 1923
<i>Brachymyrmex heeri aphidicola</i> Forel, 1909	ar pr br am	Santschi 1923
<i>Brachymyrmex heeri basalis</i> Wheeler W.M., 1921	gi	Santschi 1923
<i>Brachymyrmex heeri fallax</i> Santschi, 1923	ar pr	Santschi 1923
<i>Brachymyrmex heeri obscurior</i> Forel, 1893	tt vn co gu me am an ber bh	Santschi 1923
<i>Brachymyrmex heeri termitophilus</i> Forel, 1896	ar pr br	Santschi 1923
<i>Brachymyrmex incisus</i> Forel, 1912	br co	Santschi 1923
<i>Brachymyrmex laevis</i> Emery, 1895	ch	Santschi 1923
<i>Brachymyrmex laevis andinus</i> Santschi, 1923	ar	Santschi 1923
<i>Brachymyrmex laevis fusculus</i> Emery, 1906	ar	Santschi 1923
<i>Brachymyrmex longicornis</i> Forel, 1907	ar br cr	Santschi 1923
<i>Brachymyrmex longicornis hemiops</i> Santschi, 1923	br	Santschi 1923
<i>Brachymyrmex longicornis pullus</i> Santschi, 1933	ar	Santschi 1923
<i>Brachymyrmex longicornis immunis</i> Forel, 1908	pr ur br	Santschi 1923
<i>Brachymyrmex luederwaldti</i> Santschi, 1923	br	Santschi 1923
<i>Brachymyrmex luederwaldti attenuatus</i> Santschi, 1929	br	Santschi 1923
<i>Brachymyrmex micromegas</i> Emery, 1923	br	Santschi 1923
<i>Brachymyrmex minutus</i> Forel, 1893	tt co am an bh	Santschi 1923
<i>Brachymyrmex modestus</i> Santschi, 1923	br	Santschi 1923
<i>Brachymyrmex musculus</i> Forel, 1899	cr me	Santschi 1923
<i>Brachymyrmex myops</i> Emery, 1906	br	Santschi 1923
<i>Brachymyrmex obscurior</i> Forel, 1893	tt vn co gu me am an ber bh	Santschi 1923
<i>Brachymyrmex oculatus</i> Santschi, 1919	ar	Santschi 1923
<i>Brachymyrmex patagonicus</i> Mayr, 1868	ar pr bo br gi gf su vn	Santschi 1923
<i>Brachymyrmex patagonicus atratulus</i> Santschi, 1923	ar	Santschi 1923
<i>Brachymyrmex pictus</i> Mayr, 1887	br	Santschi 1923
<i>Brachymyrmex pictus balboae</i> Wheeler W.M., 1942	pn	Santschi 1923
<i>Brachymyrmex pilipes</i> Mayr, 1887	br	Santschi 1923
<i>Brachymyrmex santschii</i> Menozzi, 1927	cr	Santschi 1923
<i>Brachymyrmex tristis</i> Mayr, 1870	co	Santschi 1923
<i>Myrmelachista ambigua</i> Forel, 1893	an	Kempf 1972
<i>Myrmelachista amicta</i> Wheeler W.M., 1934	me	Kempf 1972
<i>Myrmelachista arborea</i> Forel, 1909	pr	Kempf 1972
<i>Myrmelachista arborea nasuta</i> Forel, 1912	br	Kempf 1972
<i>Myrmelachista arthuri</i> Forel, 1903	ar br	Kempf 1972
<i>Myrmelachista arthuri brunneiceps</i> Forel, 1908	br	Kempf 1972
<i>Myrmelachista bambusarum</i> Forel, 1903	br	Kempf 1972
<i>Myrmelachista bettinae</i> Forel, 1903	br	Kempf 1972
<i>Myrmelachista brevicornis</i> Wheeler W.M., 1934	pe br	Kempf 1972
<i>Myrmelachista bruchi</i> Santschi, 1922	ar	Kempf 1972
<i>Myrmelachista catharinae</i> Mayr, 1887	br	Kempf 1972
<i>Myrmelachista catharinae maior</i> Santschi, 1936	br	Kempf 1972
<i>Myrmelachista catharinae petropolitana</i> Santschi, 1936	br	Kempf 1972
<i>Myrmelachista catharinae intermedia</i> Santschi, 1929	br	Kempf 1972
<i>Myrmelachista chilensis</i> Forel, 1904	ar ch	Kempf 1972
<i>Myrmelachista costaricensis</i> Wheeler W.M., 1934	cr	Kempf 1972
<i>Myrmelachista dalmasi</i> Forel, 1912	co	Kempf 1972
<i>Myrmelachista donisthorpei</i> Wheeler W.M., 1934	gi	Kempf 1972
<i>Myrmelachista elata</i> Santschi, 1922	br	Kempf 1972
<i>Myrmelachista elongata</i> Santschi, 1925	ar	Kempf 1972
<i>Myrmelachista flavidula</i> Wheeler W.M., 1934	gi	Kempf 1972
<i>Myrmelachista gagates</i> Wheeler W.M., 1936	am	Kempf 1972
<i>Myrmelachista gagatina</i> Emery, 1894	ar br	Kempf 1972
<i>Myrmelachista gallicola</i> Mayr, 1887	ar ur br	Kempf 1972

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Myrmelachista goeldii</i> Forel, 1903	br	Kempf 1972
<i>Myrmelachista guyanensis</i> Wheeler W.M., 1934	br gi gf su	Kempf 1972
<i>Myrmelachista hoffmanni</i> Forel, 1903	ch	Kempf 1972
<i>Myrmelachista kloetersi</i> Forel, 1903	br	Kempf 1972
<i>Myrmelachista kraatzii</i> Roger, 1863	am	Kempf 1972
<i>Myrmelachista longinoda</i> Forel, 1899	pn	Kempf 1972
<i>Myrmelachista mayri</i> Forel, 1886	ch	Kempf 1972
<i>Myrmelachista mayri monticola</i> Mayr, 1887	ch	Kempf 1972
<i>Myrmelachista mexicana</i> Wheeler W.M., 1934	me	Kempf 1972
<i>Myrmelachista muelleri</i> Forel, 1903	br	Kempf 1972
<i>Myrmelachista nigella</i> (Roger, 1863)	br vn	Kempf 1972
<i>Myrmelachista nigella ruficornis</i> Santschi, 1936	ar br	Kempf 1972
<i>Myrmelachista nodigera</i> Mayr, 1887	ur br	Kempf 1972
<i>Myrmelachista nodigera flavigaster</i> Emery, 1896	pr br	Kempf 1972
<i>Myrmelachista nodigera pallida</i> Forel, 1909	ar pr	Kempf 1972
<i>Myrmelachista paderevskii</i> Forel, 1908	br	Kempf 1972
<i>Myrmelachista plebecula</i> Menozzi, 1927	cr	Kempf 1972
<i>Myrmelachista ramulorum</i> Wheeler W.M., 1908	am an	Kempf 1972
<i>Myrmelachista ramulorum fortior</i> Wheeler W.M., 1934	am	Kempf 1972
<i>Myrmelachista reclusa</i> Forel, 1903	co	Kempf 1972
<i>Myrmelachista rectinota</i> Forel, 1904	ch	Kempf 1972
<i>Myrmelachista reichenbergeri</i> Santschi, 1922	br	Kempf 1972
<i>Myrmelachista reticulata</i> Borgmeier, 1928	br	Kempf 1972
<i>Myrmelachista rogeri</i> André, 1887	am	Kempf 1972
<i>Myrmelachista rogeri manni</i> Wheeler W.M., 1934	am	Kempf 1972
<i>Myrmelachista rogeri rubriceps</i> Mann, 1920	am	Kempf 1972
<i>Myrmelachista rudolphi</i> Forel, 1903	br	Kempf 1972
<i>Myrmelachista ruzskii</i> Forel, 1903	br	Kempf 1972
<i>Myrmelachista schachovskoi</i> Kusnezov, 1951	ar	Kempf 1972
<i>Myrmelachista schumanni</i> Emery, 1890	co	Kempf 1972
<i>Myrmelachista schumanni cordincola</i> Wheeler W.M., 1934	bo	Kempf 1972
<i>Myrmelachista skwarrae</i> Wheeler W.M., 1934	me	Kempf 1972
<i>Myrmelachista skwarrae laeta</i> Wheeler W.M., 1934	me	Kempf 1972
<i>Myrmelachista skwarrae picea</i> Wheeler W.M., 1934	me	Kempf 1972
<i>Myrmelachista ulei</i> Forel, 1904	pe	Kempf 1972
<i>Myrmelachista ulei dubia</i> Forel, 1912	br	Kempf 1972
<i>Myrmelachista vicina</i> Kusnezov, 1951	ar	Kempf 1972
<i>Myrmelachista zeledoni</i> Emery, 1896	co cr	Kempf 1972
<i>Myrmelachista zeledoni thiemei</i> Emery, 1906	pe vn co	Kempf 1972
<i>Plagiolepis alluaudi</i> Emery 1894	ber	Kempf 1972
<i>Paratrechina anthracina</i> (Roger, 1863)	am	Kempf 1972
<i>Paratrechina brasiliensis</i> (Mayr, 1862)	br	Kempf 1972
<i>Paratrechina bruesii</i> Wheeler W.M., 1903	us	Kempf 1972
<i>Paratrechina burgessi</i> Trager, 1984	pe	Kempf 1972; Trager 1984
<i>Paratrechina caeciliae</i> (Forel, 1899)	gi gf su cr	Kempf 1972
<i>Paratrechina caeciliae elevata</i> Forel, 1908	pe cr	Kempf 1972
<i>Paratrechina cisipa</i> Smith D.R. & Lavigne, 1973	am	
<i>Paratrechina darlingtoni</i> Wheeler W.M., 1936	am	Kempf 1972
<i>Paratrechina dispar</i> Forel, 1909	gu	Kempf 1972
<i>Paratrechina fulva</i> (Mayr, 1862)	ar pr ur bo br gi gf su cr me am	Kempf 1972; Trager 1974
<i>Paratrechina fulva biolleyi</i> (Forel, 1908)	cr	Kempf 1972
<i>Paratrechina fulva cubana</i> (Santschi, 1930)	am	Kempf 1972
<i>Paratrechina fulva fumata</i> (Forel, 1909)	ar pr am	Kempf 1972
<i>Paratrechina fulva fumatipennis</i> (Forel, 1915)	ar br	Kempf 1972
<i>Paratrechina fulva incisa</i> (Forel, 1908)	cr	Kempf 1972
<i>Paratrechina fulva longiscapa</i> (Forel, 1908)	br cr	Kempf 1972
<i>Paratrechina fulva nesiotis</i> (Wheeler W.M., 1919)	ec	Kempf 1972
<i>Paratrechina goeldii</i> Forel, 1912	br	Kempf 1972
<i>Paratrechina guatemalensis</i> (Forel, 1885)	gu me us	Kempf 1972; Trager 1984
<i>Paratrechina guatemalensis cocoensis</i> (Forel, 1902)	cr	Kempf 1972

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Paratrechina guatemalensis edenensis</i> (Wheeler W.M., 1924)	ec	Kempf 1972
<i>Paratrechina hystrix</i> Trager, 1984	us	Kempf 1972; Trager 1984
<i>Paratrechina johannae</i> (Forel, 1912)	co	Kempf 1972
<i>Paratrechina johannae dibullana</i> (Forel, 1912)	co	Kempf 1972
<i>Paratrechina kincaidi</i> (Wheeler W.M., 1906)	ber	Kempf 1972
<i>Paratrechina laevigata</i> MacKay, 1998	me	
<i>Paratrechina lietzi</i> (Forel, 1908)	gi gf su cr	Kempf 1972
<i>Paratrechina longicornis</i> Latreille, 1802	ch pe ec br gi gf su tt vn co pn cr ni be ho me us am an bh	Kempf 1972
<i>Paratrechina mexicana</i> (Forel, 1899)	me	Kempf 1972
<i>Paratrechina microps</i> (Smith M.R., 1937)	am	Kempf 1972
<i>Paratrechina myops</i> (Mann, 1920)	am	Kempf 1972
<i>Paratrechina nodifera</i> (Mayr, 1870)	bo co	Kempf 1972
<i>Paratrechina pearsei</i> (Wheeler W.M., 1938)	me	Kempf 1972
<i>Paratrechina pubens</i> (Forel, 1893)	an	Kempf 1972
<i>Paratrechina silvestrii</i> (Emery, 1906)	ar ur	Kempf 1972
<i>Paratrechina silvestrii kuenzleri</i> (Forel, 1909)	ar	Kempf 1972
<i>Paratrechina staudingeri</i> (Forel, 1912)	pe	Kempf 1972
<i>Paratrechina steinheili</i> (Forel, 1893)	br tt co cr ho gu me am an	Kempf 1972
<i>Paratrechina steinheili minuta</i> (Forel, 1893)	co am an	Kempf 1972
<i>Paratrechina tococae</i> (Wheeler W.M. & Bequaert, 1929)	br	Kempf 1972
<i>Paratrechina troglodytes</i> (Weber, 1934)	am	Kempf 1972
<i>Paratrechina vividula</i> (Nylander, 1846)	br tt vn us am an ber	Kempf 1972
<i>Paratrechina vividula antillana</i> (Forel, 1893)	am an bh	Kempf 1972
<i>Paratrechina vividula australis</i> (Santschi, 1929)	br	Kempf 1972
<i>Paratrechina vividula docilis</i> (Forel, 1908)	br	Kempf 1972
<i>Prenolepis acuminata</i> Forel, 1911	me	Brandao 1991
<i>Prenolepis imparis</i> Say, 1836	me us	
Subfamily Myrmeciinae		Clark 1951
<i>Insertae sedis</i>		
<i>Ameghinoia piatnitzkyi</i> * Viana & Haedo Rossi, 1957	ar	Ward & Brady 2003
<i>Polanskiella</i> * sp . Rossi de García, 1983		Ward & Brady 2003
Subfamily Pseudomyrmecinae		
Tribe Pseudomyrmecini		
<i>Myrcidris epicharis</i> Ward, 1990	br	Ward 1990
<i>Pseudomyrmex acanthobius</i> (Emery, 1896)	ar pr bo br	Ward 1990
<i>Pseudomyrmex acanthobius cocae</i> (Santschi, 1922)	ar	Ward 1990
<i>Pseudomyrmex acanthobius fuscatus</i> (Emery, 1896)	pr	Ward 1990
<i>Pseudomyrmex acanthobius virgo</i> (Santschi, 1922)	ar br	Ward 1990
<i>Pseudomyrmex adustus</i> (Borgmeier, 1929)	br	Ward 1990
<i>Pseudomyrmex alternans</i> (Santschi, 1936)	gf	Ward 1990
<i>Pseudomyrmex alustratus</i> Ward, 1989	pe	Ward 1989 1990
<i>Pseudomyrmex alvarengai</i> Kempf, 1961	br	Ward 1990
<i>Pseudomyrmex antigenus</i> (Enzmann, 1944)	gu	Ward 1990
<i>Pseudomyrmex antigenus brunneipes</i> (Enzmann, 1944)	gu	Ward 1990
<i>Pseudomyrmex antiquus</i> * Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex apache</i> Creighton, 1953	us	Ward 1990
<i>Pseudomyrmex atripes</i> (Smith F., 1860)	br co	Ward 1990
<i>Pseudomyrmex avitus</i> * Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex baros</i> * Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex beccarii</i> (Menozzi, 1935)	gi	Ward 1990
<i>Pseudomyrmex boopis</i> (Roger, 1863)	ec tt vn pn cr gu me	Ward 1990
<i>Pseudomyrmex browni</i> Kempf, 1967	br co pn	Ward 1990
<i>Pseudomyrmex brunneus</i> (F. Smith, 1877)	cr me us am	Ward 1990
<i>Pseudomyrmex caeciliae</i> (Forel, 1913)	pn cr ni ho gu me us am	Ward 1989 1990
<i>Pseudomyrmex championi</i> (Forel, 1899)	tt gu me	Ward 1990
<i>Pseudomyrmex cladoicus</i> (Smith F., 1858)	br	Ward 1990
<i>Pseudomyrmex colei</i> (Enzmann, 1944)	pe	Ward 1990
<i>Pseudomyrmex colei vistanus</i> (Enzmann, 1944)	pe	Ward 1990
<i>Pseudomyrmex concolor</i> (Smith F., 1860)	br gi gf su	Ward 1989 1990 1999

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pseudomyrmex coruscus</i> * Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex cretus</i> Ward, 1989	cr gu	Ward 1989 1990
<i>Pseudomyrmex crudelis</i> Ward, 1999	pe	Ward 1990 1999
<i>Pseudomyrmex cubensis</i> (Forel, 1901)	ar bo pe br vn co pn cr gu me us am bh	Ward 1989 1990
<i>Pseudomyrmex curacaensis</i> (Forel, 1912)	bo ec br gi gf su tt vn co pn an	Ward 1989 1990
<i>Pseudomyrmex deminutus</i> Ward, 1999	br	Ward 1990 1999
<i>Pseudomyrmex dendroicus</i> (Forel, 1904)	br	Ward 1989 1990 1999
<i>Pseudomyrmex denticollis</i> (Emery, 1890)	ar pr br	Ward 1990
<i>Pseudomyrmex depressus</i> (Forel, 1906)	co	Ward 1990
<i>Pseudomyrmex distinctus</i> (Smith F., 1877)	me	Ward 1990
<i>Pseudomyrmex duckei</i> (Forel, 1906)	br	Ward 1990
<i>Pseudomyrmex ecaleus</i> Ward, 1999	pe ec	Ward 1990 1999
<i>Pseudomyrmex eduardi</i> (Forel, 1912)	pr bo br vn co pn cr am	Ward 1989 1990
<i>Pseudomyrmex ejectus</i> (Smith F., 1858)	bo br cr me us am	Ward 1989 1990
<i>Pseudomyrmex elongatulus</i> (Dalle Torre, 1892)	pe br gi gf su tt vn co pn cr gu me us	Ward 1990
<i>Pseudomyrmex elongatulus longus</i> (Forel, 1912)	br	Ward 1990
<i>Pseudomyrmex elongatus</i> (Mayr, 1870)	pe br gi gf su tt vn co pn cr gu me us	Ward 1989 1990 1999
<i>Pseudomyrmex ethicus</i> (Forel, 1911)	br	Ward 1990
<i>Pseudomyrmex euryblemma</i> (Forel, 1899)	co cr	Ward 1989 1990
<i>Pseudomyrmex excisus</i> (Mayr, 1870)	co	Ward 1990
<i>Pseudomyrmex faber</i> (Smith F., 1858)	br gi gf su pn cr	Ward 1990
<i>Pseudomyrmex ferox</i> Ward, 1999	pe br co	Ward 1990 1999
<i>Pseudomyrmex ferrugineus</i> (Smith F., 1877)	cr ni be ho gu me	Ward 1989 1990
<i>Pseudomyrmex fervidus</i> (Smith F., 1877)	me	Ward 1990
<i>Pseudomyrmex fiebrigi</i> (Forel, 1908)	ar pr	Ward 1990
<i>Pseudomyrmex filiformis</i> (Fabricius, 1804)	br gi gf su tt vn co pn cr gu me	Ward 1989 1990
<i>Pseudomyrmex flavigaster</i> (Smith F., 1877)	cr ni	Ward 1989 1990
<i>Pseudomyrmex flavidulus</i> (Smith F., 1858)	ar bo br tt vn co pn cr gu me am an bh	Ward 1990
<i>Pseudomyrmex flavidulus lizeri</i> (Santschi, 1922)	bo	Ward 1990
<i>Pseudomyrmex fortis</i> (Forel, 1899)	me	Ward 1990 1999
<i>Pseudomyrmex gebelli</i> (Forel, 1899)	co pn	Ward 1990
<i>Pseudomyrmex gibbinotus</i> (Forel, 1908)	br	Ward 1990
<i>Pseudomyrmex godmani</i> (Forel, 1899)	br co pn	Ward 1990
<i>Pseudomyrmex goeldii</i> (Forel, 1912)	br	Ward 1990
<i>Pseudomyrmex gracilis</i> (Fabricius, 1804)	ar pr ur bo pe ec gi gf su tt vn co pn cr ni ho gu me am	Ward 1989 1990 1999
<i>Pseudomyrmex gracilis argentinus</i> (Santschi, 1934)	ar	Ward 1990
<i>Pseudomyrmex gracilis atrinodus</i> (Santschi, 1934)	ar pr br	Ward 1990
<i>Pseudomyrmex haytianus</i> (Forel, 1901)	am	Ward 1990
<i>Pseudomyrmex hesperius</i> Ward, 1993	me	Ward 1990
<i>Pseudomyrmex holmgreni</i> (Wheeler, 1925)	pe	Ward 1989 1990
<i>Pseudomyrmex hospitalis</i> Ward, 1999	pe	Ward 1990 1999
<i>Pseudomyrmex incurrens</i> (Forel, 1912)	br	Ward 1990
<i>Pseudomyrmex inquilinus</i> Ward, 1996	co	Ward 1990 1999
<i>Pseudomyrmex insuavis</i> Ward, 1999	pr br pn cr gu	Ward 1990 1999
<i>Pseudomyrmex ita</i> (Forel, 1906)	me	Ward 1990
<i>Pseudomyrmex janzeni</i> Ward, 1993	ar pr bo pe br tt pn cr ho me	Ward 1990 1999
<i>Pseudomyrmex kuenckeli</i> (Emery, 1890)	br	Ward 1989 1990
<i>Pseudomyrmex laevifrons</i> Ward, 1989	br gi gf su co cr	Ward 1990
<i>Pseudomyrmex laevigatus</i> (Smith F., 1877)	br co	Ward 1990
<i>Pseudomyrmex laevivertex</i> (Forel, 1906)	ch	Ward 1990
<i>Pseudomyrmex lynceus</i> (Spinola, 1851)	ch	Ward 1990
<i>Pseudomyrmex macrops</i> * Ward, 1992	ar pe br gi gf su tt	Ward 1990
<i>Pseudomyrmex maculatus</i> (Smith F., 1855)	me	Ward 1990
<i>Pseudomyrmex major</i> (Forel, 1899)	gi gf su tt	Ward 1989 1990 1999
<i>Pseudomyrmex malignus</i> (Wheeler, 1921)		

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Pseudomyrmex mandibularis</i> (Spinola, 1851)	br	Ward 1990
<i>Pseudomyrmex mixtecus</i> Ward, 1993	me	Ward 1990
<i>Pseudomyrmex monochrous</i> (Dalle Torre, 1892)	br	Ward 1990
<i>Pseudomyrmex mordax</i> (Warming, 1894)	vn co pn	Ward 1990 1999
<i>Pseudomyrmex nexilis*</i> Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex niger</i> (Donisthorpe, 1940)	gi	Ward 1990
<i>Pseudomyrmex nigrocinctus</i> (Emery, 1890)	gi gf su cr	Ward 1990
<i>Pseudomyrmex nigropilosus</i> (Emery, 1890)	cr	Ward 1990
<i>Pseudomyrmex oculatus</i> (Smith F., 1855)	bo pe ec br gi gf su vn co pn cr ni be ho gu me co pn cr	Ward 1990
<i>Pseudomyrmex oki</i> (Forel, 1906)	gu	Ward 1990
<i>Pseudomyrmex opaciceps</i> Ward, 1993	ad	Ward 1990
<i>Pseudomyrmex oryzetus*</i> Ward, 1992	pn cr	Ward 1989 1990
<i>Pseudomyrmex osurus</i> (Forel, 1911)	pe vn co	Ward 1990
<i>Pseudomyrmex pallens</i> (Mayr, 1870)	co	Ward 1990
<i>Pseudomyrmex pallens landolti</i> (Forel, 1912)	ar pr ur bo br tt co cr be ho gu me us am	Ward 1990
<i>Pseudomyrmex pallidus</i> (Smith F., 1855)	cr	Ward 1990
<i>Pseudomyrmex particeps</i> Ward, 1993	am	Ward 1989 1990
<i>Pseudomyrmex pazosi</i> (Santschi, 1909)	ve su br gi	Ward 1990 1999
<i>Pseudomyrmex penetrator</i> (Smith F., 1877)	gu	Ward 1990
<i>Pseudomyrmex peperi</i> (Forel, 1913)	pe br tt cr me	Ward 1989 1990
<i>Pseudomyrmex perboscii</i> (Guérin-Méneville, 1844)	pe	Ward 1990
<i>Pseudomyrmex peruvianus</i> (Wheeler W.M., 1925)	ar pr ur br tt	Ward 1990
<i>Pseudomyrmex phyllophilus</i> (Smith F., 1858)	br	Ward 1990 1999
<i>Pseudomyrmex pictus</i> (Stitz, 1913)	pe br	Ward 1989 1990
<i>Pseudomyrmex pisinnus</i> Ward, 1989	ad	Ward 1990
<i>Pseudomyrmex prioris*</i> Ward, 1992	br	Ward 1990
<i>Pseudomyrmex pupa</i> (Forel, 1911)	ni	Ward 1990
<i>Pseudomyrmex reconditus</i> Ward, 1993	br	Ward 1990
<i>Pseudomyrmex rochai</i> (Forel, 1912)	br	Ward 1990
<i>Pseudomyrmex rubiginosus</i> (Stitz, 1913)	br	Ward 1990 1999
<i>Pseudomyrmex rufiventris</i> (Forel, 1911)	pr	Ward 1989 1990
<i>Pseudomyrmex rufomedius</i> (Smith F., 1877)	gu	Ward 1989 1990
<i>Pseudomyrmex salvini</i> (Forel, 1899)	gu me	Ward 1990
<i>Pseudomyrmex santschii</i> (Enzmann, 1944)	pe	Ward 1990
<i>Pseudomyrmex satanicus</i> (Wheeler W.M., 1942)	pn	Ward 1990
<i>Pseudomyrmex schuppi</i> (Forel, 1901)	br	Ward 1989 1990
<i>Pseudomyrmex seminole</i> Ward, 1985	me us	Ward 1990
<i>Pseudomyrmex sericeus</i> (Mayr, 1870)	br co pn cr be gu	Ward 1990
<i>Pseudomyrmex sericeus cordiae</i> (Forel, 1904)	bo pe	Ward 1990
<i>Pseudomyrmex sericeus lisa</i> (Enzmann, 1944)	pe	Ward 1990
<i>Pseudomyrmex sericeus longior</i> (Forel, 1904)	pe	Ward 1990
<i>Pseudomyrmex simplex</i> (Smith F., 1877)	pe ec br tt co pn cr es be ho gu us am bh	Ward 1989 1990
<i>Pseudomyrmex simulans</i> Kempf, 1958	pn	Ward 1990
<i>Pseudomyrmex solisi</i> (Santschi, 1916)	ar	Ward 1990
<i>Pseudomyrmex solisi belgranoi</i> (Santschi, 1916)	ar	Ward 1990
<i>Pseudomyrmex spiculus</i> Ward, 1989	pe br gi gf su co pn cr me	Ward 1989 1990
<i>Pseudomyrmex spinicola</i> (Emery, 1890)	pn cr ni be me	Ward 1989 1990
<i>Pseudomyrmex squamifer</i> (Emery, 1890)	br	Ward 1990
<i>Pseudomyrmex subater</i> (Wheeler W.M. & Mann, 1914)	am	Ward 1990
<i>Pseudomyrmex subtilissimus</i> (Emery, 1890)	cr ni me	Ward 1989 1990
<i>Pseudomyrmex succinus*</i> Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex tachigaliae</i> (Forel, 1904)	pe br	Ward 1989 1990 1999
<i>Pseudomyrmex tenuis</i> (Fabricius, 1804)	bo pe br gi gf su co cr	Ward 1989 1990
<i>Pseudomyrmex tenuissimus</i> (Emery, 1906)	bo pe ec br gi gf su tt vn co pn cr ni es be ho me am	Ward 1989 1990
<i>Pseudomyrmex terminalis</i> (Smith F., 1877)	br	Ward 1990
<i>Pseudomyrmex termitarius</i> (Smith F., 1855)	pr bo pe br gi gf su tt vn co pn an	Ward 1990

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pseudomyrmex thecolor*</i> Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex triplaridis</i> (Forel, 1904)	br gi gf su tt vn co pn cr gu me	Ward 1989 1990 1999
<i>Pseudomyrmex triplarinus</i> (Weddell, 1850)	bo pe ec br	Ward 1990 1999
<i>Pseudomyrmex ultrix</i> Ward, 1999	ec	Ward 1990 1999
<i>Pseudomyrmex unicolor</i> (Smith F., 1855)	pr pe br gi gf su	Ward 1990
<i>Pseudomyrmex urbanus</i> (Smith F., 1877)	ar pr bo pe ec br gi gf su tt vn co pn	Ward 1989 1990
<i>Pseudomyrmex veneficus</i> (Wheeler, 1942)	me	Ward 1990
<i>Pseudomyrmex venustus</i> (Smith F., 1858)	pe br co pn	Ward 1990
<i>Pseudomyrmex vicinus*</i> Ward, 1992	ad	Ward 1990
<i>Pseudomyrmex viduus</i> (Smith F., 1858)	br gi gf su	Ward 1989 1990 1999
<i>Pseudomyrmex villosus</i> Ward, 1989	br	Ward 1989 1990
<i>Pseudomyrmex vinneni</i> (Forel, 1906)	br	Ward 1990 1999
<i>Pseudomyrmex vitabilis</i> Ward, 1999	co	Ward 1990 1999
<i>Pseudomyrmex vojtoskii</i> (Enzmann, 1944)	pe	Ward 1990
<i>Pseudomyrmex vojtoskii costaricensis</i> (Enzmann, 1944)	cr	Ward 1990
<i>Pseudomyrmex weberi</i> (Enzmann, 1944)	gu	Ward 1990
<i>Pseudomyrmex wheeleri</i> (Enzmann, 1944)	pe	Ward 1990
Subfamily Cerapachyinae		
Tribe Acanthostichini		
<i>Acanthostichus bentoni</i> MacKay, 1996	br	MacKay 1996
<i>Acanthostichus brevicornis</i> Emery, 1894	ar br gi gf su	MacKay 1996
<i>Acanthostichus brevinodis</i> MacKay, 1996	br	MacKay 1996
<i>Acanthostichus concavinodis</i> MacKay, 1996	bo pe	MacKay 1996
<i>Acanthostichus emmae</i> MacKay, 1996	me	MacKay 1996
<i>Acanthostichus femoralis</i> Kusnezov, 1962	ar br	MacKay 1996
<i>Acanthostichus flexuosus</i> MacKay, 1996	br	MacKay 1996
<i>Acanthostichus hispaniolicus</i> de Andrade 1998	ad	MacKay 1996
<i>Acanthostichus kirbyi</i> Emery, 1895	ar pr bo pe br	MacKay 1996
<i>Acanthostichus laevigatus</i> MacKay, 1996	vn	MacKay 1996
<i>Acanthostichus laticornis</i> Forel, 1908	ar pr ur br	MacKay 1996
<i>Acanthostichus lattkei</i> MacKay, 1996	vn	MacKay 1996
<i>Acanthostichus punctiscapus</i> MacKay, 1996	me	MacKay 1996
<i>Acanthostichus quadratus</i> Emery, 1895	ar bo pe ec br	MacKay 1996
<i>Acanthostichus quirozi</i> MacKay, 1996	me	MacKay 1996
<i>Acanthostichus sanchezorum</i> MacKay, 1985	co	MacKay 1996
<i>Acanthostichus serratulus</i> (Smith, 1858)	ar br	MacKay 1996
<i>Acanthostichus skwarrae</i> Wheeler W.M., 1934	me	MacKay 1996
<i>Acanthostichus texanus</i> Forel, 1904	me us	MacKay 1996
<i>Acanthostichus truncatus</i> MacKay, 1996	co	MacKay 1996
Tribe Cerapachyini		
<i>Cerapachys biroi</i> Forel, 1907	am	Brown 1975
<i>Cerapachys hondurianus</i> Mann, 1922	be ho	Brown 1975
<i>Cerapachys neotropicus</i> Weber, 1939	ec	Brown 1975
<i>Cerapachys splendens</i> Borgmeier, 1957	br	Brown 1975
<i>Cerapachys toltecus</i> Forel, 1909	gu	Brown 1975
<i>Sphinctomyrmex stali</i> Mayr, 1866	br	Brown 1975
Tribe Cylindromyrmeini		
<i>Cylindromyrmex antillanus*</i> de Andrade, 1998	ad	De Andrade, 1998
<i>Cylindromyrmex boliviiae</i> Wheeler W.M., 1924	bo pe vn co	De Andrade, 1998
<i>Cylindromyrmex brasiliensis</i> Emery, 1901	pr br	De Andrade, 1998
<i>Cylindromyrmex brevitarsus</i> Santschi, 1925	pe ec br vn cr	De Andrade, 1998
<i>Cylindromyrmex darlingtoni</i> Wheeler W.M., 1937	am	De Andrade, 1998
<i>Cylindromyrmex electronius*</i> de Andrade, 1998	ad	De Andrade, 1998
<i>Cylindromyrmex escobari de</i> Andrade, 1998	co cr	De Andrade, 1998
<i>Cylindromyrmex godmani</i> Forel, 1899	ec co pn cr	De Andrade, 1998
<i>Cylindromyrmex inopinatus*</i> de Andrade, 2001	ad	De Andrade, 1998 2001
<i>Cylindromyrmex longiceps</i> André, 1892	br	De Andrade, 1998
<i>Cylindromyrmex meinerti</i> Forel, 1905	br vn pn cr	De Andrade, 1998
<i>Cylindromyrmex striatus</i> Mayr, 1870	br gi gf su	De Andrade, 1998

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Cylindromyrmex whymperi</i> Cameron (1891)	ch bo pe ec br co cr gu	De Andrade, 1998
Subfamily Ecitoninae		
Tribe Cheliomyrmecini		
<i>Cheliomyrmex andicola</i> Emery, 1894	pe br co	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Cheliomyrmex audax</i> Santschi, 1921	ec co	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Cheliomyrmex megalonyx</i> Wheeler W.M., 1921	gi gf su vn	Borgmeier 1955; Watkins 1976
<i>Cheliomyrmex morosus</i> (Smith F., 1859)	gi gf su be ho me	Borgmeier 1955; Watkins 1976; Palacio 1999
Tribe Ecitonini		
<i>Ecton burchellii</i> (Westwood, 1842)	pr ur bo ec br tt vn co pn cr ni ho gu me	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton drepanophorum</i> Smith F., 1858	bo pe ec br gi gf su	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton dulcius</i> Forel, 1912	ar pr br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton hamatum</i> (Fabricius, 1782)	bo pe ec br gi gf su co pn cr ni be me	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton jansoni</i> Forel, 1912	ec co pn cr ni	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton lucanoides</i> Emery, 1894	bo pe br co pn cr ni	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton mexicanum</i> Roger, 1863	ar pr bo pe br co pn cr be gu me	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton quadriglume</i> (Haliday, 1836)	ar pr pe br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton rapax</i> Smith F., 1855	bo pe ec br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Ecton setigaster</i> Borgmeier, 1953	bo br	Borgmeier 1955; Watkins 1976
<i>Ecton uncinatum</i> Borgmeier, 1953	ec	Borgmeier 1955; Watkins 1976
<i>Ecton vagans</i> (Olivier, 1792)	br gi gf su vn cr gu me	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Labidus auropubens</i> (Santschi, 1920)	gf	Borgmeier 1955; Watkins 1976
<i>Labidus coecus</i> (Latreille, 1802)	ar pr ur bo pe ec br gi gf su vn co pn cr ni be ho gu me us	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Labidus curvipes</i> (Emery, 1900)	ec cr	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Labidus mars</i> (Forel, 1912)	br	Borgmeier 1955; Watkins 1976
<i>Labidus nero</i> (Santschi, 1930)	br	Borgmeier 1955; Watkins 1976
<i>Labidus praedator</i> (Smith F., 1858)	ar pr bo pe br co pn cr be ho gu me	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Labidus spininodis</i> (Emery, 1890)	br co cr	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Labidus truncatidens</i> (Santschi, 1920)	br gi gf su	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex adnepos</i> (Wheeler W.M., 1922)	ec br tt cr	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex agilis</i> Borgmeier, 1953	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex alfaroi</i> (Emery, 1890)	cr	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex andrei</i> (Emery, 1901)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex angulimandibulatus</i> Watkins, 1974	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex angustinodis</i> (Emery, 1888)	ar pr br gi gf su	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex antillanus</i> (Forel, 1897)	an	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex asper</i> Borgmeier, 1955	cr	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex balzani</i> (Emery, 1894)	bo	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex bohlisi</i> (Emery, 1896)	ar pr pe br gi gf su	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex bruchi</i> (Forel, 1912)	ar	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex carettei</i> (Forel, 1913)	ar	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex carinifrons</i> Borgmeier, 1953	br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex chameleensis</i> Watkins, 1986	me	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex clavifemur</i> Borgmeier, 1953	pe br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex cloosae</i> (Forel, 1912)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex compressinodis</i> Borgmeier, 1953	cr	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex cornutus</i> Watkins, 1975	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex crassiscapus</i> Watkins, 1990	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex cratensis</i> Borgmeier, 1953	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex cristatus</i> (André, 1889)	ec br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex curvinotus</i> Watkins, 1994	pe	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex densespunctatus</i> (Borgmeier, 1933)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex detectus</i> Borgmeier, 1953	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex diabolus</i> (Forel, 1912)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex diana</i> (Forel, 1912)	ec br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex digitistipus</i> Watkins, 1975	cr	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex diversinodis</i> (Borgmeier, 1933)	ar pr bo br	Borgmeier 1955; Watkins 1976

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Neivamyrmex dorbignii</i> (Shuckard, 1840)	ar ur br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex ectopus*</i> Wilson, 1985	ad	Bolton 1995
<i>Neivamyrmex emersoni</i> (Wheeler W.M., 1921)	ec br gi gf su tt co	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex emeryi</i> (Santschi, 1921)	bo pe	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex erichsonii</i> (Westwood, 1842)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex falciferus</i> (Emery, 1900)	bo pe ec br gi gf su	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex foveolatus</i> Borgmeier, 1953	pn	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex fumosus</i> (Forel, 1913)	cr gu me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex fuscipennis</i> (Smith M.R., 1942)	us	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex genalis</i> Borgmeier, 1953	bo	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex gibbatus</i> Borgmeier, 1953	ec br gi gf su co pn cr	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex goeldii</i> (Forel, 1901)	ar br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex gracillae</i> (Mann, 1926)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex gracilis</i> Borgmeier, 1955	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex gradualis</i> Borgmeier, 1953	bo	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex guerinii</i> (Shuckard, 1840)	br gi gf su	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex guyanensis</i> (Santschi, 1916)	gi gf su vn co pn	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex halidayi</i> (Shuckard, 1840)	ar pr bo ec br gi gf su tt	Borgmeier 1955; Watkins 1976; Palacio 1999
	vn co pn cr ni ho me	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex harrisi</i> (Haldeman, 1852)	me us	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex hetschkoii</i> (Mayr, 1886)	ar pr ur br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex hopei</i> (Shuckard, 1840)	ar bo br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex humilis</i> (Borgmeier, 1939)	cr	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex iheringi</i> (Forel, 1908)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex imbellis</i> (Emery, 1900)	pe	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex inca</i> (Santschi, 1921)	pe	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex inflatus</i> Borgmeier, 1958	me	Watkins 1976
<i>Neivamyrmex iridescens</i> Borgmeier, 1950	gi gf su	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex jermannii</i> Forel, 1901	ar pr ur bo br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex klugii</i> Shuckard, 1840	an	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex laevigatus</i> (Borgmeier, 1948)	ar pr br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex laticapus</i> (Emery, 1901)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex legionis</i> (Smith F., 1855)	ar pr br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex leonardi</i> (Wheeler W.M., 1915)	us	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex leptognathus</i> (Emery, 1900)	bo br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex lieselae</i> (Forel, 1913)	ar	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex longiscapus</i> Borgmeier, 1953	pn cr gu me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex macrodentatus</i> (Menozzi, 1931)	cr	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex macropterus</i> Borgmeier, 1953	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex manni</i> (Wheeler W.M., 1914)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex maxillosus</i> (Emery, 1900)	br gi gf su	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex megatrichia</i> Kempf, 1961	ec gi gf su	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex melanocephalus</i> (Emery, 1895)	gu me us	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex melshaemeri</i> (Haldeman, 1852)	cr gu me us	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex micans</i> Borgmeier, 1953	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex minensis</i> (Borgmeier, 1928)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex modestus</i> (Borgmeier, 1933)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex mojave</i> Smith M.R., 1943	us	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex nigrescens</i> (Cresson, 1872)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex nordenskioldii</i> (Holmgren, 1908)	pe	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex nyensis</i> Watkins, 1977	us	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex opacithorax</i> (Emery, 1894)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex orthonotus</i> (Borgmeier, 1933)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex pacificus</i> Borgmeier, 1955	pe	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex pauxillus</i> (Wheeler W.M., 1903)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex perplexus</i> Borgmeier, 1953	br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex pertii</i> (Shuckard, 1840)	ar ch pr ur bo br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex physognathus</i> (Emery, 1900)	bo	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex pilosus</i> (Smith F., 1858)	ar pe br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex piraticus</i> Borgmeier, 1953	br	Borgmeier 1955; Watkins 1976

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Neivamyrmex planidens</i> Borgmeier, 1953	ec	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex planidorsus</i> (Emery, 1906)	pr br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex postangustatus</i> (Borgmeier, 1934)	br gi gf su	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex postcarinatus</i> Borgmeier, 1953	pn	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex pseudops</i> (Forel, 1909)	ar pr bo br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex puerulus</i> Borgmeier, 1955	pn	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex pulchellus</i> Borgmeier, 1955	pn	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex pullus</i> Borgmeier, 1953	pn	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex punctaticeps</i> (Emery, 1894)	ar pr bo br gi gf su tt co cr me es	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex quadraooccipitus</i> Watkins, 1975	pe	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex radoszkowskii</i> (Emery, 1900)	ar br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex raptor</i> (Forel, 1911)	ar pr bo br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex shuckardi</i> (Shuckard, 1840)	ec co pn gu	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex rosenbergi</i> (Forel, 1911)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex rugulosus</i> Borgmeier, 1953	pn	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex scutellaris</i> Borgmeier, 1953	ar pr bo br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex shuckardi</i> (Emery, 1900)	pn cr	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex spatulatus</i> (Borgmeier, 1939)	co	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex spinolai</i> (Westwood, 1843)	pn cr me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex spoliator</i> (Forel, 1899)	ar ur bo br	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex sulcatus</i> (Mayr, 1868)	ni gu me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex sumichrasti</i> (Norton, 1868)	ar pr ec br vn co pn cr me us	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Neivamyrmex swainsonii</i> (Shuckard, 1840)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex tenuis</i> (Borgmeier, 1953)	me	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex texanus</i> Watkins, 1972	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex tristis</i> (Forel, 1901)	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex vicinus</i> Borgmeier, 1953	br	Borgmeier 1955; Watkins 1976
<i>Neivamyrmex walkeri</i> (Westwood, 1842)	ar bo pe br gi gf su	Borgmeier 1955; Watkins 1976
<i>Nomamyrmex esenbeckii</i> (Westwood, 1842)	ar pr br gi gf su	Borgmeier 1955; Watkins 1976; Palacio 1999
<i>Nomamyrmex hartigii</i> (Westwood, 1842)	br gi gf su tt co pn	Borgmeier 1955; Watkins 1976; Palacio 1999
Subfamily Leptanilloidinae		
<i>Asphinctanilloides amazona</i> Brandão, Diniz, Agosti & Delabie, 1999	br	Brandão, Diniz, Agosti & Delabie, 1999
<i>Asphinctanilloides ana</i> Brandão, Diniz, Agosti & Delabie, 1999	br	Brandão, Diniz, Agosti & Delabie, 1999
<i>Asphinctanilloides manauara</i> Brandão, Diniz, Agosti & Delabie, 1999	br	Brandão, Diniz, Agosti & Delabie, 1999
<i>Leptanilloides biconstricta</i> Mann, 1923	bo	Brandão, Diniz, Agosti & Delabie, 1999
<i>Leptanilloides improvisa</i> Brandão, Diniz, Agosti & Delabie, 1999	ec	Brandão, Diniz, Agosti & Delabie, 1999
<i>Leptanilloides legionaria</i> Brandão, Diniz, Agosti & Delabie, 1999	co	Brandão, Diniz, Agosti & Delabie, 1999
<i>Leptanilloides mckennae</i> Longino, 2003	cr	Longino 2003b
<i>Leptanilloides sculpturata</i> Brandão, Diniz, Agosti & Delabie, 1999	co	Brandão, Diniz, Agosti & Delabie, 1999
Subfamily Amblyoponinae		
Tribe Amblyoponini		
<i>Amblyopone agostii</i> Lacau & Delabie, 2002	br	Lacau & Delabie 2002
<i>Amblyopone armigera</i> Mayr, 1887	ar br	Lattke 1991
<i>Amblyopone bierigi</i> (Santschi, 1930)	am	Lattke 1991
<i>Amblyopone chilensis</i> Mayr, 1887	ch	Lattke 1991
<i>Amblyopone cleiae</i> Delabie, 2002	br	Lacau & Delabie 2002
<i>Amblyopone degenerata</i> Borgmeier, 1957	br su	Lattke 1991
<i>Amblyopone egregia</i> (Kusnezov, 1955)	ar	Lattke 1991
<i>Amblyopone elongata</i> (Santschi, 1912)	ar ur br	Lattke 1991
<i>Amblyopone falcata</i> Lattke, 1991	am	Lattke 1991
<i>Amblyopone heraldoi</i> Lacau & Delabie, 2002	br	Lacau & Delabie 2002
<i>Amblyopone lurilabes</i> Lattke, 1991	br vn	Lattke 1991
<i>Amblyopone monrosi</i> Brown, 1960	ch	Lattke 1991
<i>Amblyopone mystriops</i> Brown, 1960	gu	Lattke 1991

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Amblyopone orizabana</i> Brown, 1960	me	Lattke 1991
<i>Paraprisopelta minima</i> Kusnezov, 1955	ar	Bolton 1995
<i>Prionopelta amabilis</i> Borgmeier, 1949	cr	Brown 1960
<i>Prionopelta antillana</i> Forel, 1909	bo br gi gf su tt pn cr ni ho gu an	Brown 1960
<i>Prionopelta marthae</i> Forel, 1909	vn	Brown 1960
<i>Prionopelta modesta</i> Forel, 1909	gu me	Brown 1960
<i>Prionopelta punctulata</i> Mayr, 1866	ar br	Brown 1960
Subfamily Ponerinae		
Tribe Ponerini		
<i>Anochetus altisquamis</i> Mayr, 1887	ar ur br	Brown 1978
<i>Anochetus ambiguus*</i> de Andrade, 1994	ad	Bolton 1995
<i>Anochetus bispinosus</i> (Smith F., 1858)	bo br gi gf su co	Brown 1978
<i>Anochetus brevidentatus*</i> MacKay, 1991	ad	Bolton 1995
<i>Anochetus conisquamis*</i> de Andrade, 1994	ad	Bolton 1995
<i>Anochetus corayi*</i> Baroni Urbani, 1980	ad	Bolton 1995
<i>Anochetus diegensis</i> Forel, 1912	br co pn	Brown 1978
<i>Anochetus dubius*</i> de Andrade, 1994	ad	Bolton 1995
<i>Anochetus elegans</i> Lattke, 1987	co	Bolton 1995
<i>Anochetus emarginatus</i> (Fabricius, 1804)	gi gf su tt vn co an	Brown 1978
<i>Anochetus extinctus*</i> de Andrade, 1994	ad	Bolton 1995
<i>Anochetus haytianus</i> Wheeler W.M. y Mann, 1914	am	Brown 1978
<i>Anochetus horridus</i> Kempf, 1964	br	Brown 1978
<i>Anochetus inca</i> Wheeler W.M., 1925	pe	Brown 1978
<i>Anochetus inermis</i> André, 1889	tt vn co an	Brown 1978
<i>Anochetus intermedius*</i> de Andrade, 1994	ad	Bolton 1995
<i>Anochetus kempfi</i> Brown, 1978	am	Brown 1978
<i>Anochetus longispinus</i> Wheeler W.M., 1936	am	Brown 1978
<i>Anochetus lucidus*</i> de Andrade, 1994	ad	Bolton 1995
<i>Anochetus micans</i> Brown, 1978	cr be ho bh	Brown 1978
<i>Anochetus minans</i> Mann, 1922	cr ho	Brown 1978
<i>Anochetus neglectus</i> Emery, 1894	ar ur br	Brown 1978
<i>Anochetus orchidicola</i> Brown, 1978	me	Brown 1978
<i>Anochetus oriens</i> Kempf, 1964	br	Brown 1978
<i>Anochetus simoni</i> Emery, 1890	ec gi gf su vn co	Brown 1978
<i>Anochetus striatulus</i> Emery, 1890	vn cr	Brown 1978
<i>Anochetus targioni</i> Emery, 1894	bo ec br tt	Brown 1978
<i>Anochetus testaceus</i> Forel, 1893	be an bh	Brown 1978
<i>Anochetus vallensis</i> Lattke, 1987	co	Brown 1978
<i>Anochetus vexator</i> Kempf, 1964	br	Brown 1978
<i>Belonopelta attenuata</i> Mayr, 1870	co	Baroni Urbani 1975
<i>Belonopelta deletrix</i> Mann, 1922	ho me	Baroni Urbani 1975
<i>Centromyrmex alfaroi</i> Emery, 1890	bo br cr	Kempf 1967
<i>Centromyrmex brachycola</i> (Roger, 1861)	bo br	Kempf 1967
<i>Centromyrmex gigas</i> Forel, 1911	ar br	Kempf 1967
<i>Cryptopone guatemalensis</i> (Forel, 1899)	ni gu	Bolton 1995
<i>Dinoponera australis</i> Emery, 1901	ar pr ur bo br	Kempf 1971
<i>Dinoponera australis bucki</i> Borgmeier, 1937	br	Kempf 1971
<i>Dinoponera australis nigricolor</i> Borgmeier, 1937	br	Kempf 1971
<i>Dinoponera gigantea</i> (Perty, 1833)	pe br	Kempf 1971
<i>Dinoponera longipes</i> Emery, 1901	pe br	Kempf 1971
<i>Dinoponera lucida</i> Emery, 1901	br	Kempf 1971
<i>Dinoponera mutica</i> Emery, 1901	bo br	Kempf 1971
<i>Dinoponera quadriceps</i> Kempf, 1971	br	Kempf 1971
<i>Hypoponera agilis</i> (Borgmeier, 1934)	su	Kempf 1972; Bolton 1995
<i>Hypoponera aliena</i> (Smith F., 1858)	br	Kempf 1972; Bolton 1995
<i>Hypoponera argentina</i> (Santschi, 1922)	ar	Kempf 1972; Bolton 1995
<i>Hypoponera beebei</i> (Wheeler W.M., 1924)	ec	Kempf 1972; Bolton 1995
<i>Hypoponera clavatula</i> (Emery, 1906)	ar	Kempf 1972; Bolton 1995
<i>Hypoponera collegiana</i> (Santschi, 1925)	br	Kempf 1972; Bolton 1995

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Hypoponera collegiana paranensis</i> Santschi, 1925	br	Kempf 1972; Bolton 1995
<i>Hypoponera creola</i> (Menozzi, 1931)	cr	Kempf 1972; Bolton 1995
<i>Hypoponera distinguenda</i> (Emery, 1890)	pr bo br vn co	Kempf 1972; Bolton 1995
<i>Hypoponera distinguenda dispar</i> (Santschi, 1925)	br	Kempf 1972; Bolton 1995
<i>Hypoponera distinguenda histrio</i> (Forel, 1912)	ar br	Kempf 1972; Bolton 1995
<i>Hypoponera distinguenda inexpedita</i> (Forel, 1911)	br	Kempf 1972; Bolton 1995
<i>Hypoponera distinguenda vana</i> (Forel, 1909)	gu	Kempf 1972; Bolton 1995
<i>Hypoponera ergatandria</i> (Forel, 1893)	pr cr me am an bh	Kempf 1972; Bolton 1995
<i>Hypoponera faceta</i> (Menozzi, 1931)	cr	Kempf 1972; Bolton 1995
<i>Hypoponera fenestralis</i> (Gallardo, 1918)	ar	Kempf 1972; Bolton 1995
<i>Hypoponera fiebrigi antoniensis</i> (Forel, 1912)	co	Kempf 1972; Bolton 1995
<i>Hypoponera fiebrigi famini</i> (Forel, 1912)	co	Kempf 1972; Bolton 1995
<i>Hypoponera fiebrigi transiens</i> (Santschi, 1925)	ar	Kempf 1972; Bolton 1995
<i>Hypoponera foeda sarolae</i> (Forel, 1912)	br	Kempf 1972; Bolton 1995
<i>Hypoponera foreli</i> (Mayr, 1887)	br	Kempf 1972; Bolton 1995
<i>Hypoponera gracilicornis</i> (Menozzi, 1931)	cr	Kempf 1972; Bolton 1995
<i>Hypoponera ideletiae</i> (Santschi, 1923)	br	Kempf 1972; Bolton 1995
<i>Hypoponera ignigera</i> (Menozzi, 1927)	cr	Kempf 1972; Bolton 1995
<i>Hypoponera inexorata</i> (Wheeler W.M., 1903)	cr me us	Kempf 1972; Bolton 1995
<i>Hypoponera inexorata fallax</i> (Forel, 1909)	gu	Kempf 1972; Bolton 1995
<i>Hypoponera iheringi</i> (Forel, 1908)	br	Kempf 1972; Bolton 1995
<i>Hypoponera leninei</i> (Santschi, 1925)	br	Kempf 1972; Bolton 1995
<i>Hypoponera menozzii</i> (Santschi, 1932)	cr	Kempf 1972; Bolton 1995
<i>Hypoponera neglecta</i> (Santschi, 1923)	br	Kempf 1972; Bolton 1995
<i>Hypoponera nitidula</i> (Emery, 1890)	cr ho me	Kempf 1972; Bolton 1995
<i>Hypoponera opaciceps</i> (Mayr, 1887)	ar ur pe br gi tt gu me us am an bh am	Kempf 1972; Bolton 1995
<i>Hypoponera opaciceps cubana</i> (Santschi, 1930)	co	Kempf 1972; Bolton 1995
<i>Hypoponera opaciceps gaigei</i> (Forel, 1914)	am	Kempf 1972; Bolton 1995
<i>Hypoponera opaciceps jamaicensis</i> (Aguayo, 1932)	ar	Kempf 1972; Bolton 1995
<i>Hypoponera opaciceps pampana</i> (Santschi, 1925)	ar	Kempf 1972; Bolton 1995
<i>Hypoponera opaciceps postangustata</i> (Forel, 1908)	pr	Kempf 1972; Bolton 1995
<i>Hypoponera opacioc</i> (Forel, 1893)	ar ch tt me am an	Kempf 1972; Bolton 1995
<i>Hypoponera parva</i> (Forel, 1909)	gu	Kempf 1972; Bolton 1995
<i>Hypoponera perplexa</i> (Mann, 1922)	pn cr ho	Kempf 1972; Bolton 1995
<i>Hypoponera punctatissima</i> (Roger, 1859)	tt am	Kempf 1972; Bolton 1995
<i>Hypoponera reichenbergeri</i> (Santschi, 1923)	br	Kempf 1972; Bolton 1995
<i>Hypoponera schmalzi</i> (Emery, 1896)	br	Kempf 1972; Bolton 1995
<i>Hypoponera schmalzi fugitans</i> (Forel, 1912)	br	Kempf 1972; Bolton 1995
<i>Hypoponera schmalzi paulina</i> (Forel, 1913)	br	Kempf 1972; Bolton 1995
<i>Hypoponera schwebeli</i> (Forel, 1913)	br	Kempf 1972; Bolton 1995
<i>Hypoponera stoica</i> (Santschi, 1912)	ur	Kempf 1972; Bolton 1995
<i>Hypoponera trigona</i> (Mayr, 1887)	ar pr bo br cr	Kempf 1972; Bolton 1995
<i>Hypoponera trigona cauta</i> (Forel, 1912)	br	Kempf 1972; Bolton 1995
<i>Hypoponera vernacula</i> (Kempf, 1962)	br	Kempf 1972; Bolton 1995
<i>Hypoponera viri</i> (Santschi, 1923)	br	Kempf 1972; Bolton 1995
<i>Hypoponera wilsoni</i> (Santschi, 1924)	pr br	Kempf 1972; Bolton 1995
<i>Leptogenys amazonica</i> Borgmeier, 1930	br	Kempf 1972; Bolton 1995
<i>Leptogenys ambigua</i> Santschi, 1931	pn	Kempf 1972; Bolton 1995
<i>Leptogenys analecti</i> Borgmeier, 1930	br	Kempf 1972; Bolton 1995
<i>Leptogenys antennata</i> Wheeler W.M. & Mann, 1914	am	Kempf 1972; Bolton 1995
<i>Leptogenys arcuata</i> Roger, 1861	gi gf su	Kempf 1972; Bolton 1995
<i>Leptogenys arcuata deletangi</i> Santschi, 1921	bo	Kempf 1972; Bolton 1995
<i>Leptogenys australis</i> (Emery, 1888)	ar ur	Kempf 1972; Bolton 1995
<i>Leptogenys bohlsi</i> Emery, 1896	pr br	Kempf 1972; Bolton 1995
<i>Leptogenys bohlsi weiseri</i> Santschi, 1925	ar	Kempf 1972; Bolton 1995
<i>Leptogenys consanguinea</i> Wheeler W.M., 1909	me	Kempf 1972; Bolton 1995
<i>Leptogenys crudelis</i> (Smith F., 1858)	br	Kempf 1972; Bolton 1995
<i>Leptogenys dasygyna</i> Wheeler W.M., 1923	gi gf su	Kempf 1972; Bolton 1995
<i>Leptogenys donisthorpei</i> Mann, 1922	ho	Kempf 1972; Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Leptogenys famelica</i> Emery, 1896	cr	Kempf 1972; Bolton 1995
<i>Leptogenys gaigei</i> Wheeler W.M., 1923	gi gf su	Kempf 1972; Bolton 1995
<i>Leptogenys gaigei defuga</i> Wheeler, 1923	gi gf su	Kempf 1972; Bolton 1995
<i>Leptogenys gagates</i> Mann, 1922	ho	Kempf 1972; Bolton 1995
<i>Leptogenys guianensis</i> Wheeler W.M., 1923	gi	Kempf 1972; Bolton 1995
<i>Leptogenys hansenii</i> Borgmeier, 1930	br	Kempf 1972; Bolton 1995
<i>Leptogenys honduriana</i> Mann, 1922	ho	Kempf 1972; Bolton 1995
<i>Leptogenys iheringi</i> Forel, 1911	br	Kempf 1972; Bolton 1995
<i>Leptogenys imperatrix</i> Mann, 1922	ho	Kempf 1972; Bolton 1995
<i>Leptogenys ingens</i> Mayr, 1866	co	Kempf 1972; Bolton 1995
<i>Leptogenys langi</i> Wheeler, 1923	gi gf su	Kempf 1972; Bolton 1995
<i>Leptogenys linearis</i> (Smith F., 1858)	br	Kempf 1972; Bolton 1995
<i>Leptogenys luederwaldti</i> Forel, 1913	br	Kempf 1972; Bolton 1995
<i>Leptogenys maxillosa</i> (Smith F., 1858)	an	Kempf 1972; Bolton 1995
<i>Leptogenys maxillosa falcata</i> Roger, 1861	br am	Kempf 1972; Bolton 1995
<i>Leptogenys melzeri</i> Borgmeier, 1930	br	Kempf 1972; Bolton 1995
<i>Leptogenys mexicana</i> Mayr, 1870	me	Kempf 1972; Bolton 1995
<i>Leptogenys mucronata</i> Forel, 1893	an	Kempf 1972; Bolton 1995
<i>Leptogenys mucronata columbica</i> Forel, 1901	co	Kempf 1972; Bolton 1995
<i>Leptogenys peninsularis</i> Mann, 1926	me	Kempf 1972; Bolton 1995
<i>Leptogenys pubiceps</i> Emery, 1890	vn co	Kempf 1972; Bolton 1995
<i>Leptogenys pubiceps cubensis</i> Santschi, 1930	am	Kempf 1972; Bolton 1995
<i>Leptogenys pubiceps vincentensis</i> Forel, 1901	an	Kempf 1972; Bolton 1995
<i>Leptogenys punctaticeps</i> Emery, 1890	cr am an	Kempf 1972; Bolton 1995
<i>Leptogenys pusilla</i> (Emery, 1890)	co cr	Kempf 1972; Bolton 1995
<i>Leptogenys pusilla panamana</i> Wheeler W.M., 1923	pn	Kempf 1972; Bolton 1995
<i>Leptogenys quiriguana</i> Wheeler W.M., 1923	gu	Kempf 1972; Bolton 1995
<i>Leptogenys ritae</i> Forel, 1899	pn	Kempf 1972; Bolton 1995
<i>Leptogenys rufa</i> Mann, 1922	ho	Kempf 1972; Bolton 1995
<i>Leptogenys unistimulosa</i> Roger, 1863	br gi gf su	Kempf 1972; Bolton 1995
<i>Leptogenys unistimulosa bahiana</i> Santschi, 1929	br	Kempf 1972; Bolton 1995
<i>Leptogenys unistimulosa trinidadensis</i> Forel, 1901	tt vn	Kempf 1972; Bolton 1995
<i>Leptogenys venatrix</i> Forel, 1899	pn	Kempf 1972; Bolton 1995
<i>Leptogenys vogeli</i> Borgmeier, 1933	br	Kempf 1972; Bolton 1995
<i>Leptogenys wheeleri</i> Forel, 1901	me	Kempf 1972; Bolton 1995
<i>Odontomachus affinis</i> Guérin-Méneville, 1844	br	Brown 1976
<i>Odontomachus allolabis</i> Kempf, 1974	ec br co	Brown 1976
<i>Odontomachus bauri</i> Emery, 1892	pr bo pe ec br vn co pn cr am	Brown 1976
<i>Odontomachus biolleyi</i> Forel, 1908	cr	Brown 1976
<i>Odontomachus biumbonatus</i> Brown, 1976	pe ec br gi gf su	Brown 1976
<i>Odontomachus bradleyi</i> Brown, 1976	pe ec	Brown 1976
<i>Odontomachus brunneus</i> (Patton, 1894)	br	Brown 1976
<i>Odontomachus caelalus</i> Brown, 1976	pe ec gi gf su	Brown 1976
<i>Odontomachus chelifer</i> (Latreille, 1802)	ar pr bo pe br gi gf su	Brown 1976
	vn co pn cr ho me	Brown 1976
<i>Odontomachus clarus</i> Roger, 1861	me	Brown 1976
<i>Odontomachus cornutus</i> Stitz, 1933	ec	Brown 1976
<i>Odontomachus erythrocephalus</i> Emery, 1890	co pn cr	Brown 1976
<i>Odontomachus haematodus</i> (Linnaeus, 1758)	ar bo pe ec br gi gf su tt vn co	Brown 1976
	pn cr ni ho gu me am an bh	Brown 1976
<i>Odontomachus insularis</i> Guérin-Méneville, 1844	am an	Brown 1976
<i>Odontomachus hastatus</i> (Fabricius, 1804)	pe br gi gf su co pn cr	Brown 1976
<i>Odontomachus laticeps</i> Roger, 1861	bo br gi gf su cr ho gu me	Brown 1976
<i>Odontomachus mayi</i> Mann, 1912	br gi gf su	Brown 1976
<i>Odontomachus meinerti</i> Forel, 1905	pr br vn	Brown 1976
<i>Odontomachus mormo</i> Brown, 1976	ec	Brown 1976
<i>Odontomachus opaciventris</i> Forel, 1899	ec br co pn cr gu me	Brown 1976
<i>Odontomachus panamensis</i> Forel, 1899	br co pn	Brown 1976
<i>Odontomachus ruginodis</i> Smith M.R., 1937	bh	Brown 1976
<i>Odontomachus sculptus</i> Brown, 1978	ec gi gf su	Bolton 1995

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Odontomachus spissus</i> Kempf, 1962	br	Brown 1976
<i>Odontomachus yucatecus</i> Brown, 1976	be gu me	Brown 1976
<i>Pachycondyla aenescens</i> (Mayr, 1870)	co cr	Kempf 1972; Bolton 1995
<i>Pachycondyla apicalis</i> (Latreille, 1802)	pe ec br gi gf su tt vn co pn cr ho gu me	Kempf 1972; Bolton 1995
<i>Pachycondyla arhuaca</i> Forel, 1901	bo br gi gf su vn co pn	Kempf 1972; Bolton 1995
<i>Pachycondyla atrovirens</i> (Mayr, 1866)	ec co	Kempf 1972; Bolton 1995
<i>Pachycondyla bucki</i> (Borgmeier, 1927)	br	Kempf 1972; Bolton 1995
<i>Pachycondyla bugabensis</i> (Forel, 1899)	pn	Kempf 1972; Bolton 1995
<i>Pachycondyla carbonaria</i> (Smith F., 1858)	ec	Kempf 1972; Bolton 1995
<i>Pachycondyla cauta</i> Mann 1922	br ho	Kempf 1972; Bolton 1995
<i>Pachycondyla carinulata</i> (Roger, 1861)	br gi gf su co pn cr	Kempf 1972; Bolton 1995
<i>Pachycondyla cavinodis</i> Mann, 1916	br	Kempf 1972; Bolton 1995
<i>Pachycondyla chyzeri</i> Forel, 1907	co	Kempf 1972; Bolton 1995
<i>Pachycondyla cognata</i> Emery 1896	cr	Kempf 1972; Bolton 1995
<i>Pachycondyla commutata</i> Roger, 1860	pe ec br gi gf su vn co	Kempf 1972; Bolton 1995
<i>Pachycondyla constricta</i> (Mayr, 1884)	pe br gi gf su tt co pn cr ho	Kempf 1972; Bolton 1995
<i>Pachycondyla crassinoda</i> (Latreille, 1802)	br gi gf su tt vn	Kempf 1972; Bolton 1995
<i>Pachycondyla crenata</i> (Roger, 1861)	ar pr bo br gi gf su vn co cr ho me	Kempf 1972; Bolton 1995
<i>Pachycondyla eleonorae</i> Forel, 1921	ec	Kempf 1972; Bolton 1995
<i>Pachycondyla emiliae</i> Forel, 1901	vn	Kempf 1972; Bolton 1995
<i>Pachycondyla fauveli</i> (Emery, 1896)	bo pe ec co	Kempf 1972; Bolton 1995
<i>Pachycondyla ferruginea</i> (Smith F., 1858)	me	Kempf 1972; Bolton 1995
<i>Pachycondyla foetida</i> (Olivier, 1791)	br gi gf su co pn cr	Kempf 1972; Bolton 1995
<i>Pachycondyla fuscoatra</i> Roger, 1861	bo ec br vn co cr	Kempf 1972; Bolton 1995
<i>Pachycondyla gilberti</i> Kempf 1960	br su	Kempf 1972; Bolton 1995
<i>Pachycondyla goeldii</i> Forel, 1912	br	Kempf 1972; Bolton 1995
<i>Pachycondyla guianensis</i> Weber, 1939	gi	Kempf 1972; Bolton 1995
<i>Pachycondyla haskinsi</i> Weber, 1939	br co pn	Kempf 1972; Bolton 1995
<i>Pachycondyla harpax</i> (Fabricius, 1804)	pr bo ec br gi gf su tt vn co pn cr ni be ho gu me us am pe br gi gf su	Kempf 1972; Bolton 1995
<i>Pachycondyla holmgreni</i> Wheeler W.M., 1925	bo pe ec br gi gf su tt vn cr ec br gi gf su tt co pn cr	Kempf 1972; Bolton 1995
<i>Pachycondyla impressa</i> Roger, 1861	br	Kempf 1972; Bolton 1995
<i>Pachycondyla laevigata</i> (Smith F., 1858)	vn	Kempf 1972; Bolton 1995
<i>Pachycondyla lenis</i> Kempf 1962	pn cr me	Kempf 1972; Bolton 1995
<i>Pachycondyla lenkoi</i> Kempf, 1962	bo	Kempf 1972; Bolton 1995
<i>Pachycondyla leveillei</i> Emery 1890	br	Kempf 1972; Bolton 1995
<i>Pachycondyla lineaticeps</i> (Mayr, 1866)	br	Kempf 1972; Bolton 1995
<i>Pachycondyla luteola</i> (Roger, 1861)	br	Kempf 1972; Bolton 1995
<i>Pachycondyla marginata</i> Roger 1861	br	Kempf 1972; Bolton 1995
<i>Pachycondyla magnifica</i> Borgmeier, 1929	br	Kempf 1972; Bolton 1995
<i>Pachycondyla mesonotalis</i> Santschi, 1923	br	Kempf 1972; Bolton 1995
<i>Pachycondyla metanotalis</i> Luederwaldt, 1918	br	Kempf 1972; Bolton 1995
<i>Pachycondyla oberthueri</i> (Emery, 1890)	bo br	Kempf 1972; Bolton 1995
<i>Pachycondyla obscuricornis</i> (Emery, 1890)	br gi gf su pn cr me	Kempf 1972; Bolton 1995
<i>Pachycondyla obsoleta</i> Menozzi 1931	cr	Kempf 1972; Bolton 1995
<i>Pachycondyla procidua</i> Emery, 1890	br gi gf su	Kempf 1972; Bolton 1995
<i>Pachycondyla rostrata</i> (Emery, 1890)	vn	Kempf 1972; Bolton 1995
<i>Pachycondyla stigma</i> (Fabricius, 1804)	ar bo pe ec br gi gf su tt co pn cr be ho gu am an bh ar pr ur br	Kempf 1972; Bolton 1995
<i>Pachycondyla striata</i> Smith F., 1858	pe br co pn cr gu	Kempf 1972; Bolton 1995
<i>Pachycondyla striatinodis</i> (Emery, 1890)	am	Kempf 1972; Bolton 1995
<i>Pachycondyla succedanea</i> Roger 1963	pe br pn	Kempf 1972; Bolton 1995
<i>Pachycondyla theresiae</i> (Forel, 1899)	pe ec br gi gf su tt	Kempf 1972; Bolton 1995
<i>Pachycondyla unidentata</i> (Mayr, 1862)	co pn cr ho gu me	Kempf 1972; Bolton 1995
<i>Pachycondyla venusta</i> Forel, 1912	br	Kempf 1972; Bolton 1995
<i>Pachycondyla villosa</i> (Fabricius, 1804)	ar pr bo pe ec br gi gf su vn co pn cr ho me	Kempf 1972; Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Simopelta bicolor</i> (Borgmeier, 1950)	br	Kempf 1972; Bolton 1995
<i>Simopelta curvata</i> Mayr, 1887	br	Kempf 1972; Bolton 1995
<i>Simopelta jeckylli</i> Mann, 1916	br	Kempf 1972; Bolton 1995
<i>Simopelta laticeps</i> (Gotwald & Brown, 1967)	pe	Kempf 1972; Bolton 1995
<i>Simopelta manni</i> (Wheeler W.M., 1935)	ec	Kempf 1972; Bolton 1995
<i>Simopelta minima</i> Brandão, 1989	br	Bolton 1995
<i>Simopelta oculata</i> (Gotwald y Brown, 1967)	cr	Kempf 1972; Bolton 1995
<i>Simopelta paeminosa</i> (Snelling, 1971)	cr	Bolton 1995
<i>Simopelta pergandei</i> Forel, 1909	cr gu	Kempf 1972; Bolton 1995
<i>Simopelta williamsi</i> (Wheeler W.M., 1935)	ec	Kempf 1972; Bolton 1995
Tribe Thaumatomyrmecini		
<i>Thaumatomyrmex atrox</i> Weber, 1939	gi	Kempf 1975; Longino 1988
<i>Thaumatomyrmex baryai</i> Fontenla, 1995		
<i>Thaumatomyrmex cochlearis</i> Creighton, 1928	am	Kempf 1975; Longino 1988
<i>Thaumatomyrmex contumax</i> Kempf, 1975	br	Kempf 1975; Longino 1988
<i>Thaumatomyrmex ferox</i> Mann, 1922	ho	Kempf 1975; Longino 1988
<i>Thaumatomyrmex mutilatus</i> Mayr, 1887	br	Kempf 1975; Longino 1988
Tribe Platythyreini		
<i>Platythyrea angusta</i> Forel, 1901	pe br gi gf su vn	Brown 1975
<i>Platythyrea exigua</i> Kempf, 1964	br	Brown 1975
<i>Platythyrea pilosula</i> (Smith F., 1858)	br vn co	Brown 1975
<i>Platythyrea prizo</i> Kugler, 1977	cr	Brown 1975
<i>Platythyrea punctata</i> (Smith F., 1858)	br gi gf su co pn ni ho gu me us am an bh bo pe br gi gf su vn	Brown 1975
<i>Platythyrea sinuata</i> (Roger, 1860)	am	Brown 1975
<i>Platythyrea strenua</i> Wheeler W.M. y Mann, 1914	ec	Brown 1975
<i>Platythyrea zodion</i> Brown, 1975		
Subfamily Ectatomminae		
Tribe Ectatommini		
<i>Ectatomma confine</i> Mayr, 1870	br co pn ho	Kugler & Brown 1982
<i>Ectatomma edentatum</i> Roger, 1863	ar pr ur bo pe br gi gf su vn co pn cr	Kugler & Brown 1982
<i>Ectatomma gibbum</i> Kugler & Brown, 1982	cr ni ho	Kugler & Brown 1982
<i>Ectatomma gonionion</i> Kugler & Brown, 1982	ec co	Kugler & Brown 1982
<i>Ectatomma lugens</i> Emery, 1894	br gi gf su	Kugler & Brown 1982
<i>Ectatomma muticum</i> Mayr, 1870	br	Kugler & Brown 1982
<i>Ectatomma opaciventre</i> Roger, 1861	ar pr br	Kugler & Brown 1982
<i>Ectatomma permagnum</i> Forel, 1908	ar pr bo br	Kugler & Brown 1982
<i>Ectatomma planidens</i> Borgmeier, 1939	br	Kugler & Brown 1982
<i>Ectatomma quadridens</i> (Fabricius, 1793)	ar pr bo pe ec br gi gf su co pn	Kugler & Brown 1982
<i>Ectatomma ruidum</i> Roger, 1860	br gi gf su tt vn co cr ni ho me	Kugler & Brown 1982
<i>Ectatomma suzanae</i> Almeida, 1986	br	Bolton 1995
<i>Ectatomma tuberculatum</i> (Olivier, 1792)	ar pr br gi gf su tt vn co pn cr ho gu me	Kugler & Brown 1982
<i>Gnamptogenys acuminata</i> Emery, 1896	bo br gi gf su vn	Lattke 1995
<i>Gnamptogenys acuta</i> (Brown, 1956)	bo pe	Lattke 1995
<i>Gnamptogenys alfaroi</i> Emery, 1894	cr	Lattke 1995
<i>Gnamptogenys ammophila</i> Lattke, 1990	vn	Lattke 1995
<i>Gnamptogenys andina</i> Lattke, 1995	co ec	Lattke 1995
<i>Gnamptogenys annulata</i> Mayr, 1887	bo pe ec br gi gf su vn co pn ho	Lattke 1995
<i>Gnamptogenys banksii</i> (Wheeler W.M., 1930)	pn	Lattke 1995
<i>Gnamptogenys biquetra</i> Lattke, 2002	pn	Lattke 2002
<i>Gnamptogenys bispinosa</i> (Emery, 1890)	pn cr	Lattke 1995
<i>Gnamptogenys bisulca</i> Kempf y Brown, 1968	co	Lattke 1995
<i>Gnamptogenys bruchi</i> (Santschi, 1922)	ar br	Lattke 1995
<i>Gnamptogenys boliviensis</i> Lattke, 1995	bo	Lattke 1995
<i>Gnamptogenys brunnea</i> Lattke, 1995	co	Lattke 1995
<i>Gnamptogenys brunoi*</i> Lattke, 2002	ad	Lattke 2002
<i>Gnamptogenys bufonis</i> (Mann, 1926)	me	Lattke 1995
<i>Gnamptogenys caelata</i> Kempf, 1967	pr br	Lattke 1995

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Gnamptogenys casca*</i> Lattke, 2002	ad	Lattke 2002
<i>Gnamptogenys concinna</i> (Smith F., 1858)	bo pe br tt vn co pn cr	Lattke 1995
<i>Gnamptogenys continua</i> Mayr, 1887	bo pe br vn pn me am	Lattke 1995
<i>Gnamptogenys cuneiforma</i> Lattke, 1995	pn	Lattke 1995
<i>Gnamptogenys curvoclypeata</i> Lattke, 1990	vn	Lattke 1995
<i>Gnamptogenys ejuncida</i> Lattke, 1995	co	Lattke 1995
<i>Gnamptogenys ericae</i> Forel, 1912	co	Lattke 1995
<i>Gnamptogenys extra</i> Lattke, 1995	co ec	Lattke 1995
<i>Gnamptogenys falcaria</i> Lattke, 2002	ad	Lattke 2002
<i>Gnamptogenys falcifera</i> Kempf, 1967	pe vn	Lattke 1995
<i>Gnamptogenys fernandezi</i> Lattke, 1990	vn	Lattke 1995
<i>Gnamptogenys fieldi</i> Lattke, 1990	vn	Lattke 1995
<i>Gnamptogenys gentryi</i> Lattke, 1995	co	Lattke 1995
<i>Gnamptogenys gracilis</i> (Santschi, 1929)	gi gf su vn	Lattke 1995
<i>Gnamptogenys laticephalia</i> Lattke, 1995	ec	Lattke 1995
<i>Gnamptogenys haenschei</i> (Emery, 1902)	bo pe ec br cr	Lattke 1995
<i>Gnamptogenys hartmani</i> (Wheeler W.M., 1915)	ho us	Lattke 1995
<i>Gnamptogenys haytiana</i> (Wheeler W.M. & Mann, 1914)	am	Lattke 1995
<i>Gnamptogenys horni</i> Santschi, 1929	bo br gi gf su tt vn co pn	Lattke 1995
<i>Gnamptogenys ilimani</i> Lattke, 1995	bo	Lattke 1995
<i>Gnamptogenys ingeborgae</i> Brown, 1993	co	Lattke 1995
<i>Gnamptogenys insularis</i> Lattke, 2002	am	Lattke 2002
<i>Gnamptogenys interrupta</i> Mayr, 1887	br vn ho me	Lattke 1995
<i>Gnamptogenys kempfi</i> Lenko, 1964	br	Lattke 1995
<i>Gnamptogenys lanei</i> Kempf, 1960	br	Lattke 1995
<i>Gnamptogenys lavra</i> Lattke, 2002	br	Lattke 2002
<i>Gnamptogenys levinates*</i> Baroni Urbani, 1980	ad	Lattke 2002
<i>Gnamptogenys lineolata</i> Brown, 1993	an	Lattke 1995
<i>Gnamptogenys lucaris</i> Kempf, 1968	br	Lattke 1995
<i>Gnamptogenys mecotyle</i> Brown, 1958	bo br	Lattke 1995
<i>Gnamptogenys mediatrix</i> Brown, 1958	br	Lattke 1995
<i>Gnamptogenys menozzii</i> (Borgmeier, 1928)	br	Lattke 1995
<i>Gnamptogenys mina</i> (Brown, 1956)	bo pe vn	Lattke 1995
<i>Gnamptogenys minuta</i> (Emery, 1896)	bo br gi gf su co pn cr ho	Lattke 1995
<i>Gnamptogenys moelleri</i> (Forel, 1912)	br	Lattke 1995
<i>Gnamptogenys mordax</i> (Smith F., 1858)	ar br gi gf su co pn cr ho	Lattke 1995
<i>Gnamptogenys nana</i> Kempf, 1960	ar br	Lattke 1995
<i>Gnamptogenys nigrivitrea</i> Lattke, 1995	co	Lattke 1995
<i>Gnamptogenys perspicax</i> Kempf & Brown, 1970	co	Lattke 1995
<i>Gnamptogenys petiscapa</i> Lattke, 1990	vn	Lattke 1995
<i>Gnamptogenys pilosa</i> Lattke, 1995	co	Lattke 1995
<i>Gnamptogenys pittieri</i> Lattke, 1990	vn	Lattke 1995
<i>Gnamptogenys pleurodon</i> (Emery, 1896)	bo pe br tt vn	Lattke 1995
<i>Gnamptogenys porcata</i> (Emery, 1896)	vn cr	Lattke 1995
<i>Gnamptogenys pristina*</i> Baroni Urbani, 1980	ad	Lattke 2002
<i>Gnamptogenys rastrata</i> (Mayr, 1866)	br	Lattke 1995
<i>Gnamptogenys regularis</i> Mayr, 1870	pr pe ec br gi gf su vn pn cr ho me	Lattke 1995
<i>Gnamptogenys reichenbergeri</i> (Santschi, 1929)	br	Lattke 1995
<i>Gnamptogenys relicta</i> (Mann, 1916)	br	Lattke 1995
<i>Gnamptogenys rimulosa</i> (Roger, 1861)	br	Lattke 1995
<i>Gnamptogenys rumba</i> Lattke, 2002	am	Lattke 2002
<i>Gnamptogenys schmitti</i> (Forel, 1901)	am	Lattke 1995
<i>Gnamptogenys semiferox</i> Brown, 1958	am	Lattke 1995
<i>Gnamptogenys siapensis</i> Lattke, 1995	ve	Lattke 1995
<i>Gnamptogenys simplicoides</i> (Forel, 1908)	br	Lattke 1995
<i>Gnamptogenys simulans</i> (Emery, 1896)	cr	Lattke 1995
<i>Gnamptogenys stellae</i> Lattke, 1995	cr	Lattke 1995
<i>Gnamptogenys striatula</i> Mayr, 1884	ar pr ur bo br gi gf su tt am an	Lattke 1995
<i>Gnamptogenys strigata</i> (Norton, 1871)	gi gf su co pn cr ho gu me	Lattke 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Gnamptogenys striolata</i> (Borgmeier, 1957)	br	Lattke 1995
<i>Gnamptogenys sulcata</i> (Smith F., 1858)	pr br gi gf su vn pn cr	Lattke 1995
<i>Gnamptogenys tortuolosa</i> (Smith F., 1858)	pe ec br gi gf su vn	Lattke 1995
<i>Gnamptogenys transversa</i> Lattke, 1995	pn	Lattke 1995
<i>Gnamptogenys triangularis</i> Mayr, 1887	ar bo br	Lattke 1995
<i>Gnamptogenys volcano</i> Lattke, 1995	cr	Lattke 1995
<i>Gnamptogenys vriesi</i> Brandão & Lattke, 1990	ec	Lattke 1995
Tribe Typhlomyrmecini		
<i>Typhlomyrmex bruchi</i> Santschi, 1922	ar	Brown 1965
<i>Typhlomyrmex clavicornis</i> Emery, 1906	pr bo br gi gf su	Brown 1965
<i>Typhlomyrmex foreli</i> Santschi, 1925	br	Brown 1965
<i>Typhlomyrmex major</i> Santschi, 1923	br	Brown 1965
<i>Typhlomyrmex prolatus</i> Brown, 1965	cr	Brown 1965
<i>Typhlomyrmex pusillus</i> Emery, 1894	pr bo br cr ho	Brown 1965
<i>Typhlomyrmex rogenhoferi</i> Mayr, 1862	ar bo pe gi gf su vn co pn cr ho gu me	Brown 1965
Subfamily Heteroponerinae		
Tribe Heteroponerini		
<i>Acanthoponera goeldii</i> Forel, 1912	br	Brown 1958
<i>Acanthoponera minor</i> Forel, 1899	ec co cr gu me	Brown 1958
<i>Acanthoponera mucronata</i> (Roger, 1860)	ar bo br vn	Brown 1958
<i>Acanthoponera peruviana</i> Brown, 1958	pe	Brown 1958
<i>Heteroponera angulata</i> Borgmeier, 1959	br	Brown 1958; Kempf 1962
<i>Heteroponera carinifrons</i> Mayr, 1887	ch	Brown 1958; Kempf 1962
<i>Heteroponera dentinodis</i> (Mayr, 1887)	bo br	Brown 1958; Kempf 1962
<i>Heteroponera dolo</i> (Roger, 1861)	ar pr ur br	Brown 1958; Kempf 1962
<i>Heteroponera flava</i> Kempf, 1962	br	Brown 1958; Kempf 1962
<i>Heteroponera georgesii</i> Perrault, 1998	gf	Perrault 1998
<i>Heteroponera inca</i> Brown, 1958	co	Brown 1958; Kempf 1962
<i>Heteroponera inermis</i> (Emery, 1894)	br	Brown 1958; Kempf 1962
<i>Heteroponera mayri</i> Kempf, 1962	br	Brown 1958; Kempf 1962
<i>Heteroponera microps</i> Borgmeier, 1957	br vn co	Brown 1958; Kempf 1962
<i>Heteroponera monticola</i> Kempf & Brown, 1970	co	Bolton 1995
<i>Heteroponera panamensis</i> (Forel, 1899)	pn	Brown 1958; Kempf 1962
<i>Heteroponera robusta</i> Kempf, 1962	br	Brown 1958; Kempf 1962
Subfamily Paraponerinae		
Tribe Paraponerini		
<i>Paraponera clavata</i> (Fabricius, 1775)	pr bo pe ec br gi gf su co pn cr ni	Brown 1958
<i>Paraponera dieteri*</i> Baroni Urbani, 1994	ad	Bolton 1995
Subfamily Proceratiinae		
Tribe Proceratiini		
<i>Discothyrea denticulata</i> Weber, 1939	gi gf su	Bolton 1995
<i>Discothyrea horni</i> Menozzi, 1927	cr	Bolton 1995
<i>Discothyrea humilis</i> Weber, 1939	pn	Bolton 1995
<i>Discothyrea icta</i> Weber, 1939	tt	Bolton 1995
<i>Discothyrea isthmica</i> Weber, 1940	pn	Bolton 1995
<i>Discothyrea neotropica</i> Bruch, 1919	ar	Bolton 1995
<i>Discothyrea sexarticulata</i> Borgmeier, 1954	br	Bolton 1995
<i>Discothyrea testacea</i> Roger, 1863	us	Bolton 1995
<i>Proceratium brasiliense</i> Borgmeier, 1959	br	Baroni Urbani & de Andrade 2003
<i>Proceratium catio</i> de Andrade, 2003	co	Baroni Urbani & de Andrade 2003
<i>Proceratium colombicum</i> de Andrade, 2003	co	Baroni Urbani & de Andrade 2003
<i>Proceratium compitale</i> de Andrade, 2003	me	Baroni Urbani & de Andrade 2003
<i>Proceratium conviciceps</i> Borgmeier, 1957	me pn	Baroni Urbani & de Andrade 2003
<i>Proceratium cubanum</i> de Andrade, 2003	am	Baroni Urbani & de Andrade 2003
<i>Proceratium denticulatum*</i> Lattke, 1991	ad	Baroni Urbani & de Andrade 2003
<i>Proceratium dominicanum*</i> de Andrade, 2003	ad	Baroni Urbani & de Andrade 2003
<i>Proceratium ecuadoriense</i> de Andrade, 2003	ec	Baroni Urbani & de Andrade 2003
<i>Proceratium gibberum*</i> de Andrade, 2003	ad	Baroni Urbani & de Andrade 2003

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Proceratium goliath</i> Kempf & Brown, 1968	cr	Baroni Urbani & de Andrade 2003
<i>Proceratium lattkei</i> de Andrade, 2003	ve	Baroni Urbani & de Andrade 2003
<i>Proceratium longiscapus</i> de Andrade, 2003	am	Baroni Urbani & de Andrade 2003
<i>Proceratium mancum</i> Mann, 1922	ho	Baroni Urbani & de Andrade 2003
<i>Proceratium mexicanum</i> de Andrade, 2003	me	Baroni Urbani & de Andrade 2003
<i>Proceratium micrommatum</i> (Roger, 1863)	pn ho me co vn	Baroni Urbani & de Andrade 2003
<i>Proceratium panamense</i> de Andrade, 2003	pn	Baroni Urbani & de Andrade 2003
<i>Proceratium poinari</i> * Baroni Urbani & de Andrade, 2003	ad	Baroni Urbani & de Andrade 2003
<i>Proceratium taino</i> de Andrade, 2003	am	Baroni Urbani & de Andrade 2003
<i>Proceratium tio</i> Snelling & Cover, 1992	me	Baroni Urbani & de Andrade 2003
<i>Proceratium transitionis</i> de Andrade, 2003	co	Baroni Urbani & de Andrade 2003
Tribe Probolomyrmecini		
<i>Probolomyrmex boliviensis</i> Mann, 1923	bo pe co pn	Agosti 1995
<i>Probolomyrmex brujitae</i> Agosti, 1994	ar	Agosti 1995
<i>Probolomyrmex guanacastensis</i> Agosti, 1997	cr	Agosti 1995
<i>Probolomyrmex petiolatus</i> Weber, 1940	vn co pn	Agosti 1995
Subfamily Agroecomyrmecinae		
Tribe Agroecomyrmecini		
<i>Tatuidris tatusia</i> Brown & Kempf, 1968	es	Brown & Kempf, 1968
Subfamily Myrmicinae		
Tribe Adelomyrmecini		
<i>Adelomyrmex betoi</i> Fernández, 2003	me	Fernández 2003b
<i>Adelomyrmex brevispinosus</i> Fernández, 2003	cr me	Fernández 2003b
<i>Adelomyrmex costatus</i> Fernández, 2003	co	Fernández 2003b
<i>Adelomyrmex cristiani</i> Fernández, 2003	co	Fernández 2003b
<i>Adelomyrmex foveolatus</i> Fernández, 2003	cr	Fernández 2003b
<i>Adelomyrmex grandis</i> Fernández, 2003	co	Fernández 2003b
<i>Adelomyrmex laevigatus</i> MacKay, 2003	pn cr	Fernández 2003b
<i>Adelomyrmex mackayi</i> Fernández, 2003	me	Fernández 2003b
<i>Adelomyrmex micans</i> Fernández, 2003	me	Fernández 2003b
<i>Adelomyrmex microps</i> Fernández, 2003	cr	Fernández 2003b
<i>Adelomyrmex minimus</i> Fernández, 2003	cr	Fernández 2003b
<i>Adelomyrmex myops</i> (Wheeler W.M., 1910)	pn cr ho gu	Fernández 2003b
<i>Adelomyrmex robustus</i> Fernández, 2003	me	Fernández 2003b
<i>Adelomyrmex silvestrii</i> (Menozzi, 1931)	cr gu me	Fernández 2003b
<i>Adelomyrmex striatus</i> Fernández, 2003	br	Fernández 2003b
<i>Adelomyrmex tristani</i> (Menozzi, 1931)	cr gu me	Fernández 2003b
<i>Adelomyrmex vaderi</i> Fernández, 2003	co	Fernández 2003b
Tribe Attini		
<i>Acromyrmex ambiguus</i> Emery, 1888	ar ur	Gonçalves 1961; Bolton 1995
<i>Acromyrmex aspersus</i> (Smith F., 1858)	ar pe br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex aspersus fuhrmanni</i> Forel, 1914	co	Gonçalves 1961; Bolton 1995
<i>Acromyrmex balzani</i> Emery, 1890	br	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Acromyrmex coronatus</i> (Fabricius, 1804)	bo pe br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex coronatus andicola</i> Emery, 1922	bo pe ec	Gonçalves 1961; Bolton 1995
<i>Acromyrmex coronatus globoculis</i> Kempf, 1972	gi gf su	Gonçalves 1961; Bolton 1995
<i>Acromyrmex coronatus importunus</i> Santschi, 1925	cr	Gonçalves 1961; Bolton 1995
<i>Acromyrmex coronatus panamensis</i> Forel, 1899	pe co pn cr	Gonçalves 1961; Bolton 1995
<i>Acromyrmex coronatus angustata</i> Forel, 1908	cr	Gonçalves 1961; Bolton 1995
<i>Acromyrmex coronatus rectispinus</i> Forel, 1908	cr	Gonçalves 1961; Bolton 1995
<i>Acromyrmex crassispinus</i> Forel, 1909	ar pr br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex disciger</i> (Mayr, 1887)	pr br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex echinatior</i> Forel, 1899	ec co pn cr gu me	Gonçalves 1961; Bolton 1995
<i>Acromyrmex fracticornis</i> Forel, 1909	ar pr ur br	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Acromyrmex heyeri</i> (Forel, 1899)	ar pr ur br	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Acromyrmex hispidus</i> Santschi, 1925	ur bo	Gonçalves 1961; Bolton 1995
<i>Acromyrmex hispidus fallax</i> Santschi, 1925	ar pr br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex hispidus formosus</i> Santschi, 1925	ar br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex hystrix</i> (Latreille, 1802)	ur br gi gf su	Gonçalves 1961; Bolton 1995
<i>Acromyrmex hystrix ajax</i> Forel, 1909	br	Gonçalves 1961; Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Acromyrmex insinuator</i> Schultz, Bekkevold & Boomsma, 1998	pn	Schultz, Bekkevold & Boomsma, 1998
<i>Acromyrmex laticeps</i> Emery, 1905	ar pr ur bo br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex laticeps nigrosetosus</i> Forel, 1908	bo br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex landolti</i> Forel, 1885	ar bo br vn co	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Acromyrmex lobicornis</i> Emery, 1888	ar pr ur bo br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lobicornis cochlearis</i> Santschi, 1933	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lobicornis ferrugineus</i> Emery, 1905	ar pr	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lobicornis penceensis</i> Forel, 1914	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lobicornis pruinosior</i> Santschi, 1916	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii</i> (Guérin-Meneville, 1838)	ar pr ur br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii parallelus</i> Santschi, 1916	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii boliviensis</i> Emery, 1905	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii boliviensis corallina</i> Santschi, 1919	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii boliviensis nigripes</i> Santschi, 1929	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii carli</i> Gonçalves, 1961	br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii decolor</i> Emery, 1905	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii pubescens</i> Emery, 1905	ar pr ur	Gonçalves 1961; Bolton 1995
<i>Acromyrmex lundii pubescens chacoensis</i> Santschi, 1916	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex niger</i> (Smith F., 1858)	br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex nobilis</i> Santschi, 1939	br vn	Gonçalves 1961; Bolton 1995
<i>Acromyrmex octospinosus</i> (Reich, 1793)	br gi gf su tt vn me	Gonçalves 1961; Bolton 1995
<i>Acromyrmex octospinosus cubanus</i> Wheeler W.M., 1937	am	Gonçalves 1961; Bolton 1995
<i>Acromyrmex octospinosus echinatior</i> Forel, 1899	ec co pn cr gu me	Gonçalves 1961; Bolton 1995
<i>Acromyrmex octospinosus ekchua</i> Wheeler W.M., 1937	me	Gonçalves 1961; Bolton 1995
<i>Acromyrmex octospinosus inti</i> Wheeler W.M., 1937	pe	Gonçalves 1961; Bolton 1995
<i>Acromyrmex octospinosus volcanus</i> Wheeler W.M., 1937	cr	Gonçalves 1961; Bolton 1995
<i>Acromyrmex pulvereus</i> Santschi, 1919	ar	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Acromyrmex rugosus</i> (Smith F., 1858)	ar ur bo pe br co	Gonçalves 1961; Bolton 1995
<i>Acromyrmex rugosus rochai</i> Forel, 1904	br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex rugosus santschii</i> Forel, 1912	co	Gonçalves 1961; Bolton 1995
<i>Acromyrmex silvestrii</i> Emery, 1905	ar ur	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Acromyrmex silvestrii bruchi</i> Forel, 1912	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex striatus</i> (Roger, 1863)	ar pr ur bo br	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Acromyrmex subterraneus</i> Forel, 1893	ar pr bo pe br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex subterraneus brunneus</i> Forel, 1912	br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex subterraneus molestans</i> Santschi, 1925	br	Gonçalves 1961; Bolton 1995
<i>Acromyrmex subterraneus ogloblini</i> Santschi, 1933	ar	Gonçalves 1961; Bolton 1995
<i>Acromyrmex subterraneus peruvianus</i> Borgmeier, 1940	pe	Gonçalves 1961; Bolton 1995
<i>Acromyrmex versicolor</i> (Pergande, 1894)	me	Gonçalves 1961; Fowler 1988; Bolton 1995
<i>Apterostigma abdita</i> Mann, 1922	ho	Lattke 1997
<i>Apterostigma acre</i> Lattke, 1997	br	Lattke 1997
<i>Apterostigma ancilonodum</i> Lattke, 1997	co	Lattke 1997
<i>Apterostigma andense</i> Lattke, 1997	vn pe	Lattke 1997
<i>Apterostigma angustum</i> Lattke, 1997	vn	Lattke 1997
<i>Apterostigma auriculatum</i> Wheeler W.M., 1925	br gi gf su tt vn co pn	Lattke 1997
<i>Apterostigma avium</i> Lattke, 1997	vn	Lattke 1997
<i>Apterostigma boliviannum</i> Weber, 1938	ec bo pe vn	Lattke 1997
<i>Apterostigma bruchi</i> Santschi, 1919	br	Lattke 1997
<i>Apterostigma callipygium</i> Lattke, 1997	co ec	Lattke 1997
<i>Apterostigma calverti</i> Wheeler W.M., 1911	cr	Lattke 1997
<i>Apterostigma carinatum</i> Lattke, 1997	co	Lattke 1997
<i>Apterostigma collare</i> Emery, 1896	co pn cr ni ho	Lattke 1997
<i>Apterostigma convexum</i> Lattke, 1997	vn	Lattke 1997
<i>Apterostigma chocoense</i> Lattke, 1997	cr pn co	Lattke 1997
<i>Apterostigma dentigerum</i> Wheeler W.M., 1925	pn cr co	Lattke 1997
<i>Apterostigma depressum</i> Lattke, 1997	ec pe	Lattke 1997
<i>Apterostigma discrepans</i> Forel, 1912	br	Lattke 1997
<i>Apterostigma dorothae</i> Weber, 1937	gi	Lattke 1997
<i>Apterostigma epinotale</i> Weber, 1937	gi br	Lattke 1997
<i>Apterostigma goniodes</i> Lattke, 1997	cr	Lattke 1997

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Apterostigma ierense</i> Weber, 1937	tt co ec pe br	Lattke 1997
<i>Apterostigma jubatum</i> Wheeler W.M., 1925	gi br	Lattke 1997
<i>Apterostigma madidiense</i> Weber, 1938	pe - ar	Lattke 1997
<i>Apterostigma manni</i> Weber, 1938	co - br	Lattke 1997
<i>Apterostigma mayri</i> Forel, 1893	tt	Lattke 1997
<i>Apterostigma megacephala</i> Lattke, 1999	pe co	Lattke 1997
<i>Apterostigma mexicanum</i> Lattke, 1997	me	Lattke 1997
<i>Apterostigma moelleri</i> Forel, 1892	br	Lattke 1997
<i>Apterostigma pallidum</i> Weber, 1937	tt	Lattke 1997
<i>Apterostigma pariense</i> Lattke, 1997	vn	Lattke 1997
<i>Apterostigma peruvianum</i> Wheeler W.M., 1925	pe br	Lattke 1997
<i>Apterostigma pilosum</i> Mayr, 1865	ar pr br cr me	Lattke 1997
<i>Apterostigma reburrum</i> Lattke, 1997	co	Lattke 1997
<i>Apterostigma robustum</i> Emery, 1896	bo br gi gf su cr	Lattke 1997
<i>Apterostigma scutellare</i> Forel, 1885	me	Lattke 1997
<i>Apterostigma serratum</i> Lattke, 1997	br	Lattke 1997
<i>Apterostigma spiculum</i> Lattke, 1997	pe	Lattke 1997
<i>Apterostigma steigeri</i> Santschi, 1911	br ur ar	Lattke 1997
<i>Apterostigma tachirense</i> Lattke, 1997	vn	Lattke 1997
<i>Apterostigma tholiforme</i> Lattke, 1997	ec	Lattke 1997
<i>Apterostigma tramitis</i> Weber, 1940	pn	Lattke 1997
<i>Apterostigma tropicoxa</i> Lattke, 1997	pe br	Lattke 1997
<i>Apterostigma trapeziforme</i> Lattke, 1997	es	Lattke 1997
<i>Apterostigma turgidum</i> Lattke, 1997	pn co	Lattke 1997
<i>Apterostigma urichii</i> Forel, 1893	bo br gi gf su tt	Lattke 1997
<i>Apterostigma wasmanni</i> Forel, 1892	br gi gf su tt	Lattke 1997
<i>Apterostigma zip</i> Weber, 1938	bo	Lattke 1997
<i>Atta bisphaerica</i> Forel, 1908	br	Borgmeier 1959
<i>Atta capiguara</i> Gonçalves, 1944	br	Borgmeier 1959
<i>Atta cephalotes</i> (Linnaeus, 1758)	bo pe ec br gi gf su tt vn co pn cr be ho gu me an co pn cr ni gu br am pr bo br gi gf su vn co es gu me us br br ar ar pr ur bo pe ec br gi gf su vn co pn cr br ar pr ur bo br	Borgmeier 1959; MacKay & MacKay 1986
<i>Atta columbica</i> Guérin-Méneville, 1845	br	Borgmeier 1959; MacKay & MacKay 1986
<i>Atta goiana</i> Gonçalves, 1942	am	Borgmeier 1959; MacKay & MacKay 1986
<i>Atta insularis</i> Guérin-Méneville, 1844	pr bo br gi gf su vn co	Borgmeier 1959
<i>Atta laevigata</i> (Smith F., 1858)	es gu me us	Borgmeier 1959
<i>Atta mexicana</i> (Smith F., 1858)	br	Borgmeier 1959
<i>Atta opaciceps</i> Borgmeier, 1939	br	Borgmeier 1959
<i>Atta robusta</i> Borgmeier, 1939	br	Borgmeier 1959
<i>Atta saltensis</i> Forel, 1913	ar	Borgmeier 1959
<i>Atta sexdens</i> (Linnaeus, 1758)	ar pr ur bo pe ec br gi gf su vn co pn cr br ar pr ur bo br	Borgmeier 1959; MacKay & MacKay 1986
<i>Atta silvae</i> Gonçalves, 1983	br	Borgmeier 1959
<i>Atta vollenweideri</i> Forel, 1893	br	Borgmeier 1959
<i>Cyphomyrmex auritus</i> Mayr, 1887	co	Kempf 1964b
<i>Cyphomyrmex bicarinatus</i> Snelling & Longino, 1992	br	Snelling & Longino 1992
<i>Cyphomyrmex bicornis</i> Forel, 1896	br	Kempf 1964b
<i>Cyphomyrmex bigibbosus</i> Emery, 1894	br gi gf su tt	Kempf 1964b
<i>Cyphomyrmex bruchi</i> Santschi, 1917	ar	Kempf 1966
<i>Cyphomyrmex castanei</i> MacKay & Baena, 1993	pn	Bolton 1995
<i>Cyphomyrmex cornutus</i> Kempf, 1968	co	Kempf 1966
<i>Cyphomyrmex costatus</i> Mann, 1922	co pn ho	Kempf 1964b
<i>Cyphomyrmex daguerrei</i> Santschi, 1933	ar	Kempf 1966
<i>Cyphomyrmex dixus</i> Snelling & Longino, 1992	cr	Snelling & Longino 1992
<i>Cyphomyrmex faunulus</i> Wheeler W.M., 1925	bo br gi gf su tt	Snelling & Longino 1992
<i>Cyphomyrmex flavidus</i> Pergande, 1896	me	Kempf 1966
<i>Cyphomyrmex foxi</i> André, 1892	am	Kempf 1966
<i>Cyphomyrmex hamulatus</i> Weber, 1938	bo br vn co pn	Kempf 1966
<i>Cyphomyrmex kirbyi</i> Mayr, 1887	ec vn co	Kempf 1966
<i>Cyphomyrmex laevigatus</i> Weber, 1938	bo pe br gi gf su	Kempf 1964b
<i>Cyphomyrmex lectus</i> Forel, 1911	ar br	Kempf 1964b

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Cyphomyrmex lilloanus</i> Kusnezov, 1949	ar	Kempf 1966
<i>Cyphomyrmex longiscapus</i> Weber, 1940	co pn cr	Kempf 1966
<i>Cyphomyrmex major</i> Forel, 1901	br gu	kempf 1966
<i>Cyphomyrmex minutus</i> Mayr, 1862	gi gf su tt vn co cr ho me us am an bh	Kempf 1964b
<i>Cyphomyrmex morschi</i> Emery, 1888	ar br	Kempf 1966
<i>Cyphomyrmex muelleri</i> Schultz & Solomon, 2002	ec co pn	Schultz & Solomon 2002
<i>Cyphomyrmex nemei</i> Kusnezov, 1957	ar	Kempf 1966
<i>Cyphomyrmex nesiotes</i> Snelling & Longino, 1992	ec	kempf 1966
<i>Cyphomyrmex occultus</i> Kempf, 1964	br	Kempf 1964b
<i>Cyphomyrmex olitor</i> Forel, 1893	ar br	Kempf 1964b
<i>Cyphomyrmex panicus</i> Wheeler W.M., 1925	br	Kempf 1966
<i>Cyphomyrmex peltatus</i> Kempf, 1966	br	Kempf 1964d
<i>Cyphomyrmex plaumannii</i> Kempf, 1962	br	Kempf 1966
<i>Cyphomyrmex podargus</i> Snelling & Longino, 1992	am	Snelling & Longino 1992
<i>Cyphomyrmex rimosus</i> (Spinola, 1851)	ar pr ur bo pe br gi gf su tt vn co pn cr ni be gu me am an	Kempf 1966, Snelling & Longino 1992
<i>Cyphomyrmex salvini</i> Forel, 1899	co pn cr	Kempf 1964b
<i>Cyphomyrmex strigatus</i> Mayr, 1887	br	Kempf 1966
<i>Cyphomyrmex transversus</i> Emery, 1894	ar pr br	Kempf 1964b
<i>Cyphomyrmex vallensis</i> Kusnezov, 1949	ar	Kempf 1966
<i>Cyphomyrmex vorticis</i> Weber, 1940	bo br vn	Kempf 1966
<i>Cyphomyrmex wheeleri</i> Forel, 1900	us	Kempf 1966
<i>Mycetagoicus cerradensis</i> Brandão & Mayhé-Nunes, 2001	br	Brandão & Mayhé-Nunes, 2001
<i>Mycetagoicus triangularis</i> Brandão & Mayhé-Nunes, 2001	br	Brandão & Mayhé-Nunes, 2001
<i>Mycetagoicus urbanus</i> Brandão & Mayhé-Nunes, 2001	br	Brandão & Mayhé-Nunes, 2001
<i>Mycetarotes acutus</i> Mayhé-Nunes, 1995	br	Kempf 1961; Mayhé-Nunes 1995
<i>Mycetarotes carinatus</i> Mayhé-Nunes, 1995	vn br	Kempf 1961; Mayhé-Nunes 1995
<i>Mycetarotes parallelus</i> (Emery, 1906)	br	Kempf 1961; Mayhé-Nunes 1995
<i>Mycetarotes senticosus</i> Kempf, 1960	br	Kempf 1961; Mayhé-Nunes 1995
<i>Mycetophylax bruchi</i> (Santschi, 1916)	ar	Bolton 1995
<i>Mycetophylax bruchi pauper</i> Santschi, 1923	ar	Bolton 1995
<i>Mycetophylax conformis</i> Mayr, 1884	br gi gf su tt am an	Bolton 1995
<i>Mycetophylax cristulatus</i> Santschi, 1922	ar	Bolton 1995
<i>Mycetophylax cristulatus emmae</i> Santschi, 1929	ar	Bolton 1995
<i>Mycetophylax emeryi</i> Forel, 1907	vn co	Bolton 1995
<i>Mycetophylax emeryi arenicola</i> Forel, 1912	ar	Bolton 1995
<i>Mycetophylax emeryi argentina</i> Santschi, 1916	ar	Bolton 1995
<i>Mycetophylax emeryi bolivari</i> Weber, 1948	vn	Bolton 1995
<i>Mycetophylax emeryi fortis</i> Forel, 1912	ar pr	Bolton 1995
<i>Mycetophylax emeryi gallardoi</i> Santschi, 1922	ar ur	Bolton 1995
<i>Mycetophylax emeryi hubrichi</i> Santschi, 1925	ar	Bolton 1995
<i>Mycetophylax emeryi weiseri</i> Santschi, 1929	ar	Bolton 1995
<i>Mycetophylax glaber</i> Weber, 1948	bo	Bolton 1995
<i>Mycetophylax simplex</i> Emery, 1888	pr br	Bolton 1995
<i>Mycetosoritis aspera</i> (Mayr, 1887)	ar br	Bolton 1995
<i>Mycetosoritis clorindae</i> Kusnezov, 1949	ar pr br	Bolton 1995
<i>Mycetosoritis explicata</i> Kempf, 1968	br	MacKay 1998
<i>Mycetosoritis vinsoni</i> Mackay, 1998	ar pr ur br	Bolton 1995
<i>Mycoceropurus goeldii</i> Forel, 1893	br	Kempf 1963
<i>Mycoceropurus curvispinosus</i> MacKay, 1998	ar pr bo pe br gi gf su co	MacKay 1998
<i>Mycoceropurus obsoletus</i> Emery, 1913	pn cr ni ho me am an	Kempf 1963
<i>Mycoceropurus smithii</i> Forel, 1893	pn	Kempf 1963
<i>Mycoceropurus tardus</i> Weber, 1940	bo	Bolton 1995
<i>Myrmicocrypta boliviiana</i> Weber, 1938	ar	Bolton 1995
<i>Myrmicocrypta bruchi</i> Santschi, 1936	gi gf su tt	Bolton 1995
<i>Myrmicocrypta buenzlii</i> Borgmeier, 1934	pe	Bolton 1995
<i>Myrmicocrypta collaris</i> Emery, 1913	pn cr ni be ho me	Bolton 1995
<i>Myrmicocrypta dilacerata</i> Forel, 1885		

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Myrmicocrypta dilacerata cornuta</i> Forel, 1899	me	Bolton 1995
<i>Myrmicocrypta ednaella</i> Mann, 1922	pn cr ho	Bolton 1995
<i>Myrmicocrypta elisabethae</i> Weber, 1937	gi	Bolton 1995
<i>Myrmicocrypta foreli</i> Mann, 1916	br	Bolton 1995
<i>Myrmicocrypta foreli surianensis</i> Wheeler W.M., 1925	bo	Bolton 1995
<i>Myrmicocrypta godmani</i> Forel, 1899	ni	Bolton 1995
<i>Myrmicocrypta guianensis</i> Weber, 1937	gi	Bolton 1995
<i>Myrmicocrypta infuscata</i> Weber, 1946	gi	Bolton 1995
<i>Myrmicocrypta longinoda</i> Weber, 1938	br	Bolton 1995
<i>Myrmicocrypta microphthalmia</i> Borgmeier, 1948	pe	Bolton 1995
<i>Myrmicocrypta occipitalis</i> Weber, 1938	bo	Bolton 1995
<i>Myrmicocrypta ogloblini</i> Santschi, 1936	ar	Bolton 1995
<i>Myrmicocrypta rufidiscapta</i> Emery, 1913	bo	Bolton 1995
<i>Myrmicocrypta spinosa</i> Weber, 1937	gi	Bolton 1995
<i>Myrmicocrypta squamosa</i> (Smith F., 1860)	ar pr br tt	Bolton 1995
<i>Myrmicocrypta squamosa uncinata</i> Mayr, 1887	pr br gi gf su	Bolton 1995
<i>Myrmicocrypta subnitida</i> Forel, 1899	pn	Bolton 1995
<i>Myrmicocrypta triangulata</i> Forel, 1912	gu	Bolton 1995
<i>Myrmicocrypta triangulata peruviana</i> Emery, 1913	bo pe	Bolton 1995
<i>Myrmicocrypta tuberculata</i> Weber, 1938	bo	Bolton 1995
<i>Myrmicocrypta unidentata</i> Weber, 1937	gi	Bolton 1995
<i>Myrmicocrypta urichi</i> Weber, 1937	tt an	Bolton 1995
<i>Myrmicocrypta weyrauchi</i> Borgmeier, 1948	pe	Bolton 1995
<i>Pseudoatta argentina</i> Gallardo, 1916	ar	Bolton 1995
<i>Sericomyrmex amabilis</i> Wheeler W.M., 1925	pn	Bolton 1995
<i>Sericomyrmex aztecus</i> Forel, 1885	me	Bolton 1995
<i>Sericomyrmex beniensis</i> Weber, 1938	bo	Bolton 1995
<i>Sericomyrmex bondari</i> Borgmeier, 1937	br	Bolton 1995
<i>Sericomyrmex burchelli</i> Forel, 1905	br	Bolton 1995
<i>Sericomyrmex diego</i> Forel, 1912	co	Bolton 1995
<i>Sericomyrmex harenkulli</i> Weber, 1937	gi	Bolton 1995
<i>Sericomyrmex harenkulli arawakensis</i> Weber, 1937	gi	Bolton 1995
<i>Sericomyrmex impexus</i> Wheeler W.M., 1925	gi	Bolton 1995
<i>Sericomyrmex luederwaldti</i> Santschi, 1925	br	Bolton 1995
<i>Sericomyrmex lutzi</i> Wheeler, 1916	gi gf su	Bolton 1995
<i>Sericomyrmex mayri</i> Forel, 1912	br	Bolton 1995
<i>Sericomyrmex moreirai</i> Santschi, 1925	br	Bolton 1995
<i>Sericomyrmex myersi</i> Weber, 1937	gi	Bolton 1995
<i>Sericomyrmex opacus</i> Mayr, 1865	br	Bolton 1995
<i>Sericomyrmex opacus muelleri</i> Forel, 1912	br	Bolton 1995
<i>Sericomyrmex parvulus</i> Forel, 1912	br	Bolton 1995
<i>Sericomyrmex saussurei</i> Emery, 1894	br	Bolton 1995
<i>Sericomyrmex scrobifer</i> Forel, 1911	br	Bolton 1995
<i>Sericomyrmex urichi</i> Forel, 1912	tt	Bolton 1995
<i>Sericomyrmex urichi maracas</i> Weber, 1937	tt	Bolton 1995
<i>Sericomyrmex zacapanus</i> Wheeler W.M., 1925	gu	Bolton 1995
<i>Trachymyrmex agudensis</i> Kempf, 1967	br	Bolton 1995
<i>Trachymyrmex bugnioni</i> Forel, 1912	gi gf su tt vn co pn cr ho	Bolton 1995
<i>Trachymyrmex carib</i> Weber, 1945	tt	Bolton 1995
<i>Trachymyrmex cornetzi</i> Forel, 1912	gi gf su tt vn co pn	Bolton 1995
<i>Trachymyrmex dichrous</i> Kempf, 1967	br	Bolton 1995
<i>Trachymyrmex diversus</i> Mann, 1916	br gi gf su vn	Bolton 1995
<i>Trachymyrmex echinus</i> Weber, 1938	bo	Bolton 1995
<i>Trachymyrmex farinosus</i> Emery, 1894	br	Bolton 1995
<i>Trachymyrmex fiebrigi</i> Santschi, 1916	pr	Bolton 1995
<i>Trachymyrmex gaigei</i> Forel, 1914	co	Bolton 1995
<i>Trachymyrmex guianensis</i> Weber, 1937	gi	Bolton 1995
<i>Trachymyrmex holmgreni</i> Wheeler W.M., 1925	bo	Bolton 1995
<i>Trachymyrmex intermedius</i> Forel, 1909	be gu me	Bolton 1995
<i>Trachymyrmex irmgardae</i> Forel, 1912	co	Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Trachymyrmex isthmicus</i> Santschi, 1931	pn	Bolton 1995
<i>Trachymyrmex jamaicensis</i> (André, 1893)	am bh	Bolton 1995
<i>Trachymyrmex jamaicensis frontalis</i> Santschi, 1925	am	Bolton 1995
<i>Trachymyrmex itheringi</i> (Emery, 1887)	br	Bolton 1995
<i>Trachymyrmex levis</i> Weber, 1938	bo	Bolton 1995
<i>Trachymyrmex mandibularis</i> Weber, 1938	bo	Bolton 1995
<i>Trachymyrmex oetkeri</i> Forel, 1908	br	Bolton 1995
<i>Trachymyrmex opulentus</i> (Mann, 1922)	ho	Bolton 1995
<i>Trachymyrmex papulatus</i> Santschi, 1922	ar	Bolton 1995
<i>Trachymyrmex phaleratus</i> Wheeler, 1925	gi	Bolton 1995
<i>Trachymyrmex pruinosis</i> Emery, 1905	ar ur	Bolton 1995
<i>Trachymyrmex relictus</i> Borgmeier, 1934	br gi gf su tt	Bolton 1995
<i>Trachymyrmex rufiae</i> Weber, 1937	tt	Bolton 1995
<i>Trachymyrmex saussurei</i> (Forel, 1884)	be me	Bolton 1995
<i>Trachymyrmex squamulifer</i> Emery, 1896	cr	Bolton 1995
<i>Trachymyrmex tucumanus</i> Forel, 1914	ar	Bolton 1995
<i>Trachymyrmex uruchi</i> Forel, 1893	br gi gf su tt vn co pn	Bolton 1995
Tribe Basicerotini		
<i>Basiceros convexus</i> Mayr, 1887	br	Brown & Kempf 1960
<i>Basiceros manni</i> Brown & Kempf, 1960	cr ni ho	Brown & Kempf 1960
<i>Basiceros redux</i> (Donisthorpe, 1939)	gi	Brown & Kempf 1960
<i>Basiceros singularis</i> (Smith F., 1858)	br gi gf su tt	Brown & Kempf 1960
<i>Basiceros scambognathus</i> Brown, 1949	br	Brown & Kempf 1960
<i>Eurhopalothrix alopeciosa</i> Brown & Kempf, 1960	ec tt	Brown & Kempf 1960
<i>Eurhopalothrix aphrodisia</i> Snelling, 1968	es	Bolton 1995
<i>Eurhopalothrix balaui</i> (Mayr, 1870)	br gi gf su pn ho	Brown & Kempf 1960
<i>Eurhopalothrix bruchi</i> Santschi, 1922	ar	Brown & Kempf 1960
<i>Eurhopalothrix clypeata</i> Brown & Kempf, 1960	gi	Brown & Kempf 1960
<i>Eurhopalothrix gravis</i> (Mann, 1922)	br vn cr ni ho me an	Brown & Kempf 1960
<i>Eurhopalothrix lenkoi</i> Kempf, 1967	br	Brown & Kempf 1960
<i>Eurhopalothrix pilulifera</i> Brown & Kempf, 1960	br me	Brown & Kempf 1960
<i>Eurhopalothrix speciosa</i> Brown & Kempf, 1960	br	Brown & Kempf 1960
<i>Eurhopalothrix spectabilis</i> Kempf, 1962	bo ec br gi gf su tt co	Brown & Kempf 1960
<i>Ocotostruma balzani</i> (Emery, 1894)	pn cr ni gu me an	Brown & Kempf 1960
<i>Ocotostruma batesi</i> Emery, 1894	br gi gf su	Brown & Kempf 1960
<i>Ocotostruma impressa</i> Palacio, 1997		Palacio 1997
<i>Ocotostruma inca</i> Brown y Kempf, 1960	bo pe co	Brown & Kempf 1960
<i>Ocotostruma itheringi</i> (Emery, 1888)	br tt vn co pn cr gu am	Brown & Kempf 1960
<i>Ocotostruma petiolata</i> (Mayr, 1887)	br	Brown & Kempf 1960
<i>Ocotostruma rugifera</i> (Mayr, 1887)	ar br	Brown & Kempf 1960
<i>Ocotostruma rugiferaoides</i> Brown & Kempf, 1960	me	Brown & Kempf 1960
<i>Ocotostruma stenognatha</i> Brown & Kempf, 1960	br	Brown & Kempf 1960
<i>Ocotostruma stenoscapa</i> Palacio, 1997		Palacio 1997
<i>Ocotostruma wheeleri</i> (Mann, 1922)	gu	Brown & Kempf 1960
<i>Protalaridris armata</i> Brown, 1980	ec co	Brown 1980
<i>Rhopalothrix acutipilis</i> Kempf, 1962	br	Bolton 1995
<i>Rhopalothrix ciliata</i> Mayr, 1870	co	Brown & Kempf 1960
<i>Rhopalothrix isthmica</i> Weber, 1941	pn	Brown & Kempf 1960
<i>Rhopalothrix kusnezovi</i> Brown & Kempf, 1960	ar	Brown & Kempf 1960
<i>Rhopalothrix plaumanni</i> Brown & Kempf, 1960	br	Brown & Kempf 1960
<i>Rhopalothrix stannardi</i> Brown & Kempf, 1960	me	Brown & Kempf 1960
<i>Rhopalothrix weberi</i> Brown & Kempf, 1960	am	Brown & Kempf 1960
<i>Talaridris mandibularis</i> Weber, 1941	tt	Brown & Kempf 1960
Tribe Blepharidattini		
<i>Blepharidatta brasiliensis</i> André, 1893	ar br	Bolton 1995
<i>Blepharidatta conops</i> Kempf, 1967	br	Bolton 1995
<i>Wasmaniella affinis</i> Santschi, 1929	br	Longino & Fernández 2004
<i>Wasmaniella auropunctata</i> (Roger, 1863)	ur pe ec br	Longino & Fernández 2004
<i>Wasmaniella itheringi</i> Forel, 1908	br	Longino & Fernández 2004

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Wasemannia lutzi</i> Forel, 1908	br	Longino & Fernández 2004
<i>Wasemannia rochai</i> Forel, 1912	br gi gf su tt pn	Longino & Fernández 2004
<i>Wasemannia scrobifera</i> Kempf, 1961	gi	Longino & Fernández 2004
<i>Wasemannia sigmoidea</i> (Mayr, 1884)	br gi gf su an	Longino & Fernández 2004
<i>Wasemannia sulcaticeps</i> Emery, 1894	ar	Longino & Fernández 2004
<i>Wasemannia villosa</i> Emery, 1894	br	Longino & Fernández 2004
<i>Wasemannia williamsoni</i> Kusnezov, 1952	ar	Longino & Fernández 2004
Tribe Cephalotini		
<i>Cephalotes adolphi</i> Emery, 1905	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes alfaroi</i> (Emery, 1890)	pn cr	De Andrade & Baroni Urbani 1999
<i>Cephalotes alveolatus*</i> (Vierbergen & Scheven, 1995)	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes angustus</i> (Mayr, 1862)	ar br	De Andrade & Baroni Urbani 1999
<i>Cephalotes argentiventris</i> Andrade, 1999	am	De Andrade & Baroni Urbani 1999
<i>Cephalotes atratus</i> (Linnaeus, 1758)	ar pr ur bo pe ec br gi gf su tt vn co pn cr be	De Andrade & Baroni Urbani 1999
<i>Cephalotes auriger</i> Andrade, 1999	me	De Andrade & Baroni Urbani 1999
<i>Cephalotes auricomus</i> (Wheeler, 1936)	am	De Andrade & Baroni Urbani 1999
<i>Cephalotes basalis</i> (Smith, 1876)	ec co pn cr ni es ho gu me	De Andrade & Baroni Urbani 1999
<i>Cephalotes betoii</i> Andrade, 1999	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes biguttatus</i> (Emery, 1890)	cr be ho gu me	De Andrade & Baroni Urbani 1999
<i>Cephalotes bimaculatus</i> (Smith F., 1860)	me	De Andrade & Baroni Urbani 1999
<i>Cephalotes bivestitus</i> (Santschi, 1922)	ar	De Andrade & Baroni Urbani 1999
<i>Cephalotes bloosi*</i> Baroni Urbani & Andrade, 1999	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes bohlsi</i> (Emery, 1896)	ar pr br	De Andrade & Baroni Urbani 1999
<i>Cephalotes borgmeieri</i> (Kempf, 1890)	ar pr br	De Andrade & Baroni Urbani 1999
<i>Cephalotes brevispinus*</i> Andrade, 1999	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes bruchi</i> (Forel, 1912)	ar br	De Andrade & Baroni Urbani 1999
<i>Cephalotes caribicus*</i> Andrade, 1999	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes chacomul</i> Snelling, 1999	me	De Andrade & Baroni Urbani 1999
<i>Cephalotes christopherseni</i> (Forel, 1912)	vn co pn	De Andrade & Baroni Urbani 1999
<i>Cephalotes clypeatus</i> (Fabricius, 1804)	ar pr bo pe br gi gf su tt vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes coffeeae</i> (Kempf, 1953)	co	De Andrade & Baroni Urbani 1999
<i>Cephalotes columbica</i> (Forel, 1912)	vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes complanatus</i> (Guérin-Méneville, 1844)	bo pe br gf su co	De Andrade & Baroni Urbani 1999
<i>Cephalotes conspersus</i> (Smith F., 1867)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes cordatus</i> (Smith F., 1853)	bo pe br gf co	De Andrade & Baroni Urbani 1999
<i>Cephalotes cordiae</i> (Stitz, 1913)	bo pe br	De Andrade & Baroni Urbani 1999
<i>Cephalotes cordiventris</i> (Santschi, 1931)	pn cr	De Andrade & Baroni Urbani 1999
<i>Cephalotes crenaticeps</i> (Mayr, 1866)	vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes cristatus</i> (Emery, 1890)	co cr be ho gu me	De Andrade & Baroni Urbani 1999
<i>Cephalotes curvistriatus</i> (Forel, 1899)	cr	De Andrade & Baroni Urbani 1999
<i>Cephalotes decolor</i> Andrade, 1999	vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes decoloratus</i> Andrade, 1999	an	De Andrade & Baroni Urbani 1999
<i>Cephalotes dentidorsum</i> Andrade, 1999	pe	De Andrade & Baroni Urbani 1999
<i>Cephalotes depressus</i> (Klug, 1824)	ar pr bo ec br gi vn	De Andrade & Baroni Urbani 1999
<i>Cephalotes dieteri*</i> Baroni Urbani, 1999	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes duckei</i> (Forel, 1906)	br co	De Andrade & Baroni Urbani 1999
<i>Cephalotes ecuadorialis</i> Andrade, 1999	ec	De Andrade & Baroni Urbani 1999
<i>Cephalotes eduarduli</i> (Forel, 1920)	ar pr bo br	De Andrade & Baroni Urbani 1999
<i>Cephalotes emeryi</i> (Forel, 1912)	an	De Andrade & Baroni Urbani 1999
<i>Cephalotes femoralis</i> (Smith F., 1853)	vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes fiebrigi</i> (Forel, 1906)	ar pr br	De Andrade & Baroni Urbani 1999
<i>Cephalotes flavigaster</i> Andrade, 1999	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes foliaceus</i> (Emery, 1905)	bo pe co pn	De Andrade & Baroni Urbani 1999
<i>Cephalotes fossithorax</i> (Santschi, 1921)	ar	De Andrade & Baroni Urbani 1999
<i>Cephalotes frigidus</i> (Kempf, 1958)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes goeldii</i> (Forel, 1912)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes goniodontus</i> Andrade, 1999	me	De Andrade & Baroni Urbani 1999
<i>Cephalotes grandinosus</i> (Smith F., 1860)	gi tt vn co pn cr	De Andrade & Baroni Urbani 1999
<i>Cephalotes guayaki</i> Andrade, 1999	pr br	De Andrade & Baroni Urbani 1999

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Cephalotes hamulus</i> (Roger, 1863)	an	De Andrade & Baroni Urbani 1999
<i>Cephalotes hirsutus</i> Andrade, 1999	me	
<i>Cephalotes hispaniolicus*</i> Andrade, 1999	ad	
<i>Cephalotes inaequalis</i> (Mann, 1916)	br co	De Andrade & Baroni Urbani 1999
<i>Cephalotes inca</i> (Santschi, 1911)	pe	De Andrade & Baroni Urbani 1999
<i>Cephalotes incertus</i> (Emery, 1905)	ar pr br	De Andrade & Baroni Urbani 1999
<i>Cephalotes insularis</i> (Wheeler W.M., 1934)	me	De Andrade & Baroni Urbani 1999
<i>Cephalotes integerrimus*</i> (Vierbergen & Scheven, 1995)	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes jamaicensis</i> (Forel, 1922)	am	De Andrade & Baroni Urbani 1999
<i>Cephalotes jansei*</i> (Vierbergen & Scheven, 1995)	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes jheringi</i> (Emery, 1894)	ar pr br	De Andrade & Baroni Urbani 1999
<i>Cephalotes klugi</i> (Emery, 1894)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes kukulcan</i> Snelling, 1999	cr be me	De Andrade & Baroni Urbani 1999
<i>Cephalotes laminitus</i> (Smith F., 1860)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes lanuginosus</i> (Santschi, 1919)	ar	De Andrade & Baroni Urbani 1999
<i>Cephalotes lenca</i> Andrade, 1999	ho	De Andrade & Baroni Urbani 1999
<i>Cephalotes liepini</i> Andrade, 1999	br	
<i>Cephalotes liogaster</i> (Santschi, 1916)	ar	De Andrade & Baroni Urbani 1999
<i>Cephalotes maculatus</i> (Smith F., 1876)	ar pr bo pe ec br gi tt co pn cr ni ho gu me	De Andrade & Baroni Urbani 1999
<i>Cephalotes manni</i> (Kempf, 1951)	pe br gi co	De Andrade & Baroni Urbani 1999
<i>Cephalotes marginatus</i> (Fabricius, 1804)	br gi gf su co	De Andrade & Baroni Urbani 1999
<i>Cephalotes maya*</i> Andrade, 1999	me	
<i>Cephalotes membranaceus</i> (Klug, 1824)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes minutus</i> (Fabricius, 1804)	ar pr bo pe br gi tt vn co pn cr ni es be ho gu me an	De Andrade & Baroni Urbani 1999
<i>Cephalotes mompox</i> Andrade, 1999	vn co	
<i>Cephalotes multispinosus</i> (Norton, 1868)	pn cr me	De Andrade & Baroni Urbani 1999
<i>Cephalotes nilpiei</i> Andrade, 1999	br	
<i>Cephalotes notatus</i> (Mayr, 1866)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes obscurus*</i> (Vierbergen & Scheven, 1995)	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes olmecus*</i> Andrade, 1999	me	
<i>Cephalotes oculatus</i> (Spinola, 1853)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes opacus</i> Santschi, 1920	bo pe ec br gi gf su vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes pallens</i> (Klug, 1824)	br gi gf su tt vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes pallidicephalus</i> (Smith F., 1876)	pn cr ni gu me an	
<i>Cephalotes pallidoides</i> Andrade, 1999	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes pallidus</i> Andrade, 1999	bo br gi tt vn	
<i>Cephalotes palta</i> Andrade, 1999	bo pe br gi su	
<i>Cephalotes palustris</i> Andrade, 1999	co	
<i>Cephalotes patei</i> (Kempf, 1951)	br gi	
<i>Cephalotes patellaris</i> (Mayr, 1866)	co	De Andrade & Baroni Urbani 1999
<i>Cephalotes pavonii</i> (Latreille, 1809)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes pellans</i> Andrade, 1999	pe br gi gf su co	De Andrade & Baroni Urbani 1999
<i>Cephalotes persimilis</i> Andrade, 1999	pr bo br	
<i>Cephalotes persimplex</i> Andrade, 1999	ar pr br	
<i>Cephalotes peruviensis</i> Andrade, 1999	bo br	
<i>Cephalotes pileini</i> Andrade, 1999	pe	
<i>Cephalotes pilosus</i> (Emery, 1896)	ar	
<i>Cephalotes pinelii</i> (Guérin y Méneville, 1844)	ar pr br	De Andrade & Baroni Urbani 1999
<i>Cephalotes placidus</i> (Smith F., 1860)	pr br	De Andrade & Baroni Urbani 1999
<i>Cephalotes poinari*</i> Baroni Urbani, 1999	pe br gi co	De Andrade & Baroni Urbani 1999
<i>Cephalotes porrasi</i> (Wheeler W.M., 1942)	me	
<i>Cephalotes prodigiosus</i> (Santschi, 1921)	ec co pn cr ni be me	De Andrade & Baroni Urbani 1999
<i>Cephalotes pusillus</i> (Klug, 1824)	ar	De Andrade & Baroni Urbani 1999
<i>Cephalotes quadratus</i> (Mayr, 1868)	ar pr bo pe ec br gi gf su tt vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes ramophilus</i> (Forel, 1904)	ar	De Andrade & Baroni Urbani 1999
<i>Cephalotes resinae*</i> Andrade, 1999	bo pe ec br co	Kempf 1967, De Andrade & Baroni Urbani 1999
<i>Cephalotes rohweri</i> (Wheeler W.M., 1916)	ad me us	De Andrade & Baroni Urbani 1999

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Cephalotes scutulatus</i> (Smith F., 1867)	ec vn co pn cr es be ho gu me	De Andrade & Baroni Urbani 1999
<i>Cephalotes serraticeps</i> (Smith F., 1858)	pe br co	De Andrade & Baroni Urbani 1999
<i>Cephalotes serratus*</i> (Vierbergen & Scheven, 1995)	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes setulifer</i> (Emery, 1894)	co pn cr	De Andrade & Baroni Urbani 1999
<i>Cephalotes simillimus</i> (Kempf, 1951)	bo pe br gi gf su co	De Andrade & Baroni Urbani 1999
<i>Cephalotes sobrius</i> (Kempf, 1958)	gu	De Andrade & Baroni Urbani 1999
<i>Cephalotes solidus</i> (Kempf, 1974)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes spinosus</i> (Mayr, 1862)	bo pe ec br gi gf su tt vn co	De Andrade & Baroni Urbani 1999
<i>Cephalotes squamosus*</i> (Vierbergen & Scheven, 1995)	ad	De Andrade & Baroni Urbani 1999
<i>Cephalotes sucinus*</i> Andrade, 1999	ad	
<i>Cephalotes supercilii</i> Andrade, 1999	ar	
<i>Cephalotes taino</i> Andrade, 1999	am	
<i>Cephalotes targionii</i> (Emery, 1894)	pr bo br gi tt vn	De Andrade & Baroni Urbani 1999
<i>Cephalotes texanus</i> (Santschi, 1915)	us	De Andrade & Baroni Urbani 1999
<i>Cephalotes toltecus</i> Andrade, 1999	me	
<i>Cephalotes trichophorus</i> Andrade, 1999	pe br	
<i>Cephalotes umbraculatus</i> (Fabricius, 1804)	bo br gi gf su tt vn co	De Andrade & Baroni Urbani 1999
	pn cr es ho gu me	
<i>Cephalotes unimaculatus</i> (Smith F., 1854)	an	De Andrade & Baroni Urbani 1999
<i>Cephalotes ustus</i> (Kempf, 1973)	br	De Andrade & Baroni Urbani 1999
<i>Cephalotes varians</i> (Smith F., 1876)	us am bh	De Andrade & Baroni Urbani 1999
<i>Cephalotes ventriosus*</i> Andrade, 1999	me	
<i>Cephalotes vinosus</i> (Wheeler, 1936)	am	De Andrade & Baroni Urbani 1999
<i>Cephalotes wheeleri</i> (Forel, 1901)	me	De Andrade & Baroni Urbani 1999
<i>Procryptocerus adlerzi</i> (Mayr, 1887)	br	Kempf 1951
<i>Procryptocerus attenuatus</i> (Smith F., 1876)	br gi gf su pn	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus balzani</i> Emery, 1894	co pn cr	Kempf 1951
<i>Procryptocerus batesi</i> Forel, 1899	co	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus belti</i> Forel, 1899	ec co pn cr ni be ho gu me	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus carbonarius</i> (Mayr, 1870)	co pn	Kempf 1951
<i>Procryptocerus clathratus</i> Emery, 1896	br	Kempf 1951
<i>Procryptocerus convergens</i> (Mayr, 1887)	br	Kempf 1951
<i>Procryptocerus convexus</i> Forel, 1904	br	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus coriarius</i> (Mayr, 1870)	co pn cr	Kempf 1951
<i>Procryptocerus curvistriatus</i> Kempf, 1949	br	Kempf 1951
<i>Procryptocerus eladio</i> Longino & Snelling, 2002	cr	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus elegans</i> Santschi, 1921	br	Kempf 1951
<i>Procryptocerus ferreri</i> Forel, 1912	co	Kempf 1951
<i>Procryptocerus gibbosus</i> Kempf, 1949	br	Kempf 1951
<i>Procryptocerus goeldii</i> Forel, 1899	pr br	Kempf 1951
<i>Procryptocerus gracilis</i> (Smith F., 1858)	br	Kempf 1951
<i>Procryptocerus hirsutus</i> Emery, 1896	br gi tt	Kempf 1951
<i>Procryptocerus hylaeus</i> Kempf, 1951	pr bo pe br tt vn co pn	Kempf 1951
<i>Procryptocerus impressus</i> Forel, 1899	pn cr	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus kempfi</i> Longino & Snelling, 2002	pe co pn cr	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus lenkoi</i> Kempf, 1969	br	Kempf 1951
<i>Procryptocerus lepidus</i> Forel, 1908	br	Kempf 1951
<i>Procryptocerus marginatus</i> Borgmeier, 1948	br	Kempf 1951
<i>Procryptocerus mayri</i> Forel, 1899	pe br vn co pn cr	Kempf 1951
<i>Procryptocerus montanus</i> Kempf, 1957	br	Kempf 1951
<i>Procryptocerus nalinii</i> Longino & Snelling, 2002	pe cr	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus paleatus</i> Emery, 1896	pn cr me	Kempf 1951
<i>Procryptocerus pictipes</i> Emery, 1896	bo pe ec br gi tt vn co pn cr	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus regularis</i> Emery, 1888	ar br tt	Kempf 1951
<i>Procryptocerus rufus</i> (Mayr, 1870)	co	Kempf 1951
<i>Procryptocerus sampaioi</i> Forel, 1912	br	Kempf 1951
<i>Procryptocerus scabriusculus</i> Forel, 1899	vn co pn cr ni be gu me	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus schmitti</i> Forel, 1901	br vn	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus seabrai</i> Kempf, 1964	br	Kempf 1951
<i>Procryptocerus spiniperdus</i> Forel, 1899	pe ec br tt	Kempf 1951

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Procryptocerus striatus</i> (Smith, 1870)	br	Kempf 1951
<i>Procryptocerus subpilosus</i> (Smith F., 1860)	pe ec br gi tt	Kempf 1951
<i>Procryptocerus sulcatus</i> Emery, 1894	br	Kempf 1951
<i>Procryptocerus tortuguero</i> Longino & Snelling, 2002	cr	Kempf 1951; Longino & Snelling 2002
<i>Procryptocerus victoris</i> Kempf, 1960	br	Kempf 1951
<i>Procryptocerus virgatus</i> Kempf, 1964	ec	Kempf 1951
Tribe Crematogastrini		
<i>Crematogaster abstinentis</i> Forel, 1899	ar bo br gf vn co pn	Longino 2003
<i>Crematogaster acuta</i> (Fabricius, 1804)	bo pe br gi vn co pn cr ni ho gu me	Longino 2003
<i>Crematogaster amapaensis</i> Kempf, 1960	br	Bolton 1995
<i>Crematogaster ampla</i> Forel, 1912	co pn	Longino 2003
<i>Crematogaster arata</i> Emery, 1906	ar	Bolton 1995
<i>Crematogaster arcuata</i> Forel, 1899	pe br vn co pn cr	Longino 2003
<i>Crematogaster ashmeadi</i> Mayr, 1886	us bh	Bolton 1995
<i>Crematogaster atra</i> Mayr, 1870	br me	Longino 2003
<i>Crematogaster bingi</i> Forel, 1908	br	Bolton 1995
<i>Crematogaster boliviensis</i> (Wheeler W.M., 1922)	bo	Bolton 1995
<i>Crematogaster brasiliensis</i> Mayr, 1878	bo pe ec br gi co pn cr ni	Longino 2003
<i>Crematogaster brevidentata</i> Forel, 1912	br	Longino 2003
<i>Crematogaster brevispinosa crucis carminis</i> Forel, 1912	ar	Bolton 1995
<i>Crematogaster brevispinosa minutior themis</i> Santschi, 1925	br	Bolton 1995
<i>Crematogaster brevispinosa moelleri tucumanensis</i> Forel, 1914	ar	Longino 2003
<i>Crematogaster brevispinosa sericea semisericea</i> Santschi, 1923	ar pr	Bolton 1995
<i>Crematogaster brevispinosa tumulifera convicta</i> Forel, 1912	ar	Longino 2003
<i>Crematogaster brevispinosa tumulifera fautrix</i> Forel, 1908	br	Longino 2003
<i>Crematogaster brevispinosa tumulifera scurra</i> Forel, 1914	co	Bolton 1995
<i>Crematogaster brevispinosa tumulifera tumulicula</i> Forel, 1909		pr Longino 2003
<i>Crematogaster bruchi</i> Forel, 1912	ar	Bolton 1995
<i>Crematogaster bryophila</i> Longino, 2003	ec vn co pn cr	Bolton 1995
<i>Crematogaster californica</i> Wheeler W.M., 1919	us	Bolton 1995
<i>Crematogaster carinata</i> Mayr, 1862	ec br su vn co pn cr gu an	Longino 2003
<i>Crematogaster chodati</i> Forel, 1921	pr	Longino 2003
<i>Crematogaster clydia</i> Forel, 1912	br	Bolton 1995
<i>Crematogaster corticicola</i> Mayr, 1887	ar br	Bolton 1995
<i>Crematogaster corvina</i> Mayr, 1870	me	Bolton 1995
<i>Crematogaster crinosa</i> Mayr, 1862	ar ch pr ur bo pe br gi gf su vn co pn cr ni be ho gu me us am an	Longino 2003
<i>Crematogaster cristata</i> Santschi, 1929	br	Bolton 1995
<i>Crematogaster crucis</i> Forel, 1912	co cr	Bolton 1995
<i>Crematogaster cubensis</i> Mann, 1920	am	Longino 2003
<i>Crematogaster curvispinosa</i> Mayr, 1862	ar pr ur bo pe ec br gi gf su vn co pn cr ni gu me an	Longino 2003
<i>Crematogaster delitescens</i> Wheeler W.M., 1921	gi gf su an	Bolton 1995
<i>Crematogaster depilis</i> Wheeler W.M., 1919	us	Bolton 1995
<i>Crematogaster distans</i> Mayr, 1870	ar pe ec br vn co pn cr ni gu me us	Longino 2003
<i>Crematogaster dorsidens</i> Santschi, 1925	br	Bolton 1995
<i>Crematogaster egregior</i> Forel, 1912	br	Bolton 1995
<i>Crematogaster erecta</i> Mayr, 1866	pe ec br gi gf su co pn cr	Longino 2003
<i>Crematogaster euterpe</i> Santschi, 1922	ar	Bolton 1995
<i>Crematogaster evallans</i> Forel, 1907	br co cr	Longino 2003
<i>Crematogaster flavomicrops</i> Longino, 2003	ec br vn co pn cr	Bolton 1995
<i>Crematogaster flavosensitiva</i> Longino, 2003	vn co pn	Bolton 1995
<i>Crematogaster foliolyptica</i> Longino, 2003	cr	Bolton 1995
<i>Crematogaster formosa</i> Mayr, 1870	me	Bolton 1995
<i>Crematogaster formosa aterrima</i> Wheeler W.M., 1909	me	Bolton 1995
<i>Crematogaster goeldii</i> Forel, 1903	ar br	Longino 2003
<i>Crematogaster heathi</i> Mann, 1916	br	Longino 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Crematogaster hespera</i> Buren, 1968	us	Bolton 1995
<i>Crematogaster huberi</i> (Forel, 1907)	gi	Bolton 1995
<i>Crematogaster indefensa</i> Kempf, 1968	bo	Bolton 1995
<i>Crematogaster jardinero</i> Longino, 2003	cr	Bolton 1995
<i>Crematogaster iheringi</i> Forel, 1908	ar br	Bolton 1995
<i>Crematogaster jheringi bosqi</i> Santschi, 1934	ar	Bolton 1995
<i>Crematogaster jheringi nemesis</i> Santschi, 1934	ar	Bolton 1995
<i>Crematogaster laevis</i> Mayr, 1877	br	Bolton 1995
<i>Crematogaster laevissis bruesi</i> Wheeler, 1911	an	Bolton 1995
<i>Crematogaster levior</i> Longino, 2003	bo pe ec br gi gf su vn co	Longino 2003
<i>Crematogaster limata</i> Smith F., 1858	bo pe ec br gi tt vn co pn cr ni gu me	Longino 2003
<i>Crematogaster limata parabiotica levior</i> Forel, 1911	pe br	Bolton 1995
<i>Crematogaster longispina</i> Emery, 1890	pe ec gi gf su co cr ni	Longino 2003
<i>Crematogaster longispina boliviiana</i> Wheeler W.M., 1925	bo	Bolton 1995
<i>Crematogaster longispina naumannae</i> Forel, 1921	ec	Bolton 1995
<i>Crematogaster lucayana</i> Wheeler W.M., 1905	bh	Bolton 1995
<i>Crematogaster lucayana etiolata</i> Wheeler W.M., 1905	bh	Bolton 1995
<i>Crematogaster lutzi</i> Forel, 1905	br	Bolton 1995
<i>Crematogaster lutzi florida</i> Forel, 1908	br	Bolton 1995
<i>Crematogaster magnifica</i> Santschi, 1925	br	Bolton 1995
<i>Crematogaster malevolens</i> Santschi, 1919	ar pr	Longino 2003
<i>Crematogaster mancocapaci</i> Santschi, 1911	ec	Longino 2003
<i>Crematogaster manii</i> Buren, 1968	am	Bolton 1995
<i>Crematogaster marioni</i> Buren, 1968	us	Bolton 1995
<i>Crematogaster minutissima</i> Mayr 1870	cr me us	Longino 2003
<i>Crematogaster moelleri</i> Forel, 1912	ar br cr	Longino 2003
<i>Crematogaster montana</i> Borgmeier, 1939	br	Longino 2003
<i>Crematogaster monteverdensis</i> Longino, 2003	cr	Bolton 1995
<i>Crematogaster montezumia</i> Smith F., 1858	ar pe ec br co pn cr me	Longino 2003
<i>Crematogaster mormonum</i> Wheeler W.M., 1919	us	Bolton 1995
<i>Crematogaster nigropilosa</i> Mayr, 1870	bo pe ec br vn co pn cr ni	Longino 2003
<i>Crematogaster obscurata</i> Emery, 1895	vn co pn cr be gu me us	Longino 2003
<i>Crematogaster opaca</i> Mayr, 1870	me	Bolton 1995
<i>Crematogaster opaca cedrosensis</i> Wheeler W.M., 1934	me	Bolton 1995
<i>Crematogaster opaca dentinodis</i> Forel, 1901	me	Bolton 1995
<i>Crematogaster oxygynoides</i> Santschi, 1934	ar	Bolton 1995
<i>Crematogaster parallela</i> Santschi, 1925	br	Bolton 1995
<i>Crematogaster patei</i> Buren, 1968	me	Bolton 1995
<i>Crematogaster peristerica</i> Menozzi, 1925	br	Longino 2003
<i>Crematogaster peruviana</i> Wheeler W.M., 1922	pe	Bolton 1995
<i>Crematogaster polymnia</i> Santschi, 1922	ar	Bolton 1995
<i>Crematogaster polymnia oeagria</i> Santschi, 1933	ar	Bolton 1995
<i>Crematogaster quadriformis</i> Roger, 1863	ar pr ur pe br gi gf su co	Bolton 1995
<i>Crematogaster quadriformis gracilior</i> Forel, 1901	ar pr br	Bolton 1995
<i>Crematogaster quadriformis roveretoi</i> Forel, 1913	ar pr br	Bolton 1995
<i>Crematogaster quadriformis vezeyii</i> Forel, 1907	pr	Bolton 1995
<i>Crematogaster quadrispinosa</i> Roger, 1863	me	Bolton 1995
<i>Crematogaster raptor</i> Longino, 2003	cr	Bolton 1995
<i>Crematogaster rifelna</i> Buren, 1968	me us	Bolton 1995
<i>Crematogaster rochai</i> Forel, 1903	ar bo pe br gi vn co pn cr ni ho me	Longino 2003
<i>Crematogaster rossi</i> Buren, 1968	me	Bolton 1995
<i>Crematogaster rufidula</i> Emery, 1894	ar br	Bolton 1995
<i>Crematogaster russata</i> Wheeler W.M., 1924	bo	Longino 2003
<i>Crematogaster russoi</i> Menozzi y Russo, 1930	am	Bolton 1995
<i>Crematogaster sanguinea</i> Roger, 1863	am	Bolton 1995
<i>Crematogaster sanguinea atavista</i> Mann, 1920	am	Bolton 1995
<i>Crematogaster sanguinea sotolongoi</i> Mann, 1920	am	Bolton 1995
<i>Crematogaster sanguinea torrei</i> Wheeler W.M., 1913	am	Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Crematogaster saussurei</i> Forel, 1899	me	Bolton 1995
<i>Crematogaster scapamaris</i> Santschi, 1922	ar	Bolton 1995
<i>Crematogaster sclerata</i> Santschi, 1917	ar	Bolton 1995
<i>Crematogaster sclerata taperensis</i> Borgmeier, 1937	br	Bolton 1995
<i>Crematogaster sericea</i> Forel, 1912	ar pr ur br	Longino 2003
<i>Crematogaster snellingi</i> Longino, 2003	cr	Bolton 1995
<i>Crematogaster sotobosque</i> Longino, 2003	pe ec br vn co pn cr	Longino 2003
<i>Crematogaster steinheili argentina</i> Forel, 1915	ar	Bolton 1995
<i>Crematogaster steinheili</i> Forel, 1881	am an bh	Bolton 1995
<i>Crematogaster stigmatica</i> Forel, 1911	br	Longino 2003
<i>Crematogaster stollii</i> Forel, 1885	bo pe ec br gi vn co pn cr ni gu	Longino 2003
<i>Crematogaster subtensa</i> Santschi, 1925	br	Longino 2003
<i>Crematogaster sumichrasti</i> Mayr, 1870	cr ni be gu me	Longino 2003
<i>Crematogaster tenuicula</i> Forel, 1904	bo pe br gi vn co pn cr	Longino 2003
<i>Crematogaster thalia</i> Forel, 1911	pr	Longino 2003
<i>Crematogaster torosa</i> Mayr, 1870	co pn cr ni gu me us	Longino 2003
<i>Crematogaster uncinata</i> Santschi, 1925	br	Longino 2003
<i>Crematogaster uruguayensis</i> Santschi, 1912	ar ur	Longino 2003
<i>Crematogaster vicina</i> André, 1893	am	Longino 2003
<i>Crematogaster victima</i> Smith F., 1858	ar pr ur br vn	Bolton 1995
<i>Crematogaster victimae allegrensis</i> Forel, 1901	br	Bolton 1995
<i>Crematogaster victimae cisplatinalis</i> Mayr, 1887	ur br	Bolton 1995
<i>Crematogaster victimae nitidiceps</i> Emery, 1895	ur br	Bolton 1995
<i>Crematogaster victimae nitidiceps nana</i> Santschi, 1934	ar	Bolton 1995
<i>Crematogaster victimae nitidiceps pergens</i> Forel, 1911	pr pe br	Bolton 1995
<i>Crematogaster victimae nitidiceps turbida</i> Santschi, 1925	br	Bolton 1995
<i>Crematogaster wardi</i> Longino, 2003	cr	Bolton 1995
Tribe Dacetini		
<i>Acanthognathus brevicornis</i> Smith M.R., 1944	br pn	Bolton 2000
<i>Acanthognathus latus</i> Mann, 1922	br ho	Bolton 2000
<i>Acanthognathus ocellatus</i> Mayr, 1887	pe br tt vn co pn cr	Bolton 2000
<i>Acanthognathus poinari*</i> Baroni Urbani, 1994	ad	Bolton 2000
<i>Acanthognathus rufus</i> Brown & Kempf, 1969	pr br	Bolton 2000
<i>Acanthognathus stipulosus</i> Brown & Kempf, 1969	br	Bolton 2000
<i>Acanthognathus teleductus</i> Brown & Kempf, 1969	co pn cr	Bolton 2000
<i>Dacetus armigerum</i> (Latrelle, 1802)	bo pe br gi gf su	Bolton 2000
<i>Pyramica aethegenys</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica abditivata</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica alberti</i> (Forel, 1893)	bo br gi gf su pn cr ni ho me an	Bolton 2000
<i>Pyramica appretiata</i> (Borgmeier, 1954)	br	Bolton 2000
<i>Pyramica asaphes</i> Bolton, 2000	be	Bolton 2000
<i>Pyramica auctidens</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica azteca</i> (Kempf, 1960)	be me	Bolton 2000
<i>Pyramica beebei</i> (Wheeler, 1915)	br co	Bolton 2000
<i>Pyramica brevicornis</i> (Mann, 1922)	pn cr ni be ho me	Bolton 2000
<i>Pyramica brownii</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica cassinispis</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica cincinnata</i> (Kempf, 1975)	br	Bolton 2000
<i>Pyramica comis</i> (Kempf, 1959)	br	Bolton 2000
<i>Pyramica connectens</i> (Kempf, 1958)	ec co	Bolton 2000
<i>Pyramica conspersa</i> (Emery, 1906)	ar br	Bolton 2000
<i>Pyramica convexiceps</i> (Santschi, 1931)	am	Bolton 2000
<i>Pyramica crassicornis</i> (Mayr, 1887)	ar pr br gi tt vn	Bolton 2000
<i>Pyramica crementa</i> Bolton, 2000	pn cr	Bolton 2000
<i>Pyramica dapsilis</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica decipula</i> Bolton, 2000	pe ec br	Bolton 2000
<i>Pyramica deinomastax</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica denticulata</i> (Mayr, 1887)	ar pr bo pe ec br gi gf su tt vn co	Bolton 2000
<i>Pyramica dentinasis</i> (Kempf, 1960)	br	Bolton 2000
<i>Pyramica depressiceps</i> (Weber, 1934)	pe ec gi co pn cr	Bolton 2000

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Pyramica dispalata</i> Bolton, 2000	me	Bolton 2000
<i>Pyramica dontopagis</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica doryceps</i> Bolton, 2000	ec	Bolton 2000
<i>Pyramica eggerti</i> (Emery, 1890)	pr bo pe br tt vn co cr be gu am	Bolton 2000
<i>Pyramica electrina*</i> (De Andrade, 1994)	ad	Bolton 2000
<i>Pyramica emiliae</i> (Forel, 1907)	br	Bolton 2000
<i>Pyramica enopla</i> Bolton, 2000	co	Bolton 2000
<i>Pyramica epinotalis</i> (Weber, 1934)	br cr me	Bolton 2000
<i>Pyramica excisa</i> (Weber, 1934)	cr ni be me	Bolton 2000
<i>Pyramica friderimuelleri</i> (Forel, 1886)	pr pe br tt vn co pn cr	Bolton 2000
<i>Pyramica furtiva</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica gemella</i> (Kempf, 1975)	co	Bolton 2000
<i>Pyramica glenognatha</i> Bolton, 2000	br gf vn	Bolton 2000
<i>Pyramica grytava</i> Bolton, 2000	br co	Bolton 2000
<i>Pyramica gundlachi</i> Roger, 1862	pe ec tt vn co pn cr ni me us am an bh	Bolton 2000
<i>Pyramica halosis</i> Bolton, 2000	vn	Bolton 2000
<i>Pyramica hyphata</i> (Brown, 1953)	br tt vn	Bolton 2000
<i>Pyramica inusitata</i> (Lattke, 1992)	br	Bolton 2000
<i>Pyramica jamaicensis</i> (Brown, 1959)	am	Bolton 2000
<i>Pyramica kompsomala</i> Bolton, 2000	co	Bolton 2000
<i>Pyramica kyidriformis</i> (Brown, 1964)	ar	Bolton 2000
<i>Pyramica laevipleura</i> (Kempf, 1958)	co	Bolton 2000
<i>Pyramica lalassa</i> Bolton, 2000	ec co pn cr	Bolton 2000
<i>Pyramica lilloana</i> (Brown, 1950)	ar br	Bolton 2000
<i>Pyramica longinoi</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica lygatrix</i> Bolton, 2000	ar	Bolton 2000
<i>Pyramica margaritae</i> (Forel, 1893)	tt co pn cr es be me us am bh	Bolton 2000
<i>Pyramica metopia</i> (Brown, 1959)	pe br co pn	Bolton 2000
<i>Pyramica metrix</i> Bolton, 2000	pr	Bolton 2000
<i>Pyramica microthrix</i> (Kempf, 1975)	co pn cr	Bolton 2000
<i>Pyramica minuscula</i> (Kempf, 1962)	br	Bolton 2000
<i>Pyramica mirabilis</i> (Mann, 1926)	bo br	Bolton 2000
<i>Pyramica moloch</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica myllorhapha</i> (Brown, 1959)	pn cr	Bolton 2000
<i>Pyramica necopina</i> Bolton, 2000	ec	Bolton 2000
<i>Pyramica nigrescens</i> (Weeler, 1911)	tt cr ni es be am	Bolton 2000
<i>Pyramica nitens</i> (Santschi, 1932)	am	Bolton 2000
<i>Pyramica nubila</i> (Lattke y Goitia, 1997)	vn co	Bolton 2000
<i>Pyramica orchibia</i> (Brown, 1953)	vn co	Bolton 2000
<i>Pyramica paradoxa</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica parsauga</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica pasiopsis</i> Bolton, 2000	cr	Bolton 2000
<i>Pyramica perissognatha</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica pholidota</i> Bolton, 2000	pe	Bolton 2000
<i>Pyramica prex</i> Bolton, 2000	be	Bolton 2000
<i>Pyramica probatrix</i> (Brown, 1964)	pn cr	Bolton 2000
<i>Pyramica raptans</i> Bolton, 2000	co	Bolton 2000
<i>Pyramica reticeps</i> (Kempf, 1969)	br	Bolton 2000
<i>Pyramica rogata</i> Bolton, 2000	cr me	Bolton 2000
<i>Pyramica rugithorax</i> (Kempf, 1959)	br	Bolton 2000
<i>Pyramica schleeorum*</i> (Baroni Urbani, 1994)	ad	Bolton 2000
<i>Pyramica schulzi</i> (Emery, 1894)	pe br tt vn pn cr me	Bolton 2000
<i>Pyramica simulans</i> (Santschi, 1931)	am	Bolton 2000
<i>Pyramica siagodensis</i> Bolton, 2000	pr	Bolton 2000
<i>Pyramica splendens</i> (Borgmeier, 1954)	ar br	Bolton 2000
<i>Pyramica stauroma</i> Bolton 2000	cr	Bolton 2000
<i>Pyramica stenotes</i> Bolton, 2000	br	Bolton 2000
<i>Pyramica subedentata</i> (Mayr, 1887)	gf su tt vn co pn cr ni be gu me	Bolton 2000
<i>Pyramica sublucida</i> (Brown, 1953)	br	Bolton 2000

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pyramica substricta</i> (Kempf, 1964)	br	Bolton 2000
<i>Pyramica tanymastax</i> (Brown, 1964)	pr br vn	Bolton 2000
<i>Pyramica thaxteri</i> (Wheeler, 1916)	tt	Bolton 2000
<i>Pyramica teratrix</i> Bolton, 2000	co	Bolton 2000
<i>Pyramica tlaloc</i> Bolton, 2000	me	Bolton 2000
<i>Pyramica trieces</i> (Brown, 1960)	pn cr ni	Bolton 2000
<i>Pyramica turpis</i> Bolton, 2000	me	Bolton 2000
<i>Pyramica umbroceps</i> Bolton, 2000	ec	Bolton 2000
<i>Pyramica urrhobia</i> Bolton, 2000	ec br tt co	Bolton 2000
<i>Pyramica vartana</i> Bolton, 2000	co	Bolton 2000
<i>Pyramica villiersi</i> (Perrault, 1986)	br gf tt vn co	Bolton 2000
<i>Pyramica warditeras</i> Bilton, 2000	cr	Bolton 2000
<i>Pyramica wheeleri</i> (M.R. Smith, 1944)	vn co pn cr	Bolton 2000
<i>Pyramica xenochelyna</i> Bolton, 2000	pr br	Bolton 2000
<i>Pyramica xenognatha</i> (Kempf, 1958)	co	Bolton 2000
<i>Pyramica zeteki</i> (Brown, 1959)	pe br co pn cr	Bolton 2000
<i>Strumigenys anthocera</i> Lattke y Goitia, 1997	vn	Bolton 2000
<i>Strumigenys ascita</i> Bolton, 2000	br	Bolton 2000
<i>Strumigenys bolleyi</i> Forel, 1908	ec co pn cr be ho gu me am	Bolton 2000
<i>Strumigenys boneti</i> Brown, 1959	be me us	Bolton 2000
<i>Strumigenys borgmeieri</i> Brown, 1954	br co pn cr	Bolton 2000
<i>Strumigenys calamita</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys carinithorax</i> Borgmeier, 1934	pr br su	Bolton 2000
<i>Strumigenys consanii</i> Brown, 1954	cr	Bolton 2000
<i>Strumigenys cordovensis</i> Mayr, 1887	pr br su tt vn co pn ni be me	Bolton 2000
<i>Strumigenys cosmostela</i> Kempf, 1975	pe br vn co cr me	Bolton 2000
<i>Strumigenys cultrigera</i> Mayr, 1887	br	Bolton 2000
<i>Strumigenys deletrix</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys deltisquama</i> Brown, 1957	co pn cr me	Bolton 2000
<i>Strumigenys diabola</i> Bolton, 2000	br	Bolton 2000
<i>Strumigenys diapteryx</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys dolichognata</i> Weber, 1934	br gi co	Bolton 2000
<i>Strumigenys dubitata</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys dysseides</i> Bolton, 2000	co	Bolton 2000
<i>Strumigenys elongata</i> Roger, 1863	pr bo ec gi su tt vn co pn cr ni be ho gu me	Bolton 2000
<i>Strumigenys emmae</i> (Emery, 1890)	su be us am	Bolton 2000
<i>Strumigenys emeryi</i> Mann, 1922	ho me	Bolton 2000
<i>Strumigenys epelys</i> Bolton, 2000	pr	Bolton 2000
<i>Strumigenys eversa</i> Bolton, 2000	me	Bolton 2000
<i>Strumigenys extirpa</i> Bolton, 2000	pn cr	Bolton 2000
<i>Strumigenys fairchildi</i> Brown, 1961	pn cr	Bolton 2000
<i>Strumigenys formicosa</i> Bolton, 2000	ec	Bolton 2000
<i>Strumigenys godmani</i> Forel, 1899	ec gi co pn cr	Bolton 2000
<i>Strumigenys gytha</i> Bolton, 2000	br	Bolton 2000
<i>Strumigenys hemidisca</i> Brown, 1953	co	Bolton 2000
<i>Strumigenys hindenburgi</i> Forel, 1915	ar br	Bolton 2000
<i>Strumigenys humata</i> Lattke & Goitia, 1997	pn cr	Bolton 2000
<i>Strumigenys idiogenes</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys incuba</i> Bolton, 2000	co	Bolton 2000
<i>Strumigenys infidelis</i> Santschi, 1919	ar bo br vn	Bolton 2000
<i>Strumigenys insolita</i> Bolton, 2000	pr	Bolton 2000
<i>Strumigenys interfectiva</i> Lattke & Goitia, 1997	bo pe vn co	Bolton 2000
<i>Strumigenys lacacoca</i> Brown, 1959	pn	Bolton 2000
<i>Strumigenys lanuginosa</i> Wheeler W.M., 1905	vn co pn cr me am bh	Bolton 2000
<i>Strumigenys longispinosa</i> Brown, 1958	ec pn cr	Bolton 2000
<i>Strumigenys louisianae</i> Roger, 1863	ar pr bo ec br vn co	Bolton 2000
<i>Strumigenys ludia</i> Mann, 1922	cr ni me us am bh	Bolton 2000
<i>Strumigenys mandibularis</i> Smith F., 1860	cr be ho gu me	Bolton 2000
	br	Bolton 2000

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Strumigenys manis</i> Bolton, 2000	me	Bolton 2000
<i>Strumigenys marginiventris</i> Santschi, 1931	vn co pn cr	Bolton 2000
<i>Strumigenys mcretes</i> Brown, 1959	cr	Bolton 2000
<i>Strumigenys mixta</i> Brown, 1953	gu	Bolton 2000
<i>Strumigenys moera</i> Bolton, 2000	ec	Bolton 2000
<i>Strumigenys monstra</i> Bolton, 2000	co	Bolton 2000
<i>Strumigenys nastata</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys nevermanni</i> Brown, 1959	cr me us	Bolton 2000
<i>Strumigenys obliqua</i> Bolton, 2000	ec	Bolton 2000
<i>Strumigenys ogloblini</i> Santschi, 1936	br	Bolton 2000
<i>Strumigenys pariensis</i> Lattke y Goitia, 1997	vn co cr	Bolton 2000
<i>Strumigenys perdita</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys perparva</i> Brown, 1958	bo pe br gi su tt vn co	Bolton 2000
<i>Strumigenys planeti</i> Brown, 1953	bo pe br tt vn	Bolton 2000
<i>Strumigenys platyscapa</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys precava</i> Brown, 1954	bo ec br gi vn co pn cr	Bolton 2000
<i>Strumigenys princeps</i> Kempf & Brown, 1969	br	Bolton 2000
<i>Strumigenys prospiciens</i> Emery, 1906	ar bo pe br vn	Bolton 2000
<i>Strumigenys quadrua</i> Bolton, 2000	br	Bolton 2000
<i>Strumigenys rehi</i> Forel, 1907	br co	Bolton 2000
<i>Strumigenys rogeri</i> Emery, 1890	cr me am	Bolton 2000
<i>Strumigenys ruta</i> Bolton, 2000	br co	Bolton 2000
<i>Strumigenys saliens</i> Mayr, 1887	ar br	Bolton 2000
<i>Strumigenys sanctipauli</i> Kempf, 1958	br	Bolton 2000
<i>Strumigenys schmalzi</i> Emery, 1906	pe br tt	Bolton 2000
<i>Strumigenys sevesta</i> Bolton, 2000	cr	Bolton 2000
<i>Strumigenys silvestrii</i> Emery, 1906	ar br am bh	Bolton 2000
<i>Strumigenys skia</i> Bolton, 2000	ec co	Bolton 2000
<i>Strumigenys smilax</i> Bolton, 2000	br	Bolton 2000
<i>Strumigenys smithii</i> Forel, 1886	br vn co pn cr es be an	Bolton 2000
<i>Strumigenys spathula</i> Lattke & Goitia, 1997	tt vn co pn cr be	Bolton 2000
<i>Strumigenys sublonga</i> Brown, 1958	bo br	Bolton 2000
<i>Strumigenys tachirensis</i> Lattke & Goitia, 1997	vn	Bolton 2000
<i>Strumigenys thomae</i> Kempf, 1976	br	Bolton 2000
<i>Strumigenys timicala</i> Bolton, 2000	ni es	Bolton 2000
<i>Strumigenys tococae</i> Wheeler W.M. & Bequaert, 1929	br gf	Bolton 2000
<i>Strumigenys trinidadensis</i> Wheeler W.M., 1922	br tt co pn cr	Bolton 2000
<i>Strumigenys trudifera</i> Kempf & Brown, 1969	br vn co	Bolton 2000
<i>Strumigenys usbensis</i> Lattke & Goitia, 1997	vn	Bolton 2000
<i>Strumigenys vilhenai</i> Bolton, 2000	br	Bolton 2000
<i>Strumigenys vivax</i> Bolton, 2000	ec	Bolton 2000
<i>Strumigenys xochipili</i> Bolton, 2000	me	Bolton 2000
Tribe Formicoxenini		
<i>Cardiocondyla ectopia</i> Snelling, 1974	us	MacKay 1995
<i>Cardiocondyla emeryi</i> Forel, 1881	br me am an bh	MacKay 1995
<i>Cardiocondyla venustula</i> Wheeler W.M., 1908	us am	MacKay 1995
<i>Nesomyrmex anduzei</i> Weber, 1943	br gi gf su vn co	Kempf 1959
<i>Nesomyrmex argentinus</i> Santschi, 1922	ar	Kempf 1959
<i>Nesomyrmex asper</i> Mayr, 1887	ar br tt vn	Kempf 1959
<i>Nesomyrmex brasiliensis</i> Kempf, 1958	br	Kempf 1959
<i>Nesomyrmex brimodus</i> Kempf, 1959	me	Kempf 1959
<i>Nesomyrmex claviger</i> Wheeler W.M., 1910	br	Kempf 1959
<i>Nesomyrmex costatus</i> Emery, 1896	br	Kempf 1959
<i>Nesomyrmex echinatinodis</i> Forel, 1886	ar pe br gi gf su vn co pn cr ho gu me	Kempf 1959
<i>Nesomyrmex itinerans</i> Kempf, 1959	vn	Kempf 1959
<i>Nesomyrmex mirasolis</i> Diniz, 1975	br	Kempf 1959
<i>Nesomyrmex pittieri</i> Forel, 1899	pn cr ni me	Kempf 1959
<i>Nesomyrmex pleuriticus</i> Kempf, 1959	br gi gf su vn co	Kempf 1959
<i>Nesomyrmex pulcher</i> Emery, 1917	bo	Kempf 1959

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Nesomyrmex rutilans</i> Kempf, 1958	gi gf su	Kempf 1959
<i>Nesomyrmex schwebeli</i> Forel, 1913	br	Kempf 1959
<i>Nesomyrmex sculptiventris</i> Mayr, 1887	br	Kempf 1959
<i>Nesomyrmex spininodis</i> Mayr, 1887	ar pr bo br gi gf su tt vn co	Kempf 1959
<i>Nesomyrmex tonsuratus</i> Emery, 1896	pe tt vn co pn cr	Kempf 1959
<i>Nesomyrmex tristani</i> Mayr, 1887	ar bo br	Kempf 1959
<i>Nesomyrmex vicinus</i> MacKay, 2000	me	Kempf 1959
<i>Nesomyrmex wilda</i> Smith M.R., 1943	me	Kempf 1959
<i>Temnothorax adustus</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax andersoni</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax andrei</i> Emery, 1895	me	MacKay 2000
<i>Temnothorax bicolor</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax bristoli</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax carinatus</i> Cole, 1957	me	MacKay 2000
<i>Temnothorax hispidus</i> Cole, 1957	me	MacKay 2000
<i>Temnothorax manni</i> Wheeler W.M., 1914	me	MacKay 2000
<i>Temnothorax mexicanus</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax peninsularis</i> Wheeler W.M., 1934	me	MacKay 2000
<i>Temnothorax punctaticeps</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax punctatissimus</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax punctithorax</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax rugithorax</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax rugosus</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax rugulosus</i> MacKay, 2000	me	MacKay 2000
<i>Temnothorax stollii</i> Forel, 1885	gu	MacKay 2000
<i>Temnothorax tricarinatus</i> Emery, 1895	me	MacKay 2000
<i>Temnothorax striatulus</i> Stitz, 1937	me	MacKay 2000
<i>Temnothorax whitfordi</i> Kempf, 1959	me	MacKay 2000
<i>Ochetomyrmex semipolitus</i> Mayr, 1878	ar bo br gi gf su co	Fernández 2003a
<i>Ochetomyrmex neopolitus</i> Fernández, 2003	br gi gf su vn co	Fernández 2003a
<i>Xenomyrmex floridanus</i> Emery, 1895	us am bh	Creighton 1957
<i>Xenomyrmex floridanus skwarrae</i> Wheeler W.M., 1931	me	Creighton 1957
<i>Xenomyrmex panamanus</i> (Wheeler W.M., 1922)	pn	Creighton 1957
<i>Xenomyrmex stollii</i> Forel, 1885	gu me	Creighton 1957
Tribe Lenomyrmicini		
<i>Lenomyrmex costatus</i> Fernández & Palacio, 1999	pn	Fernández & Palacio, 1999
<i>Lenomyrmex soveolatus</i> Fernández & Palacio, 1999	co	Fernández & Palacio, 1999
<i>Lenomyrmex inusitatus</i> Fernández, 2001	co	Fernández 2001
<i>Lenomyrmex mandibularis</i> Fernández & Palacio, 1999	co	Fernández & Palacio, 1999
<i>Lenomyrmex wardi</i> Fernández & Palacio, 1999	ec co	Fernández & Palacio, 1999
Tribe Myrmecinini		
<i>Myrmecina americana</i> Emery, 1895	me	Rojas-Fernández 1996
<i>Perissomyrmex snyderi</i> Smith M.R., 1947	gu	Bolton 1995
Tribe Myrmicini		
<i>Hylomyrma balzani</i> (Emery, 1894)	ar pr br	Kempf 1973a
<i>Hylomyrma blandiens</i> Kempf, 1961	pe br gi gf su tt co	Kempf 1973a
<i>Hylomyrma columbica</i> (Forel, 1912)	co	Kempf 1973a
<i>Hylomyrma dentiloba</i> (Santschi, 1931)	pn ni	Kempf 1973a
<i>Hylomyrma longiscapa</i> Kempf, 1961	ec gi gf su	Kempf 1973a
<i>Hylomyrma reitteri</i> (Mayr, 1887)	br vn	Kempf 1973a
<i>Pogonomyrmex apache</i> Wheeler W.M., 1902	us	Taber 1998
<i>Pogonomyrmex atratus</i> Santschi, 1922	ar	Taber 1998
<i>Pogonomyrmex barbatus</i> Smith, 1858	me us	Taber 1998
<i>Pogonomyrmex bicolor</i> Cole, 1968	us	Taber 1998
<i>Pogonomyrmex bispinosus</i> (Spinola, 1851)	ch	Taber 1998
<i>Pogonomyrmex bispinosus intermedia</i> Menozzi, 1935	ch	Taber 1998
<i>Pogonomyrmex bispinosus semistriata</i> Emery, 1906	ch	Taber 1998
<i>Pogonomyrmex bispinosus spinolae</i> Emery, 1906	ch	Taber 1998
<i>Pogonomyrmex brevibarbis</i> Emery, 1906	ar	Taber 1998
<i>Pogonomyrmex brevibarbis niger</i> Santschi, 1925	ar	Taber 1998

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pogonomyrmex brevibarbis silvestrii</i> Emery, 1906	ar	Taber 1998
<i>Pogonomyrmex bruchi</i> Forel, 1913	ar	Taber 1998
<i>Pogonomyrmex californicus</i> Buckley, 1867	us	Taber 1998
<i>Pogonomyrmex carbonarius</i> Mayr, 1868	ar	Taber 1998
<i>Pogonomyrmex carbonarius sanmartini</i> Kusnezov, 1953	ar	Taber 1998
<i>Pogonomyrmex catanilensis</i> Gallardo, 1931	ar	Taber 1998
<i>Pogonomyrmex coarctatus</i> Mayr, 1868	ar ur	Taber 1998
<i>Pogonomyrmex cunicularius</i> Mayr, 1887	ar pr ur bo br	Taber 1998
<i>Pogonomyrmex desertorum</i> Wheeler W.M., 1902	us	Taber 1998
<i>Pogonomyrmex guatemaltecus</i> Wheeler W.M., 1914	gu me	Taber 1998
<i>Pogonomyrmex imberbiculus</i> Wheeler W.M., 1902	us	Taber 1998
<i>Pogonomyrmex inermis</i> Forel, 1914	ar	Taber 1998
<i>Pogonomyrmex laevinodis</i> Snelling, 1982	me	Taber 1998
<i>Pogonomyrmex laticeps</i> Santschi, 1922	ar	Taber 1998
<i>Pogonomyrmex lobatus</i> Santschi, 1921	ar	Taber 1998
<i>Pogonomyrmex longibarbis</i> Gallardo, 1931	ar bo	Taber 1998
<i>Pogonomyrmex longibarbis andinus</i> Kusnezov, 1951	ar	Taber 1998
<i>Pogonomyrmex magnacanthus</i> Cole, 1968	us	Taber 1998
<i>Pogonomyrmex marcusii</i> Kusnezov, 1951	bo	Taber 1998
<i>Pogonomyrmex maricopa</i> Wheeler W.M., 1914	us	Taber 1998
<i>Pogonomyrmex mayri</i> Forel, 1899	co	Taber 1998
<i>Pogonomyrmex meridionalis</i> Kusnezov, 1951	ar	Taber 1998
<i>Pogonomyrmex meridionalis leonis</i> Kusnezov, 1951	ar	Taber 1998
<i>Pogonomyrmex micans</i> Forel, 1914	ar br	Taber 1998
<i>Pogonomyrmex montanus</i> MacKay, 1980	us	Taber 1998
<i>Pogonomyrmex pima</i> Wheeler W.M., 1909	us	Taber 1998
<i>Pogonomyrmex pronotalis</i> Santschi, 1922	ar	Taber 1998
<i>Pogonomyrmex rastratus</i> Mayr, 1868	ar	Taber 1998
<i>Pogonomyrmex rastratus pulchellus</i> Santschi, 1925	ar	Taber 1998
<i>Pogonomyrmex rugosus</i> Emery, 1895	us	Taber 1998
<i>Pogonomyrmex snellingi</i> Taber, 1998		Taber 1998
<i>Pogonomyrmex striatinodus</i> Fernández & Palacio, 1998		Fernández & Palacio, 1998
<i>Pogonomyrmex subnitidus</i> Emery, 1895	us	Taber 1998
<i>Pogonomyrmex sylvestris</i> Lattke, 1990	vn	Taber 1998
<i>Pogonomyrmex tenuispinus</i> Forel, 1914	us	Taber 1998
<i>Pogonomyrmex theresiae</i> Forel, 1899	ec	Taber 1998
<i>Pogonomyrmex uruguayensis</i> Mayr, 1887	ar ur	Taber 1998
<i>Pogonomyrmex variabilis</i> Santschi, 1916	ar	Taber 1998
<i>Pogonomyrmex vermiculatus</i> Emery, 1906	ar ch	Taber 1998
<i>Pogonomyrmex vermiculatus chubutensis</i> Forel, 1913	ar	Taber 1998
<i>Pogonomyrmex wheeleri</i> Olsen, 1934	me	Taber 1998
Tribe Phalacromyrmecini		
<i>Phalacromyrmex fugax</i> Kempf, 1960	br	Kempf 1960
Tribe Pheidolini		
<i>Aphaenogaster araneoides</i> Emery, 1890	cr	Bolton 1995
<i>Aphaenogaster araneoides canalis</i> Enzmann J., 1947	pn	Bolton 1995
<i>Aphaenogaster araneoides inermis</i> Forel, 1899	pn	Bolton 1995
<i>Aphaenogaster araneoides nitidiventris</i> Forel, 1912	cr	Bolton 1995
<i>Aphaenogaster ensifera</i> Forel, 1899	me	Bolton 1995
<i>Aphaenogaster honduriana</i> Mann, 1922	ho	Bolton 1995
<i>Aphaenogaster mexicana</i> (Pergande, 1896)	me	Bolton 1995
<i>Aphaenogaster phalangium</i> Emery, 1890	cr ho	Bolton 1995
<i>Aphaenogaster phalangium brevicollis</i> Forel, 1899	pn	Bolton 1995
<i>Messor andrei</i> Mayr, 1886	us	Rojas-Fernández 1996
<i>Messor julianus</i> (Pergande, 1894)	me	Rojas-Fernández 1996
<i>Messor pergandei</i> (Mayr, 1886)	us	Rojas-Fernández 1996
<i>Messor stoddardi</i> Emery, 1895	us	Rojas-Fernández 1996
<i>Pheidole aberrans</i> Mayr, 1868	ar bo pe br	Wilson 2003
<i>Pheidole absurda</i> Forel, 1886	cr ni ho gu me us	Wilson 2003
<i>Pheidole accinota</i> Wheeler W.M., 1925	pe	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole aciculata</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole ademonia</i> Wilson, 2003	pe ec	Wilson 2003
<i>Pheidole alacris</i> Santschi, 1923	ar	Wilson 2003
<i>Pheidole agricola</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole albipes</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole alexeter</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole alfaroi</i> Emery, 1896	co pn cr	Wilson 2003
<i>Pheidole allarmata</i> Wilson, 2003	pe br su co	Wilson 2003
<i>Pheidole alpestris</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole alpinensis</i> Forel, 1912	br	Wilson 2003
<i>Pheidole alticola</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole amabilis</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole amata</i> Forel, 1901	co	Wilson 2003
<i>Pheidole amazonica</i> Wilson, 2003	pe ec	Wilson 2003
<i>Pheidole ambigua</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole androsana</i> Wheeler W.M., 1905	bh	Wilson 2003
<i>Pheidole angulifera</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole angusta</i> Forel, 1908	br	Wilson 2003
<i>Pheidole angusticeps</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole anima</i> Wilson, 2003	ho	Wilson 2003
<i>Pheidole antillana</i> Forel, 1893	an	Wilson 2003
<i>Pheidole aper</i> Forel, 1912	br	Wilson 2003
<i>Pheidole arachnion</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole arboricola</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole arctos</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole arhuaca</i> Forel, 1901	br co	Wilson 2003
<i>Pheidole ariei</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole arietans</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole aripoensis</i> Wilson, 2003	tt	Wilson 2003
<i>Pheidole asperithorax</i> Emery, 1894	bo pe br	Wilson 2003
<i>Pheidole astur</i> Wilson, 2003	bo pe ec br gi gf su co	Wilson 2003
<i>Pheidole atticola</i> Forel, 1912	ar	Wilson 2003
<i>Pheidole auropilosa</i> Mayr, 1887	ar br	Wilson 2003
<i>Pheidole avia</i> Forel, 1908	br	Wilson 2003
<i>Pheidole azteca</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole bakeri</i> Forel, 1912	am	Wilson 2003
<i>Pheidole balzani</i> Emery, 1894	bo vn	Wilson 2003
<i>Pheidole bambusarum</i> Forel, 1908	ar br	Wilson 2003
<i>Pheidole bellatrix</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole belocephala</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole biconstricta</i> Mayr, 1870	bo pe br tt vn co pn cr ni be gu	Wilson 2003
<i>Pheidole bicarinata</i> Forel, 1899	pn cr	Wilson 2003
<i>Pheidole bidens</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole bilimeki</i> Mayr, 1870	vn co pn cr ni ho gu me am bh	Wilson 2003
<i>Pheidole binasifer</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole biolleyi</i> Forel, 1908	cr	Wilson 2003
<i>Pheidole bison</i> Wilson, 2003	bo	Wilson 2003
<i>Pheidole blumenauensis</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole boliviiana</i> Wilson, 2003	bo	Wilson 2003
<i>Pheidole boltoni</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole borgmeieri</i> Kempf, 1972	br	Wilson 2003
<i>Pheidole boruca</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole brachyops</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole brandaoi</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole brevicona</i> Mayr, 1887	br	Wilson 2003
<i>Pheidole breviscapa</i> Forel, 1899	gu	Wilson 2003
<i>Pheidole breviseta</i> Santschi, 1919	ar	Wilson 2003
<i>Pheidole browni</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole bruchella</i> Forel, 1915	ar	Wilson 2003
<i>Pheidole bruesi</i> Wheeler, 1911	br an	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole brunnescens</i> Santschi, 1929	br	Wilson 2003
<i>Pheidole bucculenta</i> Forel, 1908	br	Wilson 2003
<i>Pheidole bulliceps</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole calens</i> Forel, 1901	pe br gi vn co pn cr ho	Wilson 2003
<i>Pheidole calimana</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole cameroni</i> Mayr, 1887	ec	Wilson 2003
<i>Pheidole camilla</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole campostela</i> Kempf, 1972	br	Wilson 2003
<i>Pheidole capillata</i> Emery, 1906	pe br	Wilson 2003
<i>Pheidole caracalla</i> Wilson, 2003	pe ec	Wilson 2003
<i>Pheidole carapunco</i> Kusnezov, 1952	ar	Wilson 2003
<i>Pheidole cardiella</i> Wilson, 2003	pe ec	Wilson 2003
<i>Pheidole carinata</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole carapuna</i> Wilson, 2003	pe br gi vn co pn cr ho	Wilson 2003
<i>Pheidole cataphracta</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole cataractae</i> Wheeler W.M., 1916	pe br	Wilson 2003
<i>Pheidole caulincola</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole cavifrons</i> Wilson, 2003	ar ur br	Wilson 2003
<i>Pheidole ceibana</i> Wilson, 2003	ho	Wilson 2003
<i>Pheidole celaena</i> Wilson, 2003	cr ni	Wilson 2003
<i>Pheidole centeotl</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole cephalica</i> Wilson, 2003	bo br tt co pn cr ni gu me	Wilson 2003
<i>Pheidole cerebresior</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole cerina</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole chalca</i> Wheeler W.M., 1914	me	Wilson 2003
<i>Pheidole chalcooides</i> Wilson, 2003	ec me	Wilson 2003
<i>Pheidole charazana</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole chilensis</i> Mayr, 1862	ch pe	Wilson 2003
<i>Pheidole chloe</i> Forel, 1908	cr me	Wilson 2003
<i>Pheidole christopherseni</i> Forel, 1912	co pn cr	Wilson 2003
<i>Pheidole chrysops</i> Wilson, 2003	pr br	Wilson 2003
<i>Pheidole cielana</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole citrina</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole claviscapa</i> Santschi, 1925	bo br	Wilson 2003
<i>Pheidole cocciphaga</i> Borgmeier, 1934	gi su co cr am	Wilson 2003
<i>Pheidole coffeicicola</i> Wilson, 2003	br su	Wilson 2003
<i>Pheidole colobopsis</i> Mann, 1916	pe br co	Wilson 2003
<i>Pheidole coracina</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole cornicula</i> Wilson, 2003	bo	Wilson 2003
<i>Pheidole coveri</i> Wilson, 2003	tt vn	Wilson 2003
<i>Pheidole cramptoni</i> Wilson, 2003	pe ec co pn cr	Wilson 2003
<i>Pheidole crinita</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole crozieri</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole cubensis</i> Mayr, 1862	am	Wilson 2003
<i>Pheidole cuprina</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole cyrtostela</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole daphne</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole dasypyx</i> Wilson, 2003	pn	Wilson 2003
<i>Pheidole davidsonae</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole davisi</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole deceptrix</i> Forel, 1899	gu	Wilson 2003
<i>Pheidole defecta</i> Santschi, 1923	gu	Wilson 2003
<i>Pheidole delicata</i> Wilson, 2003	bo	Wilson 2003
<i>Pheidole demeter</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole descolei</i> Kusnezov, 1952	ar br	Wilson 2003
<i>Pheidole diabolus</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole diana</i> Forel, 1908	cr	Wilson 2003
<i>Pheidole diligens</i> Smith F., 1858	pe br	Wilson 2003
<i>Pheidole dinophila</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole dione</i> Forel, 1913	ar	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole dorsata</i> Wilson, 2003	am	Wilson 2003
<i>Pheidole drepanon</i> Wilson, 2003	am	Wilson 2003
<i>Pheidole dryas</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole dumicola</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole dwyeri</i> Gregg, 1969	me	Wilson 2003
<i>Pheidole dyctiota</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole ebenina</i> Wilson, 2003	co cr	Wilson 2003
<i>Pheidole ectatommoides</i> Wilson, 2003	ec co pn cr	Wilson 2003
<i>Pheidole ecuadorana</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole embolopyx</i> Brown, 1968	pe ec br	Wilson 2003
<i>Pheidole eparmata</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole erethizon</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole eriophora</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole erratilis</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole euruscopa</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole exarata</i> Emery, 1896	co cr	Wilson 2003
<i>Pheidole excubitor</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole exigua</i> Mayr, 1884	bo br gi gf tt co pn cr am	Wilson 2003
<i>Pheidole exquisita</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole fabricator</i> Smith F., 1858	br vn co	Wilson 2003
<i>Pheidole fariasana</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole fera</i> Santschi, 1925	br	Wilson 2003
<i>Pheidole fimbriata</i> Wilson, 2003	ar pe ec br gi gf su tt vn co pn cr ni ho gu me	Wilson 2003
<i>Pheidole fissiceps</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole flavens</i> Roger, 1863	ar pr pe br tt co pn cr me us am an bh	Wilson 2003
<i>Pheidole flavidula</i> Mayr, 1887	br	Wilson 2003
<i>Pheidole flavifrons</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole floricola</i> Wilson, 2003	ho	Wilson 2003
<i>Pheidole fowleri</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole fracticeps</i> Wilson, 2003	pr pe ec br tt	Wilson 2003
<i>Pheidole fullerae</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole gagates</i> Wilson, 2003	pe ec	Wilson 2003
<i>Pheidole gaigei</i> Forel, 1914	co	Wilson 2003
<i>Pheidole gangamon</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole gauthieri</i> Forel, 1901	co pn cr ho gu	Wilson 2003
<i>Pheidole geminata</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole gemmula</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole geraensis</i> Santschi, 1929	br	Wilson 2003
<i>Pheidole germaini</i> Emery, 1896	br	Wilson 2003
<i>Pheidole gertrudae</i> Forel, 1886	ar pr bo br	Wilson 2003
<i>Pheidole gibba</i> Mayr, 1887	br	Wilson 2003
<i>Pheidole gibbata</i> Borgmeier, 1934	su vn co	Wilson 2003
<i>Pheidole gigaslavens</i> Wilson, 2003	ar bo	Wilson 2003
<i>Pheidole gilva</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole globularia</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole glomericeps</i> Wilson, 2003	cr me	Wilson 2003
<i>Pheidole gnomus</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole godmani</i> Forel, 1893	an	Wilson 2003
<i>Pheidole goeldii</i> Forel, 1896	br	Wilson 2003
<i>Pheidole gradifer</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole grandinodus</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole grantae</i> Forel, 1908	co cr	Wilson 2003
<i>Pheidole gravida</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole grex</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole guajiranana</i> Wilson, 2003	br vn co	Wilson 2003
<i>Pheidole guayasana</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole guerreriana</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole guilelmimuelleri</i> Forel, 1886	ar br	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole glyphoderma</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole harlequina</i> Wilson, 2003	am	Wilson 2003
<i>Pheidole hasticeps</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole hazenae</i> Wilson, 2003	ec co	Wilson 2003
<i>Pheidole harrisonfordi</i> Wilson, 2003	be ho	Wilson 2003
<i>Pheidole hecate</i> Wheeler W.M., 1911	am	Wilson 2003
<i>Pheidole heyeri</i> Forel, 1899	br	Wilson 2003
<i>Pheidole heterothrix</i> Wilson, 2003	ar	Wilson 2003
<i>Pheidole hetschkoii</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole hierax</i> Wilson, 2003	pn	Wilson 2003
<i>Pheidole hispaniolae</i> Wilson, 2003	am	Wilson 2003
<i>Pheidole hirtula</i> Forel, 1899	me	Wilson 2003
<i>Pheidole hizemops</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole hoelldobleri</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole horribilis</i> Wilson, 2003	pe ec	Wilson 2003
<i>Pheidole hortonae</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole huilana</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole humida</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole inca</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole incisa</i> Mayr, 1870	co	Wilson 2003
<i>Pheidole indistincta</i> Forel, 1899	cr	Wilson 2003
<i>Pheidole inermis</i> Wilson, 2003	ni ho me	Wilson 2003
<i>Pheidole infernalis</i> Wilson, 2003	bo	Wilson 2003
<i>Pheidole inversa</i> Forel, 1901	co an	Wilson 2003
<i>Pheidole iracunda</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole jeannei</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole jivaro</i> Wilson, 2003	ec pn	Wilson 2003
<i>Pheidole kuna</i> Wilson, 2003	pn	Wilson 2003
<i>Pheidole laelaps</i> Wilson, 2003	ec co	Wilson 2003
<i>Pheidole laevifrons</i> Mayr, 1887	br	Wilson 2003
<i>Pheidole laeviventris</i> Mayr, 1870	co	Wilson 2003
<i>Pheidole laevinota</i> Forel, 1908	ar br	Wilson 2003
<i>Pheidole laidlowi</i> Mann, 1916	pe br	Wilson 2003
<i>Pheidole laevifrons</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole lancifer</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole laselva</i> Wilson, 2003	ec vn co pn cr	Wilson 2003
<i>Pheidole laticrista</i> Santschi, 1916	ar	Wilson 2003
<i>Pheidole laticornis</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole laudatana</i> Wilson, 2003	an	Wilson 2003
<i>Pheidole lemnisca</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole lemur</i> Forel, 1912	br	Wilson 2003
<i>Pheidole lignicola</i> Mayr, 1887	ar pr br	Wilson 2003
<i>Pheidole lilloi</i> Wilson, 2003	ar	Wilson 2003
<i>Pheidole longinoi</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole longior</i> Wilson, 2003	ar	Wilson 2003
<i>Pheidole longiscapa</i> Wilson, 2003	br vn	Wilson 2003
<i>Pheidole longiseta</i> Wilson, 2003	pe br	Wilson 2003
<i>Pheidole lucaris</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole lustrata</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole lutzi</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole machetula</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole mackayi</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole mallota</i> Wilson, 2003	pn	Wilson 2003
<i>Pheidole mamore</i> Mann, 1916	pe br gi gf su	Wilson 2003
<i>Pheidole manuana</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole medialis</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole megacephala</i> (Fabricius, 1793)	pe br gi gf su vn co pn cr ni ho gu me us am an bh	Wilson 2003
<i>Pheidole meinerti</i> Forel, 1905	pe ec br gi vn	Wilson 2003
<i>Pheidole meinertopsis</i> Wilson, 2003	pe	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole melastomae</i> Wilson, 2003	ec co pn	Wilson 2003
<i>Pheidole mendicula</i> Wheeler W.M., 1925	pe ec br tt co pn cr	Wilson 2003
<i>Pheidole metana</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole mincana</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole micridis</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole micon</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole microps</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole minensis</i> Santschi, 1923	br	Wilson 2003
<i>Pheidole minutula</i> Mayr, 1878	pe ec br gf vn co pn	Wilson 2003
<i>Pheidole mirabilis</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole mittermeieri</i> Wilson, 2003	bo	Wilson 2003
<i>Pheidole mixteca</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole moerens</i> Wheeler W.M., 1908	us am	Wilson 2003
<i>Pheidole moffetti</i> Wilson, 2003	gf	Wilson 2003
<i>Pheidole monteverdensis</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole morelosana</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole moseni</i> Wheeler W.M., 1925	br	Wilson 2003
<i>Pheidole mosenopsis</i> Wilson, 2003	pr bo	Wilson 2003
<i>Pheidole multispina</i> Wilson, 2003	pn cr	Wilson 2003
<i>Pheidole nana</i> Emery, 1894	br	Wilson 2003
<i>Pheidole napoensis</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole nasifer</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole nasutoides</i> Hölldobler & Wilson, 1992	cr	Wilson 2003
<i>Pheidole naylae</i> Wilson, 2003	am	Wilson 2003
<i>Pheidole nebulosa</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole neolongiceps</i> Wilson, 2003	am	Wilson 2003
<i>Pheidole nigella</i> Emery, 1894	bo pe	Wilson 2003
<i>Pheidole nigricula</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole nitella</i> Wilson, 2003	ec br co pn cr	Wilson 2003
<i>Pheidole nitidicollis</i> Emery, 1896	br cr am	Wilson 2003
<i>Pheidole nitidula</i> Emery, 1888	ar br	Wilson 2003
<i>Pheidole noar</i> Wilson, 2003	am	Wilson 2003
<i>Pheidole nubila</i> Emery, 1906	ar	Wilson 2003
<i>Pheidole nubicola</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole oaxacana</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole obnixa</i> Forel, 1912	ar	Wilson 2003
<i>Pheidole obrima</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole obscurifrons</i> Santschi, 1925	ar	Wilson 2003
<i>Pheidole obtusopilosa</i> Mayr, 1887	ar ur br	Wilson 2003
<i>Pheidole obtusospinosa</i> Pergande, 1896	me us	Wilson 2003
<i>Pheidole olsoni</i> Wilson, 2003	pn cr	Wilson 2003
<i>Pheidole onyx</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole orbica</i> Forel, 1893	am an	Wilson 2003
<i>Pheidole otisi</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole oxyops</i> Forel, 1908	ar pr br	Wilson 2003
<i>Pheidole palenquensis</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole pampana</i> Santschi, 1929	ar	Wilson 2003
<i>Pheidole pariana</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole peckorum</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole pedana</i> Wilson, 2003	pe br	Wilson 2003
<i>Pheidole peltastes</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole pepo</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole perdiligens</i> Wilson, 2003	pn cr	Wilson 2003
<i>Pheidole peregrina</i> Wheeler W.M., 1916	br	Wilson 2003
<i>Pheidole perkinsi</i> Wilson, 2003	an	Wilson 2003
<i>Pheidole perpusilla</i> Emery, 1894	br gi co pn	Wilson 2003
<i>Pheidole peruviana</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole petersoni</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole pholeops</i> Wilson, 2003	pe ec br	Wilson 2003
<i>Pheidole piceonigra</i> Wilson, 2003	cr me	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole pidax</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole pilifera</i> Wilson, 2003	us	Wilson 2003
<i>Pheidole pilispina</i> Wilson, 2003	pn	Wilson 2003
<i>Pheidole pinealis</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole pinicola</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole plato</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole polita</i> Emery, 1894	bo pe br	Wilson 2003
<i>Pheidole polymorpha</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole praeaes</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole pratatorum</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole primigenia*</i> Baroni Urbani, 1995	ad	Wilson 2003
<i>Pheidole prolixa</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole prostrata</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole psammophila</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole pubiventris</i> Mayr, 1887	br co	Wilson 2003
<i>Pheidole pugnax</i> Wilson, 2003	pn cr es ho	Wilson 2003
<i>Pheidole pullula</i> Wilson, 2003	ar	Wilson 2003
<i>Pheidole punctatissima</i> Mayr, 1870	co pn cr ni ho gu me	Wilson 2003
<i>Pheidole pygmaea</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole quadriiceps</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole quercicola</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole quiaccana</i> Wheeler, 1925	bo	Wilson 2003
<i>Pheidole radoszkowskii</i> Mayr, 1884	ar pr bo pe br gi gf su vn co pn cr ni gu am an	Wilson 2003
<i>Pheidole rectiluma</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole rectisentis</i> Wilson, 2003	ni	Wilson 2003
<i>Pheidole rectispina</i> Wilson, 2003	cr be	Wilson 2003
<i>Pheidole rectitrudis</i> Wilson, 2003	cr me	Wilson 2003
<i>Pheidole reichenspergeri</i> Santschi, 1923	br	Wilson 2003
<i>Pheidole renae</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole rhea</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole rhinoceros</i> Forel, 1899	pn cr ni ho	Wilson 2003
<i>Pheidole rhytidera</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole risii</i> Forel, 1892	ar br	Wilson 2003
<i>Pheidole riveti</i> Santschi, 1911	ec	Wilson 2003
<i>Pheidole rogeri</i> Emery, 1896	cr	Wilson 2003
<i>Pheidole rochae</i> Forel, 1912	br	Wilson 2003
<i>Pheidole rosae</i> Wilson, 2003	ar br	Wilson 2003
<i>Pheidole rosula</i> Wilson, 2003	ar	Wilson 2003
<i>Pheidole rotundiceps</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole rudigenis</i> Emery, 1906	ar br	Wilson 2003
<i>Pheidole rufipilis</i> Forel, 1908	br	Wilson 2003
<i>Pheidole rugatula</i> Santschi, 1933	ar	Wilson 2003
<i>Pheidole rugiceps</i> Wilson, 2003	pe br tt pn cr be ho gu	Wilson 2003
<i>Pheidole ruida</i> Wilson, 2003	pn cr	Wilson 2003
<i>Pheidole rutilana</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole sabella</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole sabina</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole sagana</i> Wheeler W.M., 1934	cr me	Wilson 2003
<i>Pheidole sagax</i> Wilson, 2003	co ni an	Wilson 2003
<i>Pheidole sagittaria</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole sarcina</i> Forel, 1912	ec br	Wilson 2003
<i>Pheidole sarpedon</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole scabriventralis</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole scalaris</i> Wilson, 2003	tt cr	Wilson 2003
<i>Pheidole schmalzi</i> Emery, 1894	bo br vn	Wilson 2003
<i>Pheidole schwarzmaieri</i> Borgmeier, 1939	ar br	Wilson 2003
<i>Pheidole scimitarra</i> Wilson, 2003	pe	Wilson 2003
<i>Pheidole sciophila</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole scolioceps</i> Wilson, 2003	pe br	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole scrobifera</i> Emery, 1896	cr	Wilson 2003
<i>Pheidole sculptior</i> Forel, 1893	su tt vn am an	Wilson 2003
<i>Pheidole securiger</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole seedrayersi</i> Forel, 1910	ar bo co	Wilson 2003
<i>Pheidole scitula</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole seligmani</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole senilis</i> Santschi, 1929	br	Wilson 2003
<i>Pheidole sensitiva</i> Borgmeier, 1959	br co pn cr	Wilson 2003
<i>Pheidole servilia</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole setsukoae</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole sicaria</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole sigillata</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole similigena</i> Wheeler W.M., 1937	am	Wilson 2003
<i>Pheidole simonsi</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole simplex</i> Wheeler, 1925	pe br	Wilson 2003
<i>Pheidole sitarches</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole spadonia</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole spathicornis</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole specularis</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole sphaerica</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole spilota</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole spininodis</i> Wilson, 2003	ar br	Wilson 2003
<i>Pheidole socrates</i> Forel, 1912	br gi gf su tt	Wilson 2003
<i>Pheidole sospes</i> Forel, 1908	ar pe br	Wilson 2003
<i>Pheidole steinheili</i> Forel, 1901	co	Wilson 2003
<i>Pheidole stigma</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole stomachosa</i> Wheeler W.M., 1917	am	Wilson 2003
<i>Pheidole striaticeps</i> Mayr, 1870	pe cr ho me	Wilson 2003
<i>Pheidole strigosa</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole stulta</i> Wilson, 2003	br pn	Wilson 2003
<i>Pheidole styrax</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole subaberrans</i> Kusnezov, 1952	ar	Wilson 2003
<i>Pheidole subarmata</i> Mayr, 1884	br co pn cr me am an bh	Wilson 2003
<i>Pheidole subnuda</i> Wilson, 2003	vn co	Wilson 2003
<i>Pheidole subsphaerica</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole superba</i> Wilson, 2003	co pn	Wilson 2003
<i>Pheidole synarmata</i> Wilson, 2003	br gi	Wilson 2003
<i>Pheidole tachigaliae</i> Wheeler W.M., 1921	gi gf	Wilson 2003
<i>Pheidole tachirana</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole tambopatae</i> Wilson, 2003	pe br co	Wilson 2003
<i>Pheidole tanyscapa</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole tarchon</i> Wilson, 2003	ar	Wilson 2003
<i>Pheidole taurus</i> Emery, 1906	ar	Wilson 2003
<i>Pheidole tayrona</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole tenebra</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole tenerescens</i> Wheeler W.M., 1922	tt	Wilson 2003
<i>Pheidole teneriffana</i> Forel, 1893	pe us am	Wilson 2003
<i>Pheidole tennantae</i> Wilson, 2003	pe cr	Wilson 2003
<i>Pheidole tenuis</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole tepicana</i> Pergande, 1896	me us	Wilson 2003
<i>Pheidole tepuicola</i> Wilson, 2003	vn	Wilson 2003
<i>Pheidole termitobia</i> Forel, 1901	br	Wilson 2003
<i>Pheidole terresi</i> Wheeler W.M. & Mann, 1914	am an	Wilson 2003
<i>Pheidole terribilis</i> Wilson, 2003	bo br su	Wilson 2003
<i>Pheidole taihepa*</i> Wilson, 1985	ad	Wilson 2003
<i>Pheidole tetrica</i> Forel, 1913	br	Wilson 2003
<i>Pheidole texticeps</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole tigris</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole tillandsiarum</i> Wheeler W.M., 1934	me	Wilson 2003
<i>Pheidole tisiphone</i> Wheeler W.M., 1911	gu me	Wilson 2003

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Pheidole titanis</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole torosa</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole trachyderma</i> Emery, 1906	ar	Wilson 2003
<i>Pheidole tragica</i> Wheeler W.M., 1934	me	Wilson 2003
<i>Pheidole transversostriata</i> Mayr, 1887	br gi gf su tt co an	Wilson 2003
<i>Pheidole triconstricta</i> Forel, 1886	ar pr ur bo br	Wilson 2003
<i>Pheidole trinitatis</i> Wilson, 2003	tt	Wilson 2003
<i>Pheidole tristicula</i> Wilson, 2003	bo pe ec br gf tt co ho me	Wilson 2003
<i>Pheidole tristis</i> (Smith F., 1858)	ar br	Wilson 2003
<i>Pheidole tristops</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole triumbonata</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole truncula</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole tschinkelii</i> Wilson, 2003	gu	Wilson 2003
<i>Pheidole tuxtlasana</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole tysoni</i> Wilson, 2003	me us	Wilson 2003
<i>Pheidole ulothrix</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole umphreyi</i> Wilson, 2003	pn	Wilson 2003
<i>Pheidole unicornis</i> Wilson, 2003	ec co	Wilson 2003
<i>Pheidole ursus</i> Mayr, 1870	me	Wilson 2003
<i>Pheidole vafella</i> Wheeler, 1925	bo	Wilson 2003
<i>Pheidole vafra</i> Santschi, 1923	ar pr br tt vn co pn cr ni	Wilson 2003
<i>Pheidole vallifica</i> Forel, 1901	pr ho	Wilson 2003
<i>Pheidole variceps</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole variegata</i> Emery, 1896	cr	Wilson 2003
<i>Pheidole vaslitii</i> Pergande, 1896	bo br pn	Wilson 2003
<i>Pheidole veletis</i> Wilson, 2003	bo br co pn	Wilson 2003
<i>Pheidole venatrix</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole verricula</i> Wilson, 2003	cr ho	Wilson 2003
<i>Pheidole victima</i> Wilson, 2003	br	Wilson 2003
<i>Pheidole viriosa</i> Wilson, 2003	ec	Wilson 2003
<i>Pheidole violacea</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole vestita</i> Wilson, 2003	cr	Wilson 2003
<i>Pheidole vomer</i> Wilson, 2003	co	Wilson 2003
<i>Pheidole walkeri</i> Mann, 1922	ho	Wilson 2003
<i>Pheidole wardi</i> Wilson, 2003	ho	Wilson 2003
<i>Pheidole weiseri</i> Wilson, 2003	ar	Wilson 2003
<i>Pheidole williamsi</i> Wheeler W.M., 1919	ec	Wilson 2003
<i>Pheidole xerophila</i> Wilson, 2003	us	Wilson 2003
<i>Pheidole xyston</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole yucatana</i> Wilson, 2003	me	Wilson 2003
<i>Pheidole zelata</i> Wilson, 2003	bo	Wilson 2003
<i>Pheidole zeteki</i> Smith M.R., 1947	pe ec tt co pn	Wilson 2003
<i>Pheidole zoster</i> Wilson, 2003	pe	Wilson 2003
Tribe Solenopsidini		
<i>Allomerus decemarticulatus</i> Mayr, 1878	br	Bolton 1995
<i>Allomerus decemarticulatus novemarticulatus</i> Wheeler W.M. & Mann, 1942	bo	Bolton 1995
<i>Allomerus octoarticulatus</i> Mayr, 1878	bo br	Bolton 1995
<i>Allomerus octoarticulatus angulatus</i> Wheeler W.M., 1942	bo	Bolton 1995
<i>Allomerus octoarticulatus demerarae</i> Wheeler G.C., 1935	gi	Bolton 1995
<i>Allomerus octoarticulatus exsanguis</i> Wheeler W.M., 1942	bo	Bolton 1995
<i>Allomerus octoarticulatus melanoticus</i> Wheeler W.M., 1942	bo	Bolton 1995
<i>Allomerus octoarticulatus septemarticulatus</i> Mayr, 1878	br	Bolton 1995
<i>Allomerus octoarticulatus tuberculatus</i> Forel, 1912	br	Bolton 1995
<i>Carebara angulata</i> Fernández, 2004	co ec	Fernández 2004a
<i>Carebara anopthalma</i> (Emery, 1906)	br	Fernández 2004a
<i>Carebara audita</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara bicarinata</i> Santschi, 1912	gf	Fernández 2004a
<i>Carebara brasiliensis</i> Fernández, 2004	br	Fernández 2004a
<i>Carebara brevipilosa</i> Fernández, 2004	cr - br	Fernández 2004a

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Carebara bruchi</i> (Santschi, 1933)	ar	Fernández 2004a
<i>Carebara coeca</i> Fernández, 2003	br co	Fernández 2004a
<i>Carebara elongata</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara inca</i> Fernández, 2004	cr pe	Fernández 2004a
<i>Carebara incierta</i> Fernández, 2004	br	Fernández 2004a
<i>Carebara intermedia</i> Fernández, 2004	be	Fernández 2004a
<i>Carebara globularia</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara kofana</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara longiceps</i> (Santschi, 1929)	ar	Fernández 2004a
<i>Carebara majeri</i> Fernández, 2004	br	Fernández 2004a
<i>Carebara minuta</i> Fernández, 2004	br	Fernández 2004a
<i>Carebara mayri</i> Forel, 1901	pr br gi gf su	Fernández 2004a
<i>Carebara nuda</i> Fernández, 2004	br	Fernández 2004a
<i>Carebara paya</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara panamensis</i> (Wheeler, 1925)	pn	Fernández 2004a
<i>Carebara peruviana</i> (Emery, 1906)	pe	Fernández 2004a
<i>Carebara pilosa</i> Fernández, 2004	br	Fernández 2004a
<i>Carebara reina</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara reticulata</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara semistriata</i> Fernández, 2004	co	Fernández 2004a
<i>Carebara striata</i> Fernández, 2004	co tt br	Fernández 2004a
<i>Carebara stenopterus</i> (Kusnezov, 1952)	ar	Fernández 2004a
<i>Carebara tenua</i> Fernández, 2004	br	Fernández 2004a
<i>Carebara urichi</i> (Wheeler, 1922)	gi gf su tt me am	Fernández 2004a
<i>Carebara winifredae</i> Wheeler W.M., 1922	gi	Fernández 2004a
<i>Megalomyrmex balzani</i> Emery, 1894	bo pe br co	Brandão 1990 2003
<i>Megalomyrmex bituberculatus</i> (Fabricius, 1798)	br gi gf su tt	Brandão 1990 2003
<i>Megalomyrmex drifti</i> Kempf, 1961	pe ec br gi gf su tt co pn cr me	Brandão 1990 2003
<i>Megalomyrmex emeryi</i> Forel, 1904	gi gf su	Brandão 1990 2003
<i>Megalomyrmex foreli</i> Emery, 1890	co pn cr	Brandão 1990 2003
<i>Megalomyrmex glaesarius</i> Kempf, 1970	pe ec	Brandão 1990 2003
<i>Megalomyrmex gnomus</i> Kempf, 1970	br	Brandão 1990 2003
<i>Megalomyrmex goeldii</i> Forel, 1912	br	Brandão 1990 2003
<i>Megalomyrmex incisus</i> Smith R., 1947	pe ec br tt vn co pn me	Brandão 1990 2003
<i>Megalomyrmex iheringi</i> Forel, 1911	br	Brandão 1990 2003
<i>Megalomyrmex latreillei</i> Emery, 1890	ec br co	Brandão 1990 2003
<i>Megalomyrmex leoninus</i> Forel, 1885	br gi gf su tt vn co	Brandão 1990 2003
<i>Megalomyrmex modestus</i> Emery, 1896	vn co pn cr	Brandão 1990 2003
<i>Megalomyrmex myops</i> Santschi, 1925	br	Brandão 1990 2003
<i>Megalomyrmex nasutus</i> Forel, 1912	br	Brandão 1990 2003
<i>Megalomyrmex pusillus</i> Forel, 1912	br	Brandão 1990 2003
<i>Megalomyrmex silvestrii</i> Wheeler W.M., 1909	ar pr bo pe br gi gf su tt vn co pn cr ho me	Brandão 1990 2003
<i>Megalomyrmex staudingeri</i> Emery, 1890	pe br co	Brandão 1990 2003
<i>Megalomyrmex symmetochus</i> Wheeler W.M., 1925	bo br pn cr	Brandão 1990 2003
<i>Megalomyrmex wallacei</i> Mann, 1916	br	Brandão 1990 2003
<i>Megalomyrmex wetteri</i> Brandão, 2003	cr pn	Brandão 2003
<i>Megalomyrmex weyrauchi</i> Kempf, 1970	pe	Brandão 1990 2003
<i>Monomorium bicolor</i> (Ettershank, 1965)	ch	Fernández, 2004
<i>Monomorium bidentatum</i> Mayr, 1887	ch	Fernández, 2004
<i>Monomorium brasiliense</i> Forel, 1908	br	Fernández, 2004
<i>Monomorium cekalovici</i> (Snelling, 1975)	ch	Fernández, 2004
<i>Monomorium carbonarium</i> Smith F., 1858	az	Fernández, 2004
<i>Monomorium cyaneum</i> Wheeler W.M., 1914	me	Fernández, 2004
<i>Monomorium destructor</i> (Jerdon, 1851)	ec br tt co am	Fernández, 2004
<i>Monomorium ebeninum</i> Forel, 1891	cr ho gu me am an	Fernández, 2004
<i>Monomorium ergatogyna</i> Wheeler W.M., 1904	us	Fernández, 2004
<i>Monomorium floricola</i> (Jerdon, 1851)	ch ec br gi gf su tt co cr es me us am an bh	Fernández, 2004
<i>Monomorium latastei</i> Emery, 1895	ch	Fernández, 2004

Taxon / Taxón	Neotropical Distribution / Distribución Neotropical	Reference / Referencia
<i>Monomorium minimum</i> (Buckley, 1867)	pr me us	Fernández, 2004
<i>Monomorium monomorium</i> Bolton, 1987	br us	Fernández, 2004
<i>Monomorium pharaonis</i> (Linnaeus, 1758)	ar ch pr ur bo ec br gi gf su tt co pn cr me am bh	Fernández, 2004
<i>Monomorium salomonis</i> (Linnaeus, 1758)	br	Fernández, 2004
<i>Monomorium subcoecum</i> Emery, 1894	an	Fernández, 2004
<i>Monomorium subopacum</i> Smith F., 1858	br me	Fernández, 2004
<i>Oxyepoecus browni</i> Albuquerque & Brandão, 2004	br	Albuquerque & Brandão, 2004
<i>Oxyepoecus bruchi</i> Santschi, 1926	ar	Kempf 1974b
<i>Oxyepoecus daguerrei</i> (Santschi, 1933)	ar	Kempf 1974b
<i>Oxyepoecus ephippiatus</i> Albuquerque & Brandão, 2004	br	Albuquerque & Brandão, 2004
<i>Oxyepoecus inquilinus</i> (Kusnezov, 1952)	ar br	Kempf 1974b
<i>Oxyepoecus kempfi</i> Albuquerque & Brandão, 2004	br	Albuquerque & Brandão, 2004
<i>Oxyepoecus longicephalus</i> Albuquerque & Brandão, 2004	br	Albuquerque & Brandão, 2004
<i>Oxyepoecus mandibularis</i> (Emery, 1913)	bo	Kempf 1974b
<i>Oxyepoecus punctifrons</i> (Borgmeier, 1928)	br	Kempf 1974b
<i>Oxyepoecus quadratus</i> Albuquerque & Brandão, 2004	ec	Albuquerque & Brandão, 2004
<i>Oxyepoecus rastratus</i> (Mayr, 1887)	br	Kempf 1974b
<i>Oxyepoecus vezeyi</i> (Forel, 1907)	pr br	Kempf 1974b
<i>Solenopsis albidula</i> Emery, 1905	ar	Bolton 1995
<i>Solenopsis albidula flava</i> Santschi, 1929	ar	Bolton 1995
<i>Solenopsis albidula postbrunnea</i> Forel, 1913	br	Bolton 1995
<i>Solenopsis altinodis</i> Forel, 1912	gi gf su tt vn	Bolton 1995
<i>Solenopsis alvarengai</i> (Kempf, 1975)	br	Bolton 1995
<i>Solenopsis amblychila</i> Wheeler W.M., 1915	me	Trager 1991; Bolton 1995
<i>Solenopsis amica</i> Menozzi, 1936	br	Bolton 1995
<i>Solenopsis andina</i> Santschi, 1923	ar	Bolton 1995
<i>Solenopsis angulata</i> Emery, 1894	ar ur br	Bolton 1995
<i>Solenopsis angulata carettei ardua</i> Santschi, 1929	ar	Bolton 1995
<i>Solenopsis angulata carettei</i> Forel, 1913	br	Bolton 1995
<i>Solenopsis angulata dolichops</i> Emery, 1905	ar	Bolton 1995
<i>Solenopsis angulata huasanensis</i> Forel, 1912	ar	Bolton 1995
<i>Solenopsis angulata mendozensis</i> Forel, 1914	ar	Bolton 1995
<i>Solenopsis angulata nigelloides</i> Forel, 1913	ar	Bolton 1995
<i>Solenopsis aurea</i> Wheeler W.M., 1906	me	Trager 1991; Bolton 1995
<i>Solenopsis azteca</i> Forel, 1893	an	Bolton 1995
<i>Solenopsis azteca pallida</i> Wheeler W.M., 1908	am	Bolton 1995
<i>Solenopsis basalis</i> Forel, 1896	ar br	Bolton 1995
<i>Solenopsis basalis major</i> Forel, 1913	ar	Bolton 1995
<i>Solenopsis basalis oculatior</i> Forel, 1913	ar	Bolton 1995
<i>Solenopsis basalis raptor</i> Santschi, 1919	ar	Bolton 1995
<i>Solenopsis basalis urichi</i> Forel, 1912	tt an	Bolton 1995
<i>Solenopsis basalis vittata</i> Forel, 1912	br	Bolton 1995
<i>Solenopsis bicolor</i> (Emery, 1925)	ar pr br gi gf su cr	Bolton 1995
<i>Solenopsis brasiliiana</i> Santschi, 1925	br	Bolton 1995
<i>Solenopsis brevicornis</i> Emery, 1888	br gi gf su	Bolton 1995
<i>Solenopsis brevicornis medioclara</i> Santschi, 1923	br	Bolton 1995
<i>Solenopsis brevicornis petropolitana</i> Borgmeier, 1928	br	Bolton 1995
<i>Solenopsis brevipes</i> Emery, 1906	ar br	Bolton 1995
<i>Solenopsis bruchiella</i> Emery, 1922	ar	Bolton 1995
<i>Solenopsis brucki</i> Kempf, 1973	ar	Bolton 1995
<i>Solenopsis bruesi</i> Creighton, 1930	me	Trager 1991; Bolton 1995
<i>Solenopsis castor</i> Forel, 1893	bo cr an	Bolton 1995
<i>Solenopsis clytemnestra</i> Emery, 1896	ar pr ur br	Bolton 1995
<i>Solenopsis clytemnestra bruchi</i> Forel, 1912	ar	Bolton 1995
<i>Solenopsis clytemnestra leda</i> Forel, 1913	br	Bolton 1995
<i>Solenopsis clytemnestra orestes</i> Forel, 1903	br	Bolton 1995
<i>Solenopsis clytemnestra orestes tucumana</i> Forel, 1914	br	Bolton 1995
<i>Solenopsis clytemnestra strangulata</i> Forel, 1913	br	Bolton 1995
<i>Solenopsis condei</i> (Borgmeier, 1937)	br	Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Solenopsis conjurata</i> Wheeler W.M., 1925	pn	Bolton 1995
<i>Solenopsis corticalis</i> Forel, 1881	br co pn cr us am an	Bolton 1995
<i>Solenopsis corticalis amazonensis</i> Forel, 1904	pe gi gf su	Bolton 1995
<i>Solenopsis corticalis binotata</i> Mann, 1920	am	Bolton 1995
<i>Solenopsis corticalis margotae</i> Forel, 1908	br gu	Bolton 1995
<i>Solenopsis corticalis virgula</i> Forel, 1904	am	Bolton 1995
<i>Solenopsis daguerrei</i> (Santschi, 1930)	ar ur	Bolton 1995
<i>Solenopsis decipiens</i> Emery, 1905	ar	Bolton 1995
<i>Solenopsis decipiens abjecta abjectior</i> Forel, 1909	ar	Bolton 1995
<i>Solenopsis decipiens abjecta</i> Emery, 1906	ar	Bolton 1995
<i>Solenopsis decipiens abjecta ignobilis</i> Forel, 1913	br	Bolton 1995
<i>Solenopsis decipiens scelestae</i> Forel, 1908	pr br	Bolton 1995
<i>Solenopsis dysderces</i> Snelling, 1975	ch	Bolton 1995
<i>Solenopsis egregia</i> (Kusnezov, 1953)	ar	Bolton 1995
<i>Solenopsis electra</i> Forel, 1914	bo pa ar	Trager 1991; Bolton 1995
<i>Solenopsis eximia</i> (Kusnezov, 1953)	ar	Bolton 1995
<i>Solenopsis franki</i> Forel, 1908	pr br	Bolton 1995
<i>Solenopsis franki idae</i> Forel, 1908	br	Bolton 1995
<i>Solenopsis gallardoi</i> Santschi, 1925	ar	Bolton 1995
<i>Solenopsis gayi</i> (Spinola, 1851)	ch pe	Trager 1991; Bolton 1995
<i>Solenopsis gayi bruesi</i> Creighton, 1930	pe	Bolton 1995
<i>Solenopsis geminata</i> (Fabricius, 1804)	bo br gi gf su tt vn co pn cr ni be ho gu me us am an	Trager 1991; Bolton 1995
<i>Solenopsis germaini</i> Emery, 1895	ch	Bolton 1995
<i>Solenopsis germaini schedingi</i> Forel, 1907	ch	Bolton 1995
<i>Solenopsis globularia</i> (Smith F., 1858)	br gi co cr	Bolton 1995
<i>Solenopsis globularia lucayensis</i> Wheeler W.M., 1908	bh	Bolton 1995
<i>Solenopsis globularia pacifica</i> Wheeler, 1919	ec	Bolton 1995
<i>Solenopsis globularia pacifica desecheoensis</i> Mann, 1920	am	Bolton 1995
<i>Solenopsis globularia pacifica rubida</i> Wheeler, 1919	ec	Bolton 1995
<i>Solenopsis globularia steinheili</i> Forel, 1881	am an	Bolton 1995
<i>Solenopsis goeldii</i> Forel, 1912	br	Bolton 1995
<i>Solenopsis granivora</i> Kusnezov 1957	ar	Bolton 1995
<i>Solenopsis hammari</i> Mayr, 1903	br	Bolton 1995
<i>Solenopsis hammari carhuensis</i> Forel, 1912	ar	Bolton 1995
<i>Solenopsis hayemi</i> Forel, 1908	cr	Bolton 1995
<i>Solenopsis helena</i> Emery, 1895	ch br	Bolton 1995
<i>Solenopsis helena hermione</i> Wheeler W.M., 1921	gi	Bolton 1995
<i>Solenopsis helena ultrix</i> Wheeler W.M., 1921	gi	Bolton 1995
<i>Solenopsis hostilis</i> (Borgmeier, 1959)	br	Bolton 1995
<i>Solenopsis interrupta</i> Santschi, 1916	ar bo	Trager 1991; Bolton 1995
<i>Solenopsis invicta</i> Buren, 1972	ar br us am	Trager 1991; Bolton 1995
<i>Solenopsis iheringi</i> Forel, 1908	br	Bolton 1995
<i>Solenopsis joergensi</i> Santschi, 1919	ar	Bolton 1995
<i>Solenopsis joergensi cuspisior</i> Santschi, 1923	ar	Bolton 1995
<i>Solenopsis joergensi edentula</i> Santschi, 1933	ar	Bolton 1995
<i>Solenopsis laeviceps</i> Mayr, 1870	br co	Bolton 1995
<i>Solenopsis laeviceps antoniensis</i> Forel, 1912	co	Bolton 1995
<i>Solenopsis latastei</i> Emery, 1895	ch	Bolton 1995
<i>Solenopsis latastei masora</i> Forel, 1912	ch	Bolton 1995
<i>Solenopsis latastei weiseri</i> Santschi, 1933	ar	Bolton 1995
<i>Solenopsis leptanilloides</i> Santschi, 1925	ar	Bolton 1995
<i>Solenopsis loretana</i> Santschi, 1936	ar	Bolton 1995
<i>Solenopsis maboya</i> Snelling 2001		Snelling 2001
<i>Solenopsis macdonaghi</i> Santschi, 1916	ar	Trager 1991; Bolton 1995
<i>Solenopsis macrops</i> Santschi, 1917	ar	Bolton 1995
<i>Solenopsis megergates</i> Trager, 1991	br	Trager 1991; Bolton 1995
<i>Solenopsis metanotalis argus</i> Santschi, 1923	ar	Bolton 1995
<i>Solenopsis metanotalis</i> Emery, 1896	ar	Bolton 1995
<i>Solenopsis metanotalis emiliae</i> Santschi, 1912	ar ur	Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Solenopsis metanotalis pelotana</i> Forel, 1912	ar br	Bolton 1995
<i>Solenopsis metanotalis picturata</i> Santschi, 1931	ar	Bolton 1995
<i>Solenopsis metanotalis shiptoni</i> Forel, 1914	ar	Bolton 1995
<i>Solenopsis metanotalis shiptoni steigeri</i> Santschi, 1916	ar	Bolton 1995
<i>Solenopsis metatarsalis</i> (Kusnezov, 1957)	ar	Bolton 1995
<i>Solenopsis minutissima</i> Emery, 1906	ar gi gf su tt me	Bolton 1995
<i>Solenopsis molesta</i> (Say, 1836)	us	Bolton 1995
<i>Solenopsis nigella</i> Emery, 1888	ar ur br	Bolton 1995
<i>Solenopsis nigella gensterblumi</i> Forel, 1901	ar br	Bolton 1995
<i>Solenopsis novemmaculata</i> Wheeler W.M., 1925	bo	Bolton 1995
<i>Solenopsis oculata</i> Santschi, 1925	ar	Bolton 1995
<i>Solenopsis parabiotica</i> Weber, 1943	pe br gi	Bolton 1995
<i>Solenopsis parva</i> Mayr, 1868	ar	Bolton 1995
<i>Solenopsis patagonica</i> Emery, 1906	ar ch	Bolton 1995
<i>Solenopsis patagonica medeis</i> Forel, 1912	ar br	Bolton 1995
<i>Solenopsis photophila</i> Santschi, 1923	ar	Bolton 1995
<i>Solenopsis picea</i> Emery, 1896	cr ho gu me am	Bolton 1995
<i>Solenopsis picea reducta</i> Menozzi, 1927	cr	Bolton 1995
<i>Solenopsis picea subadpressa</i> Forel, 1903	bo br	Bolton 1995
<i>Solenopsis picquarti</i> Forel, 1899	cr	Bolton 1995
<i>Solenopsis pollux</i> Forel, 1893	br am an	Bolton 1995
<i>Solenopsis pusillignis</i> Trager, 1991	br	Trager 1991; Bolton 1995
<i>Solenopsis pygmaea</i> Forel, 1901	an	Bolton 1995
<i>Solenopsis pythia</i> Santschi, 1934	ar	Trager 1991; Bolton 1995
<i>Solenopsis quinquecuspis</i> Forel, 1913	ar	Trager 1991; Bolton 1995
<i>Solenopsis reichenspergeri</i> Santschi, 1923	br	Bolton 1995
<i>Solenopsis richteri</i> Forel, 1909	ar ch ur	Trager 1991; Bolton 1995
<i>Solenopsis rugiceps</i> Mayr, 1870	co	Bolton 1995
<i>Solenopsis saevissima</i> (Smith F., 1855)	ar pr ur bo ec br gi gf su	Trager 1991; Bolton 1995
<i>Solenopsis schilleri</i> Santschi, 1923	ar	Bolton 1995
<i>Solenopsis schmalzi</i> Forel, 1901	br	Bolton 1995
<i>Solenopsis schmalzi flaveolens</i> Forel, 1901	br	Bolton 1995
<i>Solenopsis sea</i> (Kusnezov, 1953)	ar	Bolton 1995
<i>Solenopsis silvestrii</i> Emery, 1906	ur	Bolton 1995
<i>Solenopsis solenopsisidis</i> (Kusnezov, 1954)	ar	Bolton 1995
<i>Solenopsis spei</i> Forel, 1912	co	Bolton 1995
<i>Solenopsis stricta</i> Emery, 1896	bo	Bolton 1995
<i>Solenopsis stricta foederata</i> Santschi, 1923	br	Bolton 1995
<i>Solenopsis stricta foederata specularis</i> Santschi, 1923	br	Bolton 1995
<i>Solenopsis substituta</i> Santschi, 1925	br	Trager 1991; Bolton 1995
<i>Solenopsis subtilis</i> Emery, 1896	pr br vn	Bolton 1995
<i>Solenopsis succinea</i> Emery, 1890	cr am an	Bolton 1995
<i>Solenopsis succinea nicai</i> Forel, 1913	ar br	Bolton 1995
<i>Solenopsis sulfurea</i> (Roger, 1862)	gi gf su vn	Bolton 1995
<i>Solenopsis tenuis delfinoi</i> Forel, 1913	ar	Bolton 1995
<i>Solenopsis tenuis</i> Mayr, 1878	ar br tt gu	Bolton 1995
<i>Solenopsis tenuis minuiscens</i> Forel, 1912	br	Bolton 1995
<i>Solenopsis terricola</i> Menozzi, 1931	cr	Bolton 1995
<i>Solenopsis tetricantha</i> Emery, 1906	ar	Bolton 1995
<i>Solenopsis tetricantha videns</i> Forel, 1912	ar	Bolton 1995
<i>Solenopsis torresi</i> Snelling, 2001		Snelling 2001
<i>Solenopsis tridens</i> Forel, 1911	br	Trager 1991; Bolton 1995
<i>Solenopsis tridens lehmannnitschei</i> Santschi, 1916	ar	Bolton 1995
<i>Solenopsis trihasta</i> Santschi, 1923	ar	Bolton 1995
<i>Solenopsis virulens</i> Smith F., 1858	co - br	Trager 1991; Bolton 1995
<i>Solenopsis wasmannii</i> Emery, 1894	ar pr bo	Bolton 1995
<i>Solenopsis wasmanni transformis</i> Forel, 1911	ar pr br	Bolton 1995
<i>Solenopsis wasmanni transformis innocens</i> Forel, 1915	ar	Bolton 1995
<i>Solenopsis wasmanni transformis robustior</i> Santschi, 1923	ar	Bolton 1995
<i>Solenopsis weiseri</i> Forel, 1914	ar	Bolton 1995
<i>Solenopsis westwoodi</i> Forel, 1894	br	Bolton 1995

TAXON / Taxón	NEOTROPICAL DISTRIBUTION / Distribución Neotropical	REFERENCE / Referencia
<i>Solenopsis westwoodi atticola</i> Forel, 1912	ar	Bolton 1995
<i>Solenopsis westwoodi platensis</i> Emery, 1906	ar	Bolton 1995
<i>Solenopsis weyreuchi</i> Trager, 1991	pe	Trager 1991; Bolton 1995
<i>Solenopsis xyloni</i> MacCook, 1879	me	Trager 1991; Bolton 1995
<i>Solenopsis zeteki</i> Wheeler W.M., 1942	pn	Bolton 1995
<i>Tranopelta gilva</i> Mayr, 1866	bo br gi gf su tt vn co pn cr	Fernández, 2003a
<i>Tranopelta subterranea</i> (Mann, 1916)	bo ec br	Fernández, 2003a
Insertae sedis		
<i>Ilemomyrmex caecus*</i> Wilson, 1985	ad	Bolton 1995
<i>Oxyidris antillana*</i> Wilson, 1985	ad	Bolton 1995
Tribe Stegomyrmecini		
<i>Stegomyrmex connectens</i> Emery, 1912	bo pe	Diniz 1990
<i>Stegomyrmex manni</i> Smith M.R., 1946	pn	Diniz 1990
Tribe Stenammini		
<i>Bariamyrmex hispidula</i> Lattke, 1990	vn	Lattke 1990
<i>Lachnomyrmex haskinsi</i> Smith M.R., 1944	pn	Fernández & Baena 1997
<i>Lachnomyrmex pilosus</i> Weber, 1950	tt	Fernández & Baena 1997
<i>Lachnomyrmex plaumannii</i> Borgmeier, 1957	br	Fernández & Baena 1997
<i>Lachnomyrmex scrobiculatus</i> Wheeler W.M., 1910	gu me	Fernández & Baena 1997
<i>Rogeria belti</i> Mann, 1922	ho	Kugler 1994
<i>Rogeria blanda</i> (Smith F., 1858)	br gi gf su	Kugler 1994
<i>Rogeria bruchi</i> Santschi, 1922	ar	Kugler 1994
<i>Rogeria brunnea</i> Santschi, 1930	am	Kugler 1994
<i>Rogeria caraiba</i> Santschi, 1936	am	Kugler 1994
<i>Rogeria cubensis</i> Santschi, 1936	am	Kugler 1994
<i>Rogeria cubensis habanica</i> Santschi, 1936	am	Kugler 1994
<i>Rogeria curvipubens</i> Emery, 1894	ar am an	Kugler 1994
<i>Rogeria foreli</i> Emery, 1894	ar an	Kugler 1994
<i>Rogeria foreli gaigei</i> Forel, 1914	co	Kugler 1994
<i>Rogeria germaini</i> Emery, 1894	br	Kugler 1994
<i>Rogeria inermis</i> Mann, 1922	ho	Kugler 1994
<i>Rogeria lacertosa</i> Kempf, 1963	br	Kugler 1994
<i>Rogeria micromma</i> Kempf, 1961	gi gf su	Kugler 1994
<i>Rogeria minensis</i> Santschi, 1923	br	Kugler 1994
<i>Rogeria minima</i> Kusnezov, 1958	ar	Kugler 1994
<i>Rogeria pellecata</i> Kempf, 1963	br	Kugler 1994
<i>Rogeria procera</i> Emery, 1896	br	Kugler 1994
<i>Rogeria scabra</i> Weber, 1934	am	Kugler 1994
<i>Rogeria scandens</i> (Mann, 1922)	ho	Kugler 1994
<i>Rogeria sicaria</i> Kempf, 1962	br	Kugler 1994
<i>Rogeria subarmata</i> (Kempf, 1961)	br	Kugler 1994
<i>Rogeria tonduzi</i> Forel, 1899	cr	Kugler 1994
<i>Stenamma californicum</i> Snelling, 1973	us	Smith 1962
<i>Stenamma diecki</i> Emery, 1895	us	Smith 1962
<i>Stenamma diversum</i> Mann, 1922	ho	Smith 1962
<i>Stenamma expolitum</i> Smith M.R., 1962	cr	Smith 1962
<i>Stenamma felixi</i> Mann, 1922	co cr ho me	Smith 1962
<i>Stenamma heathi</i> Wheeler W.M., 1915	us	Smith 1962
<i>Stenamma manni</i> Wheeler W.M., 1914	me	Smith 1962
<i>Stenamma schmidti</i> Menozzi, 1931	cr	Smith 1962
<i>Stenamma snellingi</i> Bolton, 1995	us	Smith 1962
Tribe Tetramoriini		
<i>Tetramorium caespitum</i> (Linnaeus, 1758)	ar ch	Bolton 1979
<i>Tetramorium guineense</i> (Bernard, 1953)	pe ec br gi gf su tt co pn cr me am an	Bolton 1979
<i>Tetramorium lucayanum</i> Wheeler W.M., 1905	am bh	Bolton 1979
<i>Tetramorium spinosum</i> (Pergande, 1896)	bo ec br gi gf su co ho me am an bh	Bolton 1979
Insertae sedis en Formicidae		
Subfamily Paleosminthurinae		
<i>Cariridris bipetiolata*</i> Brandão & Martins-Neto, 1990	br	Bolton 2003

Acknowledgments / Agradecimientos

This list is the result of many friends and colleagues who of one or another form have contributed with Literature, names and copies of last hour. We are in debt with all of them: Barry Bolton (Museum of Natural History of London), Carlos Brandão Robert (University of São Paulo), William L. Brown Jr, deceased (University of Cornell), Fabiana Cuezzo (University of Tucumán), Víctor Hugo Gonzalez (University of Kansas), John S. LaPolla (University of Rutgers), John And Lattke (Central University of Venezuela), John T. Longino (Evergreen College), William P. MacKay (University of Texas in the Step), Fernando Muñoz (University of Antioquia), Patricia Red-Fernandez (Institute of Ecology, Xalapa), Ted Schultz (National Museum of Natural History in Washington), Michael J. Sharkey (University of Kentucky), Roy R. Snelling (Museum of Natural History of Los Angeles), Jeffrey Sossa (University of Maryland), Philip S. Ward (University of California), and Edward Or Wilson (University of Harvard).

Esta lista es fruto de muchos amigos y colegas que de una u otra forma han contribuido con literatura, nombres y copias de última hora. Estamos en deuda con todos ellos: Barry Bolton (Museo de Historia Natural de Londres), Carlos Roberto Brandão (Universidad de São Paulo), William L. Brown Jr., fallecido (Universidad de Cornell), Fabiana Cuezzo (Universidad de Tucumán), Victor Hugo Gonzalez (Universidad de Kansas), John S. LaPolla (Universidad de Rutgers), John E. Lattke (Universidad Central de Venezuela), John T. Longino (Evergreen College), William P. MacKay (Universidad de Texas en El Paso), Fernando Muñoz (Universidad de Antioquia), Patricia Rojas-Fernández (Instituto de Ecología, Xalapa), Ted Schultz (Museo Nacional de Historia Natural en Washington), Michael J. Sharkey (Universidad de Kentucky), Roy R. Snelling (Museo de Historia Natural de Los Angeles), Jeffrey Sossa (Universidad de Maryland), Philip S. Ward (Universidad de California), y Edward O. Wilson (Universidad de Harvard).

Literature Cited / Literatura Citada

- Agosti D. (1991) Revision of the oriental ant genus *Cladomyrma*, with an outline of the higher classification of the Formicidae (Hymenoptera: Formicidae). *Systematic Entomology* 16:293-310.
- Agosti D. (1994) The phylogeny of the ant tribe Formicinae (Hymenoptera: Formicidae) with the description of a new genus. *Systematic Entomology* 19:93-117.
- Agosti D. (1995) A revision of the South American species of the ant genus *Probolomyrmex*. *Journal of the New York Entomological Society* 102:429-434.
- Agosti D. (2004) www.AntWeb.Org
- Agosti, D., J. D. Majer, L. E. Alonso y T. R. Schultz (eds) (2000) *Ants: Standard Methods for Measuring and Monitoring Biodiversity*. Smithsonian Institution Press.
- Agosti, D., N. Johnson (2003) La nueva taxonomía de hormigas, pp. 45-48 en: F. Fernández, ed., *Introducción a las hormigas de la región Neotropical*. Instituto Humboldt, Bogotá D.C.
- Albuquerque N.L., C.R.F. Brandão. (2004) A revision of the Neotropical Solenopsidini ant genus *Oxyepoecus* Santschi, 1926 (Hymenoptera: Formicidae: Myrmicinae). 1. The vezényii species group *Papéis Avulsos de Zoologia* 44(4):55-80.
- Andrade M. L. de, C. Baroni Urbani (1999). Diversity and Adaptation in the Ant Genus *Cephalotes*, Past and Present. *Stuttgarter Beiträge zur Naturkunde Serie B (Geologie und Paläontologie)* 271:1-889.
- Andrade M. L. de. (1998) Fossil and extant species of *Cylindromyrmex* (Hymenoptera: Formicidae). *Revue Suisse de Zoologie* 105(3):581-664.
- Andrade M. L. de. (2001) A remarkable Dominican amber species of *Cylindromyrmex* with Brazilian affinities and additions to the generic revision (Hymenoptera: Formicidae). *Beiträge zur Entomologie* 51(1):51-63.
- Astruc, C., J.F. Julien, C. Errard, A. Lenoir (2004) Phylogeny of ants (Formicidae) based on morphology and DNA sequence data. *Molecular Phylogenetics and Evolution* 31:880-893.
- Ayala F. J., J. K. Wetterer, J. T. Longino, D. L. Hartl. (1996) Molecular phylogeny of *Azteca* ants (Hymenoptera: Formicidae) and the colonization of *Cecropia* trees. *Molecular Phylogenetic Evolution* 5:423-428.
- Baroni Urbani C. (1975) Contributo alla conoscenza dei generi *Belonopelta* Mayr e *Leiopelta* gen. n. (Hymenoptera: Formicidae). *Mitteilungen der Schwieizerischen Entomologischen Gesellschaft* 48:295-310.
- Baroni Urbani C. (1978) Materiali per una revisione dei *Leptothorax* neotropicali appartenenti al sottogenere *Macromischa* Roger, n. comb. (Hymenoptera: Formicidae). *Entomologica Basiliensis* 3:395-618.
- Baroni Urbani C. (2000) Rediscovery of the Baltic amber ant genus *Prionomyrmex* (Hymenoptera, Formicidae) and its taxonomic consequences. *Eclogae geol. Helv.* 93:471-480.

- Baroni Urbani C., M. L. de Andrade (1994) First description of fossil Dacetini ants with a critical analysis of the current classification of the tribe (Amber Collection Stuttgart: Hymenoptera, Formicidae. VI. Dacetini). *Stuttgarter Beiträge zur Naturkunde Serie B (Geologie und Paläontologie)* 198:1-65.
- Baroni Urbani C., M.L. de Andrade (2003) The ant genus *Proceratium* in the extant and fossil record. *Museo Regionale di Scienze Naturali Monografie* 36:1-492.
- Baroni Urbani C., B. Bolton, P. S. Ward (1992) The internal phylogeny of ants (Hymenoptera: Formicidae). *Systematic Biology* 17:301-329.
- Bolton B. (1976) The ant tribe Tetramoriini. Constituent genera, review of the smaller genera and revision of *Triglyphothrix* Forel. *Bulletin of the British Museum of Natural History (Entomol.)* 34:281-379.
- Bolton B. (1979) The ant tribe Tetramoriini. The genus *Tetramorium* Mayr in the Malagasy region and in the New World *Bulletin of the British Museum of Natural History (Entomology)* 38:129-181.
- Bolton B. (1987) A review of the *Solenopsis* genus group and revision of Afrotropical *Monomorium* Mayr *Bulletin of the British Museum of Natural History (Entomol.)* 54:263-452.
- Bolton B. (1990a) Abdominal characters and status of the cerapachyine ants (Hymenoptera, Formicidae) *Journal of Natural History* 24:53-68.
- Bolton, B. (1990b). Army ants reassessed: the phylogeny and classification of the doryline section (Hymenoptera, Formicidae) *Journal of Natural History* 24:1339-1364.
- Bolton B. (1994) *Identification guide to the ant genera of the world* Harvard University Press, Cambridge, Massachusetts, E.E.U.U., 222 pp
- Bolton B. (1995a) *A New General Catalogue of the Ants of the World* Harvard University Press, 504 pp.
- Bolton B. (1995b) A taxonomic and zoogeographical census of extant ant taxa (Hymenoptera: Formicidae) *Journal of Natural History* 29:1037-1056.
- Bolton B. (1998) Monophyly of the dacetonini tribe-group and its component tribes *Bulletin of the British Museum of Natural History (Entomology)* 67:65-78.
- Bolton B. (1999) Ant genera of the tribe Dacetonini (Hymenoptera: Formicidae) *Journal of Natural History* 23:1267-1308.
- Bolton B. (2000) The ant tribe Dacetini. *Memoirs of the American Entomological Institute* 65(parts I+II):1-1065.
- Bolton B. (2003) Synopsis and classification of Formicidae. *Memoirs of the American Entomological Institute* 71:1-370.
- Bolton B., W.L. Brown Jr. (2002) *Loboponera* gen.n. and a review of the Afrotropical *Plectroctena* genus group. *Bulletin of the Natural History Museum* 71:1-18.
- Borgmeier T. (1949) Formigas novas ou pouco conhecidas de Costa Rica e da Argentina *Revista Brasileira de Biologia* 9:201-210.
- Borgmeier T. (1955) Die Wanderameisen der Neotropischen Region *Studia Entomologica* 3:1-720.
- Borgmeier T. (1959) Revision der Gattung *Atta* Fabricius *Studia Entomologica* (N.S.) 2:321-390.
- Bourke, A.F.G, N.R. Franks (1995) *Social Evolution in Ants*. Princeton University Press, NY.
- Brady S.G. (2003) Evolution of the army ant syndrome: The origin and long-term evolutionary stasis of a complex of behavioral and reproductive adaptations *Proceedings of the National Academy of Sciences* 100(11):6575-6579.
- Brady S.G, J. Gadau, P. S. Ward (2000) Systematics of the ant genus *Camponotus* (Hymenoptera: Formicidae): a preliminary analysis using data from the mitochondrial gene cytochrome Oxidase I, pp. 131-139 En A. D. Austin, M. Dowton, eds., *Hymenoptera: Evolution, Biodiversity and Biological Control*. CSIRO Pub., Australia.
- Brandão C. R. F. (1990) Systematic revision of the Neotropical ant genus *Megalomyrmex* Forel (Hymenoptera: Formicidae) with the description of thirteen new species. *Arquivos de Zoologia* 31:411-481.
- Brandão C.R.F. (1991) Adendos ao catálogo abreviado das formigas da região neotropical (Hymenoptera: Formicidae). *Revista Brasileira de Entomologia* 35:319-412.
- Brandão C.R.F. (2003) Further revisionary studies on the ant genus *Megalomyrmex* Forel (Hymenoptera: Formicidae: Myrmicinae: Solenopsidini) *Papéis Avulsos de Zoologia* 43(8):145-159.
- Brandão C. R. F., Baroni Urbani, J., Wagensberg, C. I. Yamamoto (1998) New *Technomyrmex* in Dominican amber (Hymenoptera: Formicidae), with a reappraisal of Dolichoderinae phylogeny. *Entomologica Scandinavica* 29(4):1-10.
- Brandão C. R. F., A. Mayhe-Nunes (2001) A new fungus growing ants genus *Mycetagoicus* gen.n. with the description of three new species and comments on the monophyly of Attini (Hymenoptera: Formicidae) *Sociobiology* 38(3B):639-665.
- Brandão, C. R. F., J. L. M. Diniz, D. Agosti y J. H. Delabie. 1999. Revision of the Neotropical ant subfamily Leptanilloidinae. *Systematic Entomology* 24:17-36.
- Brown W. L. Jr. (1952) Contributions toward a reclassification of the Formicidae. I. Tribe Platythyreini (Hymenoptera) *Breviora* 6:1-6.
- Brown W. L. Jr. (1958) Contributions toward a reclassification of the Formicidae. II. Tribe Ectatommini (Hymenoptera) *Bulletin of the Museum of Comparative Zoology* 118(5):175-362.

- Brown W. L. Jr. (1960) Contributions toward a reclassification of the Formicidae. III. Tribe Amblyoponini (Hymenoptera) *Bulletin of the Museum of Comparative Zoology* 122:145-230.
- Brown W. L. Jr. (1965) Contributions toward a reclassification of the Formicidae. IV. Tribe Typhlomyrmecini (Hymenoptera) *Psyche* 72:65-78.
- Brown W. L. Jr. (1973) A comparison of the Hylean and Congo-West African rain forest ant faunas, pp 165-185 en E. J. Meggers, A. S Ayensu y W. D. Duckworth, eds., *Tropical Forest Ecosystems in Africa and South America: A Comparative Review*. Smithsonian Institution Press, Washington, D. C.
- Brown W. L. Jr. (1974) A supplement to the revision of the ant genus *Basiceros* (Hymenoptera: Formicidae) *Journal of the New York Entomological Society* 82:131-140.
- Brown W. L. Jr. (1975) Contributions toward a reclassification of the Formicidae. V. Ponerinae, tribes Platythyreini, Cerapachyini, Cylindromyrmecini, Acanthostichini, and Aenictogitini *Search Agriculture (Ithaca, New York)* 5:1-115.
- Brown W. L. Jr. (1976) Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, Tribe Ponerini, Subtribe Odontomachiti. Section A. Introduction, Subtribal characters. Genus *Odontomachus* *Studia Entomologica* 19:67-171.
- Brown W. L. Jr. (1978) Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, Tribe Ponerini, Subtribe Odontomachiti. Section A. Introduction. Genus *Anochetus* *Studia Entomologica* 20:549-652.
- Brown, W. L. Jr. (1979) A remarkable new species of *Proceratium* with dietary and other notes on the genus (Hymenoptera: Formicidae) *Psyche* 86:337-346.
- Brown W.L. (1980) *Protalaridris* genus nov. *Pilot Register of Zoology* Card No. 37.
- Brown W. L. Jr. (1981) Preliminary contributions toward a revision of the ant genus *Pheidole* (Hymenoptera: Formicidae). Part I *Journal of the Kansas Entomological Society* 54:523-530.
- Brown W. L. Jr. (1992) Two new species of *Gnamptogenys*, and an account of millipede predation by one of them *Psyche* 99:275-289.
- Brown W. L. Jr.(2000) Diversity of ants, pp. 45-79 in D. Agosti, J. D. Majer, L. E. Alonso and T. R. Schultz, eds., *Ants: Standard Methods for Measuring and Monitoring Biodiversity*. Smithsonian Institution Press.
- Brown W. L., W. W. Kempf (1960) A world revision of the ant tribe Basicerotini *Studia Entomologica* (N. S.) 3:161-250.
- Brown W. L., W. W. Kempf. (1967) *Tatuidris*, a remarkable new genus of Formicidae *Psyche* 74:183-190.
- Brown W.L., W.W. Kempf (1969) A revision of the Neotropical Dacetine ant genus *Acanthognathus* *Psyche* 76(2):87-109.
- Chiotis M., L. S. Jermiin, R. H. Crozier (2000) A Molecular Framework for the Phylogeny of the ant Subfamily Dolichoderinae *Molecular Phylogenetic Evolution* 17(1):108-116.
- Creighton W. S. (1930) The New World species of the genus *Solenopsis* Part I. *Procedures of the American Academy of Arts and Sciences* 66:39-151.
- Creighton W. S. (1957) A study of the genus *Xenomyrmex* *American Museum Novitates* 1843:1-14.
- Cuezzo F. (2000) Revisión del género *Forelius* (Hymenoptera: Formicidae: Dolichoderinae) *Sociobiology* 35(2A):197-277.
- Diniz J.L.M. (1990) Revisão sistemática da tribo Stegomyrmecini, com a descrição de uma nova espécie (Hymenoptera: Formicidae) *Revista Brasileira de Entomologia* 34:277-295.
- Du Bois M. (1986) A revision of the native New World species of the ant genus *Monomorium* (*minimum* group) (Hymenoptera: Formicidae) *University of Kansas Science Bulletin* 53:65-119.
- Emery C. (1893) Studio monografico sul genere *Azteca* Forel *Memorie della R. Accademia delle Scienze dell'Istituto di Bologna* (5)3:119-152.
- Ettershank G. (1966) A generic revision of the world Myrmicinae related to *Solenopsis* and *Pheidole* *geton Australian Journal of Zoology* 14:73-171.
- Fernández F. (2001) Hormigas de Colombia. IX: Nueva especie de *Lenomyrmex* (Formicidae: Myrmicinae). *Revista Colombiana de Entomología* 27(3-4):201-204.
- Fernández F. (2002a) Revisión de las hormigas *Camponotus* del subgénero *Dendromyrmex* (Hymenoptera: Formicidae) *Papeis Avulsos de Zoología* (São Paulo) 42(4):47-101.
- Fernández F. (2002b) New ant records for Colombia and South America (Hymenoptera: Formicidae) *Revisita Colombiana de Entomología* 28(2):215.
- Fernández F. (2003a) The myrmicine ant genera *Ochetomyrmex* Mayr and *Tranopelta* Mayr (Hymenoptera: Formicidae) *Sociobiology* 41(3):1-24.
- Fernández F. (2003b) Revision of the myrmicine ants of the *Adelomyrmex* genus-group *Zootaxa* 361:1-52.
- Fernández F. (ed) (2003c) *Introducción a las hormigas de la región Neotropical*. Instituto Humboldt, Bogotá D.C., 424 pp.
- Fernández F. (2004a) The american species of the myrmicine ant genus *Carebara* Westwood (Hymenoptera: Formicidae) *Caldasia* 26(1):191-238.

- Fernández F. (2004b) Adelomyrmecini new tribe and *Cryptomyrmex* new genus of myrmicine ants (Hymenoptera: Formicidae) *Sociobiology* 44(2):325-335.
- Fernández F. (2004c) Two new South American species of *Monomorium* Mayr (Hymenoptera: Formicidae) with taxonomic notes on the genus *Memoirs of the American Entomological Institute* 38.
- Fernández, F., E.E. Palacio (1998) Clave para las *Pogonomyrmex* (Hymenoptera: Formicidae) del Norte de Suramérica, con la descripción de una nueva especie *Revista de Biología Tropical* 45(4):1649-1661.
- Fernández, F., E.E. Palacio (1999) *Lenomyrmex*, an enigmatic new ant genus from the Neotropical Region (Hymenoptera: Formicidae: Myrmicinae) *Systematic Entomology* 24:7-16.
- Fernández F., E.E. Palacio (2003) Sistemática y filogenia de las hormigas: breve repaso a propuestas, pp. 29-49 en: F. Fernández, ed., *Introducción a las hormigas de la región Neotropical*, Instituto Humboldt. Bogotá D.C.
- Fernández F., M.L. Baena (1997) Hormigas de Colombia VII. Nuevas especies de los géneros *Lachnomyrmex* Wheeler y *Megalomyrmex* Forel (Hymenoptera: Formicidae) *Caldasia* 19(1-2):109-114.
- Fernández F., W.P. MacKay (2003) The myrmicinae ants of the *Adelomyrmex laevigatus* species complex (Hymenoptera: Formicidae) *Sociobiology* 41(2):1-12.
- Fontenla J.L. (2000) Historical biogeography and character evolution in the phylogenetic taxon "Macromischa" (Hymenoptera: Formicidae: Leptothorax) *Transactions of the American Entomological Society* 126(3-4):401-416.
- Fowler H.G. (1988) Taxa of the Neotropical grass-cutting ants, *Acromyrmex (Moellerius)* (Hymenoptera: Formicidae: Attini) *Científica* 16(2):281-296.
- Gonçalves C. R. (1961) O genero *Acromyrmex* no Brasil *Studia Entomologica* 4:113-180.
- Gotwald W. L. (1982) Army Ants, pp. 157-254 En H. R. Hermann, ed., *Social Insects*. Vol. 4.
- Gotwald W. L. (1995) *Army Ants: The Biology of Social Predation*. Cornell University Press, Ithaca.
- Gotwald W., W. L. Brown Jr. (1966) The ant genus *Simopelta* (Hymenoptera: Formicidae) *Psyche* 73:261-277.
- Grimaldi D., D. Agosti (2000) A formicine in New Jersey Cretaceous amber (Hymenoptera: Formicidae) and early evolution of the ants *Proceedings of the National Academy of Sciences* 97(25):13678-13683.
- Grimaldi, D., D. Agosti, J. M. Carpenter (1997) New and rediscovered primitive ants (Hymenoptera: Formicidae) in Cretaceous amber from New Jersey, and their phylogenetic relationships *American Museum Novitates* 3208:1-43.
- Hashmi A. A. (1973) A revision of the Neotropical ant subgenus *Myrmothrix* of genus *Camponotus* *Studia Entomologica* (N.S.) 16:1-140.
- Heterick B.E. (2001) Revision of the Australian ants of the genus *Monomorium* (Hymenoptera: Formicidae) *Invertebrate Taxonomy* 15:353-459.
- Hölldobler B., E. O. Wilson (1990) The ants, 732 pp. Belknap Press, Cambridge, Massachusetts.
- Jaffé K. (1993) *El Mundo de las Hormigas*. Equinoccio, Baruta, Venezuela.
- Keller R. (2000) Cladistics of the tribe Ectatommini (Hymenoptera: Formicidae): a reappraisal *Insect Systematics and Evolution* 31:59-69.
- Kempf, W. W. (1951) A taxonomic study on the ant tribe Cephalotini *Revista de Entomología* 22:1-244.
- Kempf W. W. (1958) New studies of the ant tribe Cephalotini *Studia Entomologica* (N.S.) 1:1-168.
- Kempf W. W. (1959) A synopsis of the New World species belonging to the *Nesomyrmex*-group of the ant genus *Leptothorax* Mayr. *Studia Entomologica* (N.S.) 2:391-432.
- Kempf W. W. (1960) *Phalacromyrmex*, a new ant genus from southern Brazil *Revista Brasileira de Biología* 20:89-92.
- Kempf W. W. (1961) A review of the ant genus *Mycetarotes* Emery *Revista Brasileira de Biología* 20:277-283.
- Kempf W. W. (1962) Retoques à classificação das formigas neotropicais do gênero *Heteroponera* Mayr (Hym., Formicidae) *Papeis Avulsos de Zoologia* 15:29-47.
- Kempf W. W. (1963) A review of the ant genus *Mycoceropurus* Forel *Studia Entomologica* 6:417-432.
- Kempf W. W. (1964a) Additions to the knowledge of the Cephalotine ants *Papeis Avulsos de Zoologia* 16:243-255.
- Kempf W. W. (1964b) A revision of the Neotropical fungus-growing ants of the genus *Cyphomyrmex* Mayr. Part I: Group of *strigatus* Mayr *Studia Entomologica* 7:1-44.
- Kempf W. W. (1966) A revision of the Neotropical fungus-growing ants of the genus *Cyphomyrmex* Mayr. Part II: Group of *rimosus* (Spinola). *Studia Entomologica* 8:161-200.
- Kempf W. W. (1966) A synopsis of the Neotropical ants of the genus *Thaumatomyrmex* Mayr (Hymenoptera: Formicidae) *Studia Entomologica* 9:401-410.
- Kempf W. W. (1967) A synopsis of the Neotropical ants of the genus *Centromyrmex* Mayr. *Studia Entomologica* 9:401-410.
- Kempf W. W. (1971) A preliminary review of the ponerine ant genus *Dinoponera* Roger (Hym. Formicidae) *Studia Entomologica* 14:369-394.
- Kempf W. W. (1972) Catálogo abreviado das formigas da Região Neotropical *Studia Entomologica* 15:3-344.

- Kempf W. W. (1973)a A revision of the Neotropical Myrmicine ant genus *Hylomyrma* Forel. *Studia Entomologica* 16:225-260.
- Kempf W. W. (1973)b A new *Zacryptocerus* from Brazil, with remarks on the generic classification of the Tribe Cephalotini (Hymenoptera, Formicidae) *Studia Entomologica* 16:449-462.
- Kempf, W. W. (1974) A review of the Neotropical ant genus *Oxyepoecus* Santschi (Hym., Formicidae). *Studia Entomol.* 17:471-512.
- Kempf W. W. (1975) A revision of the Neotropical ponerine ant genus *Thaumatomyrmex* Mayr (Hymenoptera: Formicidae) *Studia Entomologica* 18:95-126.
- Kempf W. W. (1975) Miscellaneous studies on Neotropical ants VI. (Hymenoptera, Formicidae) *Studia Entomologica* 18:95-126.
- Kempf W. W., W. L. Brown Jr. (1968) Report on some neotropical ant studies *Papéis Avulsos de Zoologia* (São Paulo) 22:89-102.
- Kempf W. W., K. Lenko (1968) Novas observações e estudos sobre *Gigantiops destructor* (Fabricius) (Hymenoptera: Formicidae). *Papeis Avulsos de Zoologia* 21(22):209-230.
- Kugler C. (1977) A new species of *Platythyrea* (Hymenoptera, Formicidae) from Costa Rica *Psyche* 83:216-221.
- Kugler C. (1994) A revision of the ant genus *Rogeria* with descriptions of the sting apparatus (Hymenoptera: Formicidae) *Journal of Hymenoptera Research* 3:17-89.
- Kugler C., W. L. Brown Jr. (1982) Revisionary and other studies on the ant genus *Ectatomma*, including the descriptions of two new species *Search Agriculture* (Ithaca, New York) 24:1-8.
- Kusnezov N. (1952a) *Lasiophanes* Emery en la Patagonia. *Acta Zoologica Lilloana* 12:89-100.
- Kusnezov N. (1952b) El género *Camponotus* en la Argentina. *Acta Zoologica Lilloana* 12:183-252.
- Kusnezov N. (1952c) El estado real del grupo *Dorymyrmex* Mayr (Hymenoptera, Formicidae) *Acta Zoologica Lilloana* 10:427-448.
- Kusnezov N. (1957) Die Solenopsidinen-Gattungen von Suedamerika *Zoologisher Anzeiger* 158:266-280.
- Kusnezov N. (1963) Zoogeografía de las hormigas en Sudamérica *Acta Zoológica Lilloana* 19:25-186.
- Kusnezov N. (1978) Hormigas Argentinas: Claves para su identificación *Instituto Miguel Lillo Miscelánea Tucumán*.
- Lacau S., J. H. C. Delabie (2002) Description de trois nouvelles espèces d'*Amblyopone* avec quelques notes biogéographiques sur le genre au Brésil (Formicidae, Ponerinae) *Bulletin de la Société Entomologique de France* 107(1):33-41.
- LaPolla J. (2004) *Acropyga* (Hymenoptera : Formicidae) of the World *Contributions of the American Entomological Institute* 33(3):1-130.
- Lattke J.E. (1990a) Revisión del género *Gnamptogenys* Mayr para Venezuela *Acta Terramaris* 2:1-47.
- Lattke J.E. (1990b) A new genus of myrmicine ants (Hymenoptera: Formicidae) from Venezuela *Entomologica Scandinavica* 21:173-178.
- Lattke J.E. (1991) Studies of neotropical *Amblyopone* Erichson (Hymenoptera: Formicidae) *Contributions in Science* 428:1-7.
- Lattke J.E. (1994) Phylogenetic relationships and classification of the Ectatommini *Entomologica Scandinavica* 25:105-119.
- Lattke J.E. (1995) Revision of the ant genus *Gnamptogenys* in the New World (Hymenoptera: Formicidae) *Journal of Hymenoptera Research* 4:137-193.
- Lattke J.E. (1997) Revisión del género *Apterostigma* (Hymenoptera: Formicidae) *Arquivos de Zoología* 34(5):121-221.
- Lattke J.E. (1999) A new species of fungus-growing ant and its implications for attine phylogeny (Hymenoptera: Formicidae) *Systematic Entomology* 24:1-6.
- Lattke J.E. (2002a) Nuevas especies de *Gnamptogenys* Roger, 1863 de América (Hymenoptera: Formicidae: Ponerinae) *Entomotropica* 17(2):135-144.
- Lattke J.E. (2004) A taxonomic revision and phylogenetic analysis of the ant genus *Gnamptogenys* Roger in Southeast Asia and Australasia (Hymenoptera: Formicidae) *University of California Publications in Entomology* 122:1-264.
- Lattke J. E., W. Goitia. (1997) El Género *Strumigenys* (Hymenoptera: Formicidae) en Venezuela *Caldasia* 19(3):367-396.
- Longino J. T. (1988) Notes on the taxonomy of the Neotropical ant genus *Thaumatomyrmex* Mayr (Hymenoptera: Formicidae), pp. 35-42 en: J. M. Trager, ed., *Advances in Myrmecology* E. J. Brill, New York.
- Longino J. T. (1989) Taxonomy of the *Cecropia*-inhabiting ants in the *Azteca alfari* species group (Hymenoptera: Formicidae): Evidence for two broadly sympatric species *Contributions in Science* 412:1-16.
- Longino J.T. (1991) Taxonomy of the *Cecropia*-inhabiting *Azteca* ants. *Journal of Natural History* 25:1571-1602.
- Longino J.T. (1996) Taxonomic characterization of some live-stem inhabiting *Azteca* in Costa Rica, with special reference to the ants of *Cordia* (Boraginaceae) and *Triplaris* (Polygonaceae) *Journal of Hymenoptera Research* 5:131-156.
- Longino J. T. (2003) The *Crematogaster* (Hymenoptera, Formicidae, Myrmicinae) of Costa Rica *Zootaxa* 151:1-150.
- Longino J. T., D. A. Hartley (1994) *Perissomyrmex snyderi* (Hymenoptera: Formicidae) is native to Central America and exhibits worker polymorphism *Psyche* 101(3-4):195-202.

- Longino J. T., P. Hanson (1995) The Ants (Formicidae), pp. 587-620 in P. Hanson and I. Gauld, eds., *The Hymenoptera of Costa Rica*. Oxford University Press, New York.
- Longino J. T., R. R. Snelling (2002) A taxonomic revision of the *Procryptocerus* (Hymenoptera: Formicidae) of Central America. *Contributions in Science* 495:1-30.
- MacKay W. P. (1993) A review of the New World ants of the genus *Dolichoderus* (Hymenoptera: Formicidae). *Sociobiology* 22(1):1-148.
- MacKay W. P. (1995) New distributional records for the ants genus *Cardiocondyla* in the New World (Hymenoptera: Formicidae). *Pan-Pacific Entomologist* 71(3):169-172.
- MacKay W. P. (1996) A revision of the ant genus *Acanthostichus* (Hymenoptera: Formicidae). *Sociobiology* 27:129-179.
- MacKay W. P. (1997) A revision of the Neotropical ants of the genus *Camponotus*, subgenus *Myrmostenus* (Hymenoptera: Formicidae). *Proceedings of the Entomological Society of Washington* 99(1):194-203.
- MacKay W. P. (1998) Dos especies nuevas de hormigas de la Tribu Attini de Costa Rica y México: *Mycetosoritis vinnsoni* y *Mycocepurus curvispinosus* (Hymenoptera: Formicidae). *Revista de Biología Tropical* 46(2):421-426.
- MacKay W. P. (2001) A Review of the New World ants of the subgenus *Myrafant*, genus *Leptocephalus* (Hymenoptera: Formicidae). *Sociobiology* 35(2):1-112.
- MacKay W.P. (2003) A new species of the ant genus *Acanthostichus* (Hymenoptera: Formicidae) from Paraguay, and a description of the female of *A. brevicornis*. Proc. Entomol. Soc. Wash. In press.
- MacKay W.P. (2005) Biology and Systematics of the Neotropical carpenter ants of the genus *Camponotus* (Hymenoptera: Formicidae).
- MacKay W. P., E. MacKay (1986) Las hormigas de Colombia: Arrieras del género *Atta* (Hymenoptera: Formicidae). *Revista Colombiana de Entomología* 12(1):23-30.
- MacKay, W. P., E. MacKay (1997a) A revision of the Neotropical ants of the *montivagus* species complex, genus *Camponotus*, subgenus *Myrmecoma* (Hymenoptera: Formicidae). *Sociobiology* 30(3):319-334.
- MacKay W. P. y E. MacKay (1997b) Una nueva especie de hormiga del género *Trachymyrmex* (Hymenoptera: Formicidae) del estado de Chihuahua, México. *Sociobiology* 30(1):43-49.
- Mackay W. P., M. L. Baena (1993) A new "horned" fungus growing ant, *Cyphomyrmex castagnei*, from Colombia. *Sociobiology* 22:31-37.
- Mayhé-Nunes A. (1995) Sinopse do genero *Mycetarotes* (Hym., Formicidae), com a descrição de duas espécies novas. *Boletín de Entomología Venezolana N.S.* 10(2):197-202.
- Moffet M. (1986) Revision of the myrmicine genus *Acanthomyrmex*. *Bulletin of the Museum of Comparative Zoology* 151:55-89.
- Mueller, U.G., S.A. Rehner y T.R. Schultz. 1998. The evolution of agriculture in ants. *Science* 281:2034-2038.
- Mueller U.G., T.R. Schultz C.R. Currie, R.M.M. Adams, D. Malloch (2001) The origin of the attine ant-fungus mutualism. *The Quarterly Review of Biology* 76(2):169-197.
- O'Keefe S., D. Agosti (1997) A new species of *Probolomyrmex* (Hymenoptera: Formicidae) from Guanacaste, Costa Rica. *Journal of the New York Entomological Society* 105:190-192.
- Okido H., K. Ogata (2000) Structures of the Labrum and Antennal Base in the Tribe Myrmecinini (Formicidae: Myrmicinae). *ANet Newsletter* 1:19-21.
- Palacio E.E. (1997) Hormigas de Colombia VI. Dos nuevas especies de *Octostruma* (Hymenoptera: Formicidae: Basicerotini). *Caldasia* 19(3):409-418.
- Palacio E. E. (1999) Hormigas legionarias (Hymenoptera: Formicidae: Ecitoninae) de Colombia, pp. 117-189 en G. Amat, G. Andrade y F. Fernández, eds., *Insectos de Colombia Vol. II*. Academia Colombiana de Ciencias Exactas, Físicas y Naturales y Facultad de Ciencias Universidad Nacional de Colombia, Bogotá D.C.
- Perrault G.H. (1988) *Octostruma betschi*, n.sp. de Guyane Française (Hymenoptera: Formicidae). *Revue Française d'Entomologie* (N.S) 10(4):303-307.
- Perrault G. (2000) Probolomyrmicinae, nouvelle sous-famille pour la genre *Probolomyrmex*. *Bulletin de la Société Entomologique de France* 105:253-272.
- Rojas-Fernández P. (1996) Formicidae (Hymenoptera), pp. 483-500 en J. Llorente, A. N. García y E. González, eds., *Biodiversidad, taxonomía y biogeografía de artrópodos de México: hacia una síntesis de su conocimiento*. Universidad Nacional Autónoma de México.
- Santschi F. (1923) Revue des fourmis du genre "Brachymyrmex" Mayr. *Anales del Museo Nacional de Historia Natural de Buenos Aires* 31:650-678.
- Santschi F. (1925) Revision du genre *Acromyrmex* Mayr. *Revue Suisse Zoologie* 31:355-398.
- Schultz T. R., R. Meier (1995) A phylogenetic analysis of the fungus-growing ants (Hymenoptera: Formicidae: Attini) based on morphological characters of the larvae. *Systematic Entomology* 20:337-370.
- Schultz T. R., D. Bekkevold, J.J. Boomsma. 1998. *Acromyrmex insinuator* new species: an incipient social parasite of fungus-growing ants. *Insectes Sociaux* 45:457-471.

- Schultz T.R., S.A. Solomon, U.G. Mueller, P. Villesen, J.J. Boomsma, R.M.M. Adams, B. Norden (2002) Cryptic speciation in the fungus growing ants *Cyphomyrmex longiscapus* Weber and *Cyphomyrmex muelleri* Schultz and Solomon, new species (Formicidae, Attini) *Insectes Sociaux* 49:331-343.
- Shattuck S. O. (1992a) Generic revision of the ant subfamily Dolichoderinae (Hymenoptera: Formicidae) *Sociobiology* 21(1):1-181.
- Shattuck S. O. (1992b) Higher classification of the ant subfamilies Aneuretinae, Dolichoderinae and Formicinae (Hymenoptera: Formicidae) *Systematic Entomology* 17:199-206.
- Shattuck S. (1994) Taxonomic catalog of the ant subfamilies Aneuretinae and Dolichoderinae *University of California Publications in Entomology* 112:1-241.
- Shattuck S. O. (1995) Generic-level relationships within the ant subfamily Dolichoderinae (Hymenoptera: Formicidae) *Systematic Entomology* 20:217-228.
- Smith M. R. (1944) The genus *Lachnomyrmex*, with the description of a second species *Proceedings of the Entomological Society of Washington* 46:225-228.
- Smith M. R. (1962) A remarkable new *Stenamma* from Costa Rica, with pertinent facts on other Mexican and Central American species *Journal of the New York Entomological Society* 70:33-38.
- Snelling R. R. (1973) The ant genus *Conomyrma* in the United States (Hymenoptera: Formicidae) *Contributions in Science* 238:1-6.
- Snelling R. R. (1975) Descriptions of the new Chilean ant taxa (Hymenoptera: Formicidae) *Contributions in Science* 274:1-19.
- Snelling R.R. (1976) A revision of the honey ants of the genus *Myrmecocystus*, *Natural History Museum of Los Angeles County Science Bulletin* 24:1-163.
- Snelling R.R. (1982) A revision of the honey ants, genus *Myrmecocystus* first supplement *Bulletin of the Southern California Academy of Sciences* 81:69-86.
- Snelling R.R. (1995) Systematics of Nearctic ants of the genus *Dorymyrmex* (Hymenoptera: Formicidae) *Contributions in Science* 454:1-14.
- Snelling R.R. (2001) Two new species of ThiefAnt (*Solenopsis*) from Puerto Rico *Sociobiology* 37:511-525.
- Snelling, R. R. y J. H. Hunt. (1975) The ants of Chile (Hymenoptera: Formicidae). *Revista Chilena de Entomología* 9:63-130.
- Snelling R. R., J. T. Longino (1992) Revisionary notes on the fungus-growing ants of the genus *Cyphomyrmex*, *rimosus*-group, pp. 479-494 en D. Quintero y A. Aiello, eds., *Insects of Panama and Mesoamerica: selected studies* Oxford University Press.
- Snelling R.R., J.A. Torres (1998) *Camponotus ustus* Forel and two similar new species from Puerto Rico *Contributions in Science* 469:1-10.
- Snelling R.R. (1975) Descriptions of new Chilean ant taxa *Contributions in Science* 274:1-19.
- Taber S. W. (1998) *The World of the Harvester Ants* Texas A&M University Press.
- Taber S. W. (2000) *Fire Ants*. Texas A&M University Press.
- Trager J. P. (1984) A revision of the genus *Paratrechina* (Hymenoptera: Formicidae) of the Continental United States *Sociobiology* 9(2):51-162.
- Trager J.C. (1991) A revision of the fire ants, *Solenopsis geminata* group *Journal of the New York Entomological Society* 99:141-198.
- Ward P. S. (1985) The Nearctic species of the genus *Pseudomyrmex* (Hymenoptera: Formicidae) *Quaestiones Entomologicae* 21:209-246.
- Ward P. S. (1988) Mesic elements in the western Nearctic ant fauna: taxonomic and biological notes on *Amblyopone*, *Proceratium*, and *Smithistruma* (Hymenoptera: Formicidae) *Journal of the Kansas Entomological Society* 61:102-124.
- Ward P. S. (1989) Systematic studies on pseudomyrmecine ants: revision of the *Pseudomyrmex oculatus* and *P. subtilissimus* species groups, with taxonomic comments on other species *Quaestiones Entomologicae* 25:393-468.
- Ward P. S. (1990) The ant subfamily Pseudomyrmecinae (Hymenoptera: Formicidae): generic revision and relationship to other formicids *Systematic Entomology* 15:449-489.
- Ward P. S. (1991) Phylogenetic analysis of pseudomyrmecine ants associated with domatia-bearing plants, pp. 335-352 en C.R. Huxley y D.F. Cutler, eds., *Ant-plant interactions*. Oxford: Oxford University Press.
- Ward P. S. (1992) Ants of the genus *Pseudomyrmex* (Hymenoptera: Formicidae) from Dominican amber, with a synopsis of the extant Antillean species *Psyche* 99:55-85.
- Ward P. S. (1993) Systematic studies on *Pseudomyrmex acacia*-ants (Hymenoptera: Formicidae: Pseudomyrmecinae) *Journal of Hymenoptera Research* 2:117-168.
- Ward P. S. (1994) *Adetomyrma*, an enigmatic new ant genus from Madagascar (Hymenoptera: Formicidae), and its implications for ant phylogeny *Systematic Entomology* 19:159-175.
- Ward P. S. (1996) A new workerless social parasite in the ant genus *Pseudomyrmex* (Hymenoptera: Formicidae), with a discussion of the origin of social parasitism in ants *Systematic Entomology* 21:253-263.
- Ward P. S. (1999) Systematics, biogeography and host plant associations of the *Pseudomyrmex vidiuus* group (Hymenoptera: Formicidae), *Triplaris*- and *Tachigali*-inhabiting ants *Zoological Journal of the Linnean Society* 126:451-540.

- Ward, P. S. (2002) Ants of the subfamily Pseudomyrmecinae. <http://entomology.ucdavis.edu/faculty/ward/pseudo.html>
- Ward P.S., S.G. Brady (2003) Phylogeny and biogeography of the ant subfamily Myrmecinae (Hymenoptera: Formicidae) *Invertebrate Systematics* 17:361-386.
- Weber N. A. (1941) Four new genera of Ethiopian and Neotropical Formicidae *Annals of the Entomological Society of America* 34:183-194.
- Weber N. A. (1972) *Gardening Ants, The Attines American Philosophical Society*, Philadelphia.
- Weber N.A. (1982) Fungus Ants, pp. 255-363 en H.R. Hermann, ed., *Social Insects* Vol. 4 Academic Press, NY.
- Wetterer J.K., T.R. Shultz, R. Meier (1998) Phylogeny of fungus-growing ants (Tribe Attini) based on mtDNA sequence and morphology *Molecular Phylogenetics and Evolution* 9(1):42-47.
- Wheeler G. C., J. Wheeler (1976) Ant larvae: review and synthesis *Memoirs of the Entomological Society of Washington* 7:1-108.
- Wheeler G.C. (1913) A revision of the ants of the genus *Formica* (Linné) Mayr *Bulletin of the Museum of Comparative Zoology* 53:379-565.
- Wild A. L. (2002) The genus *Pachycondyla* (Hymenoptera: Formicidae) in Paraguay *Boletín del Museo Nacional de Historia Natural del Paraguay* 14(1-2):1-18.
- Wilson E. O. (1985) Ants of the fominican amber (Hymenoptera: Formicidae). 3. The subfamily Dolichoderinae *Psyche* 92(1):17-37.
- Wilson E.O. (1955) A monographic revision of the ant genus *Lasius*, *Bulletin of the museum of comparative Zoology at Harvard College* 113:1-201.
- Wilson E.O. (2003) *The genus Pheidole in the New World: A Dominant, Hyperdiverse Ant Genus* Harvard University Press (+ CD).

Appendix 1 / Anexo 1

Synonymic list of supraspecific ant taxa in the Neotropical Region

Lista sinónimica de los taxones supraespecíficos de hormigas en la Región Neotropical

Subfamily Dolichoderinae

Tribe Dolichoderini Forel, 1878

- = Liometopini Dlussky & Fedoseeva, 1988
- = Tapinomini Emery, 1912

Azteca Forel, 1878

- = *Aztecum* Bertkau, 1879

Dolichoderus Lund, 1831

- = *Hypoclinea* Mayr, 1855
- = *Monacus* Roger, 1862

Dorymyrmex Mayr, 1866

- = *Ammomyrma* Santschi, 1922
- = *Araucomyrmex* Gallardo, 1919
- = *Biconomyrma* Kusnezov, 1952
- = *Conomyrma* Forel, 1913
- = *Psammomyrma* Forel, 1912
- = *Spinomyrma* Kusnezov, 1952

Forelius Emery, 1888

- = *Amrymex* Kusnezov, 1953
- = *Neoforelius* Kusnezov, 1953

Subfamily Formicinae

Tribe Camponotini

Camponotus Mayr, 1861

- = *Colobopsis* Mayr, 1861
- = *Dendromyrmex* Emery, 1895

= *Hypercolobopsis* Emery, 1920

= *Manniela* Wheeler, 1921

= *Myrmaphaenus* Emery, 1920

= *Myrmentoma* Forel, 1912

= *Myrmepomis* Forel, 1912

= *Myrmeurynota* Forel, 1912

= *Myrmobrachys* Forel, 1912

= *Myrmocladoeus* Wheeler, 1921

= *Myrmodirhachis* Emery, 1925

= *Myrmomalis* Forel, 1914

= *Myrmoplatus* Santschi, 1921

= *Myrmosiphincta* Forel, 1912

= *Myrmostenus* Emery, 1920

= *Myrmothrix* Forel, 1912

= *Pseudocolobopsis* Emery, 1920

= *Tanaemyrmex* Ashmead, 1905

Tribe Lasiini Ashmead, 1905

- = *Acanthomyopsini* Donisthorpe, 1943

Acropyga Roger, 1862

- = *Rhizomyrma* Forel, 1893

Myrmecocystus Wesmael, 1838

- = *Endiodioctes* Snelling, 1976

Tribe Plagiolepidini Forel, 1886

- = *Brachymyrmicini* Emery, 1825

= *Myrmelachistini* Forel, 1912
 = *Prenolepidini* Emery, 1916

- Brachymyrmex* Mayr, 1868
 = *Bryschia* Santschi, 1923
Myrmelachista Roger, 1863
 = *Decamera* Roger, 1863 (homonym)
 = *Hincksidris* Donisthorpe, 1944
 = *Neaphomus* Menozzi, 1935
Paratrechina Motschoulsky, 1863
 = *Nylanderia* Emery, 1906
 = *Paraparatrechina*

Subfamily Pseudomyrmecinae M.R. Smith, 1952

Tribe Pseudomyrmecini M.R. Smith, 1952

- = *Leptaleinae* M.R. Smith, 1951
 = *Pseudomyrmidae* Forel 1885

- Pseudomyrmex* Lund, 1831
 = *Apedunculata* Enzmann, 1944
 = *Clavanoda* Enzmann, 1944
 = *Latinoda* Enzmann, 1944
 = *Leptalaea* Spinola, 1851
 = *Leptalea* Erichson, 1839
 = *Myrmex* Guérin, 1845 (homonym)
 = *Ornatinoda* Enzmann, 1944
 = *Pseudomyrma* Guérin, 1844
 = *Triangulinoda* Enzmann, 1944

Subfamily Cerapachyinae Forel, 1893

Tribe Acanthostichini

- Acanthostichus* Mayr, 1887
 = *Ctenopyga* Ashmead, 1906

Tribe Cylindromyrmecini

- Cylindromyrmex* Mayr, 1870
 = *Holcoponera* Cameron, 1891
 = *Hypocylindromyrmex* Wheeler, 1924
 = *Metacylindromyrmex* Wheeler, 1924

Subfamily Ecitoninae

Tribe Ecitonini

- Eciton* Latreille, 1804
 = *Ancylognathus* Lund, 1831
 = *Camptognatha* Gray, 1832
 = *Holopone* Santschi, 1924
 = *Mayromyrmex* Ashmead, 1905

- Labidus* Jurine, 1807
 = *Nycteresia* Roger, 1861
 = *Pseudodichthadia* E. André, 1885
Neivamyrmex Borgmeier, 1940
 = *Acamatus* Emery, 1894 (homonym)
 = *Woitkowskia* Enzmann, 1952

Subfamily Amblyoponinae

Tribe Amblyoponini

- Amblyopone* Erichson 1842
 = *Ericapelta* Kusnezov, 1955

Subfamily Ponerinae

Tribe Ponerini Lepeletier, 1835

- = *Centromyrmicini* Emery, 1911
 = *Euponerinae* Emery, 1909
 = *Leptogenysii* Forel, 1893
 = *Odontomachidae* Mayr, 1862
 = *Pachycondylinae* Ashmead, 1905

- Anochetus* Mayr, 1861
 = *Stenomyrmex* Mayr, 1862

- Belonopelta* Mayr, 1870
 = *Leiopelta* Baroni Urbani, 1975

- Centromyrmex* Mayr, 1866

- Leptogenys* Roger, 1861
 = *Lobopelta* Mayr, 1862
- Pachycondyla* F. Smith, 1858
 = *Mesoponera* Emery, 1901
 = *Neoponera* Emery, 1901
 = *Termitopone* Wheeler, 1936
 = *Trachymesopus* Emery, 1911
 = *Wadeura* Weber, 1939

Subfamily Ectatomminae

Tribe Ectatommini

- Gnamptogenys* Roger, 1863
 = *Alfaria* Emery, 1896
 = *Barbourella* Wheeler, 1930
 = *Commateta* Santschi, 1929
 = *Emeryella* Forel, 1901
 = *Holcoponera* Mayr, 1887
 = *Opisthoscyphus* Mann, 1922
 = *Parectatomma* Emery, 1911
 = *Poneracantha* Emery, 1897
 = *Rhopalopone* Emery, 1897
 = *Spaniopone* Wheeler W.M. & Mann, 1914
 = *Tammoteca* Santschi, 1929

Subfamily Heteroponerinae**Tribe Heteroponerini***Acanthoponera* Mayr, 1862

- = *Anacanthoponera* Wheeler, 1923
- = *Paranomopone* Wheeler, 1915

Subfamily Proceratiinae**Tribe Proceratiini Emery, 1895**

- = *Discothyridinae* Clark, 1951

Proceratium Roger, 1863

- = *Sysphingta* Roger, 1863

Subfamily Myrmicinae**Tribe Adelomyrmecini***Adelomyrmex* Emery, 1897

- = *Apyschomyrmex* Wheeler, 1910

Tribe Attini*Atta* Fabricius, 1804

- = *Archeatta* Gonçalves 1942
- = *Epiatta* Borgmeier, 1950
- = *Myrmegis* Rafinesque, 1815 (*Nomen nudum*)
- = *Neoatta* Gonçalves 1942
- = *Oecodoma* Latreille, 1818
- = *Palaeatta* Borgmeier, 1950.

Cyphomyrmex Mayr, 1862

- = *Cyphomannia* Weber, 1938

Mycetophylax Emery, 1913

- = *Mycetopurus* Santschi, 1925
- = *Paramycetophylax* Kusnezov, 1956

Mycocerus Forel, 1893

- = *Descolemyrma* Kusnezov, 1951

Myrmicocrypta F. Smith, 1860

- = *Glyptomyrmex* Forel, 1884

Tribe Basicerotini*Basicerca* Schulz, 1906

- = *Aspididris* Weber, 1950
- = *Ceratobasis* F. Smith, 1860 (homonym)
- = *Creigthonidris* Brown, 1949

Rhopalothrix Mayr, 1870

- = *Acanthidris* Weber, 1941
- = *Heptastruma* Weber, 1934

Tribe Blepharidattini*Wasmannia* Forel, 1893

- = *Hercynia* J. Enzmann, 1947

Tribe Cephalotini M.R. Smith, 1949

- = *Cryptoceridae* F. Smith, 1854

Cephalotes Latreille, 1802

- = *Cryptocerus* Latreille, 1803
- = *Cyathomyrmex* Creighton, 1933
- = *Cyathocephalus* Emery, 1915
- = *Eucryptocerus* Kempf, 1951
- = *Exocryptocerus** Vierbergen & Scheven, 1995.
- = *Harnedia* Smith M.R., 1949
- = *Paracryptocerus* Emery, 1915
- = *Zacryptocerus* Wheeler, 1911

Tribe Crematogastrini*Crematogaster* Lund, 1831

- = *Acrocoelia* Mayr, 1852
- = *Cremastogaster* Agassiz, 1846
- = *Tranopeltoides* Wheeler, 1922

Tribe Dacetini Forel, 1892

- = *Dacetonini* Forel, 1892

Daceton Perty, 1833

- = *Dacetum* Agassiz, 1848

Pyramica Roger, 1862

- = *Epitritus* Emery, 1869
- = *Trichoscapa* Emery, 1869
- = *Cephaloxys* F. Smith, 1865
- = *Glamyromyrmex* Wheeler, 1915
- = *Codiomyrmex* Wheeler, 1916
- = *Tingimyrmex* Mann, 1926
- = *Codioxenus* Santschi, 1931
- = *Smithistruma* Brown, 1948
- = *Neostruma* Brown, 1948
- = *Dorisidris* Brown, 1948
- = *Chelystruma* Brown, 1950
- = *Borgmeierita* Brown, 1953
- = *Gymnomyrmex* Borgmeier, 1954

Strumigenys F. Smith, 1860

- = *Quadristruma* Brown, 1949

Tribe Formicoxenini Forel, 1893

- = *Cardiocondylini* Emery, 1914
- = *Leptocephalini* Emery, 1914

Nesomyrmex Wheeler, 1910

- = *Goniothorax* Emery, 1896 (homonym)
- = *Caulomyrma* Forel, 1914

- Temnothorax* Mayr 1861
 = *Macromischa* Roger, 1863
 = *Antillaemymrmex* Mann, 1920
 = *Croesomyrmex* Mann, 1920
 = *Myrafant* M.R. Smith, 1950
Ochetomyrmex Mayr, 1878
 = *Brownidris* Kusnezov, 1956
Xenomyrmex Forel 1885
 = *Myrmecinella* Forel, 1884

Tribe Myrmicini

- Hylomyrma* Forel, 1912
 = *Lundella* Emery, 1915
Pogonomyrmex Mayr, 1868
 = *Ephebomyrmex* Wheeler, 1902
 = *Forelomyrmex* Wheeler, 1913
 = *Janetia* Forel, 1899 (homonym)

Tribe Pheidolini Emery, 1877

- = *Aphaenogastrini* Enzmann, 1947
- Pheidole* Westwood, 1839
 = *Allopheidole* Forel, 1912
 = *Bruchomyrma* Santschi, 1922
 = *Cardiopheidole* Wheeler, 1914
 = *Ceratopheidole* Pergande, 1896
 = *Decapheidole* Forel, 1912
 = *Elasmopheidole* Forel, 1913
 = *Eriopheidole* Kusnezov, 1951
 = *Gallardomyrma* Bruch, 1932
 = *Hendecapheidole* Wheeler, 1922
 = *Macropheidole* Emery, 1915
 = *Oecophthora* Heer, 1852
 = *Scrobopheidole* Emery, 1915
 = *Trachypheidole* Emery, 1915

Tribe Solenopsidini Forel, 1893

- = *Megalomyrmecini* Dlussky & Fedoseeva, 1988
 = *Monomoriini* Emery, 1914
 = *Pheidologetini* Emery, 1914

Carebara Westwood, 1840

- = *Erebomyrma* Wheeler, 1903
 = *Oligomyrmex* Mayr, 1867
 = *Spelaeomyrmex* Wheeler, 1922

Megalomyrmex Forel, 1885

- = *Cepobroticus* Wheler, 1925
 = *Wheelerimyrmex* Mann, 1922

Monomorium Mayr, 1855

- = *Antichthonidris* Snelling, 1975
 = *Mitara* Emery, 1913
 = *Nothidris* Ettershank, 1966

Oxyepoecus Santschi, 1926

- = *Forelididis* M.R. Smith, 1954
 = *Martia* Forel, 1907 (homonym)

Solenopsis Westwood, 1840

- = *Bisolenopsis* Kusnezov, 1953
 = *Carebarella* Emery, 1906
 = *Carebarelloides* Borgmeier, 1937
 = *Diagyne* Santschi, 1923
 = *Diplorhoptrum* Mayr, 1855
 = *Euopthalma* Creighton, 1930
 = *Granisolenopsis* Kusnezov, 1957
 = *Labauchena* Santschi, 1930
 = *Lilidris* Kusnezov, 1957
 = *Octella* Forel, 1915
 = *Oedaleocerus* Creighton, 1930
 = *Paranamyrmex* Kusnezov, 1954
 = *Synsolenopsis* Forel, 1918

Tribe Stenammini

- Rogeria* Emery, 1894
 = *Irogera* Emery, 1915

Appendix 2 / Anexo 2

Synonymic list of Neotropical ant species. For details see Kempf (1972), Brandão (1991) and Bolton (1995a, 2003). The extensive synonymies in *Camponotus* (MacKay, in preparation) are not included.

Lista sinónímica de las especies de hormigas Neotropicales. Para detalles por favor consultar Kempf (1972), Brandão (1991) y Bolton (1995a, 2003). La extensa sinonimia en Camponotus (MacKay, en preparación) no se incluye.

Subfamily Dolichoderinae

- Genus *Azteca* Forel, 1878
Azteca alfari Emery, 1894
 = *breviscapus* Forel, 1912
 = *cecropiae* Forel, 1906
 = *curtiscpa* Forel, 1912

- = *lucidula* Forel, 1899
 = *langi* Wheeler W.M., 1942
 = *fumaticeps* Wheeler W.M., 1942
 = *lynchi* Brethes, 1914
 = *mixta* Forel, 1908
 = *virens* Forel, 1899
 = *bicolor* Emery, 1893

- Azteca aurita* Emery, 1893
 = *pilosula* Forel, 1899
- Azteca constructor* Emery, 1896
 = *emmae* Forel, 1904
 = *guianae* Wheeler W.M., 1942
- Azteca ovaticeps* Forel, 1904
 = *alfari aequilata* Forel, 1904
 = *alfari tuberosa* Forel, 1906
 = *alfari aequalis* Forel, 1906
- Azteca trigona* Emery, 1893
 = *festai* Emery, 1896
- Azteca xanthochroa* (Roger, 1863)
 = *costaricensis* Wheeler W.M., 1942
- Genus Dolichoderus Lund, 1831**
- Dolichoderus abruptus* (Smith, 1858)
 = *ursus* Mayr, 1866
- Dolichoderus attelaboides* (Fabricius, 1775)
 = *pulla* Santschi, 1923
 = *heterogaster* Santschi, 1923
 = *imbecillus* Mann, 1916
- Dolichoderus bidens* L., 1758
 = *perditor* (Fabricius, 1804)
 = *auromaculatus* Forel, 1885
 = *attenuatus* Forel, 1903
 = *bahiana* Santschi, 1921
- Dolichoderus debilis* Emery, 1890
 = *parabiotica* Forel, 1912
 = *sieversi* Forel, 1901
- Dolichoderus decollatus* Smith, 1858
 = *capitatus* Santschi, 1921
 = *neglectus* Menozzi, 1935
- Dolichoderus diversus* Emery, 1894
 = *championi* Forel, 1899
 = *garbei* Forel, 1911
 = *leviusculus* Emery, 1906
 = *mazaruni* Donisthorpe, 1939
 = *ornatus* Mann, 1916
 = *taeniatus* Wheeler W.M., 1916
 = *trinidadensis* Forel, 1899
- Dolichoderus laminatus* (Mayr, 1870)
 = *luteiventris* Emery, 1894
- Dolichoderus lutosus* (Smith, 1858)
 = *cingulata* (Mayr, 1862)
 = *nigriventris* Forel, 1893
- Dolichoderus plagiatus* (Mayr, 1870)
 = *borealis* Provancher, 1888
 = *inornatus* Wheeler W.M., 1905
- Dolichoderus quadridenticulatus* (Roger, 1862)
 = *analis* Emery, 1894
 = *gibbosoanalis* Forel, 1922
 = *integra* Forel, 1911
 = *nitidor* Emery, 1894

- Dolichoderus schulzi* Emery, 1894
 = *bolleyi* Forel, 1908
 = *columbica* Forel, 1914
- Dolichoderus spurius* Forel, 1903
 = *albatus* Vihemeyer, 1922
- Dolichoderus taschenbergi* (Mayr, 1866)
 = *aterrimus* Wheeler W.M., 1915
- Genus Dorymyrmex Mayr, 1866**
- Dorymyrmex antarcticus* Forel, 1904
 = *pallidipes* Brethes, 1914
- Dorymyrmex insanus* (Buckley, 1866)
 = *smithi* Cole, 1936
 = *wheeleri* Kusnezov, 1952
 = *medeis* Trager, 1988
 = *reginicula* Trager, 1988

- Genus Tapinoma Foester, 1850**
- Tapinoma antarcticum* Forel, 1904
 = *eidmanni* Goetsch, 1933
- Tapinoma melanocephalum* (Fabricius, 1793)
 = *familiaris* (Smith, 1860)
- Tapinoma panamense* Wheeler W.M., 1934
 = *canalis* Wheeler W.M., 1942

Subfamily Formicinae

Tribe Formicini

- Genus Formica L., 1758**
- Formica argentea* Wheeler W.M., 1912
 = *blanda* Wheeler W.M., 1913
- Formica integroides* Wheeler W.M., 1913
 = *subfasciata* Creigthon, 1940
- Formica lasioides* Emery, 1893
 = *vetula* Wheeler W.M., 1912
- Formica moki* Wheeler W.M., 1906
 = *occidua* Wheeler W.M., 1913
- Formica xerophila* Smith M.R., 1939
 = *grundmanni* Cole, 1943

Tribe Lasiini

- Genus Acropyga Roger, 1862**
- Acropyga donisthorpei* Weber, 1944
 = *oko* Weber, 1944
- = *paludis* Weber, 1944
- Acropyga fuhrmanni* (Forel, 1914)
 = *berwicki* (Wheeler, 1935)
- Acropyga exsanguis* (Wheeler, 1909)
 = *wheeleri* Mann, 1922
 = *bruchi* Santschi, 1929
 = *paramaribensis* Borgmeier, 1933
 = *robae* Donisthorpe, 1936

- Acropyga goeldii* Forel, 1893
 = *pachycera* (Emery, 1906)
 = *goeldii columbica* (Forel, 1912)
 = *goeldii tridentata* (Forel, 1912)
 = *marshalli* (Crawley, 1921)
 = *pickeli* Borgmeier, 1927
 = *rutgersi* Bünzli, 1935
 = *borgmeieri* Donisthorpe, 1939
 = *cuadriceps* Weber, 1944
 = *trinitatis* Weber, 1944
Acropyga parvidens (Wheeler & Mann, 1914)
 = *mesonotalis* Weber, 1944
Acropyga smithii Forel, 1893
 = *kathrynae* Weber, 1944
 = *urichi* Weber, 1944

Genus *Anoplolepis* Santschi, 1914

- Anoplolepis gracilipes* (Smith, 1857)
 = *longipes* (Jerdon 1851)

Genus *Lasiophanes* Forel, 1895

- Lasiophanes picinus* (Roger, 1863)
 = *bruchi* (Forel, 1915)
 = *edwardsi* (Donisthorpe, 1933)
Lasiophanes valdiviensis (Forel, 1904)
 = *bolivari* (Santschi, 1916)
 = *pilosulus* (Emery, 1922)
 = *uxorius* (Emery, 1922)

Genus *Lasius* Fabricius, 1804

- Lasius alienus* (Foester, 1850)
 = *americanus* Emery, 1893
 = *pannonica* Roszler, 1942

Genus *Myrmecocystus* Wesmael, 1838

- Myrmecocystus melliger* Forel, 1886
 = *comatus* Creigthon, 1950
Myrmecocystus mexicanus Wesmael, 1838
 = *hortusdeorum* McCook, 1881
Myrmecocystus mimicus Wheeler W.M., 1908
 = *lomaensis* Wheeler W.M., 1912

Tribus Plagiolepidini

Genus *Brachymyrmex* Mayr, 1868

- Brachymyrmex depilis* Emery, 1893
 = *nanellus* Wheeler W.M., 1903
Brachymyrmex giardi Emery, 1895
 = *nitida* Santschi, 1922

Genus *Myrmelachista* Roger, 1863

- Myrmelachista arthuri* Forel, 1903
 = *ute* Kusnezov, 1969

- Myrmelachista gagatina* Emery, 1894
 = *ina* Kusnezov, 1969
Myrmelachista gallicola Mayr, 1887
 = *rubiginosa* Emery, 1906
Myrmelachista hoffmanni Forel, 1903
 = *rectinota* Forel, 1904
Myrmelachista mayri Forel, 1886
 = *monticola* Mayr, 1887

Genus *Paratrechina* Motschoulsky, 1863

- Paratrechina longicornis* (Latreille, 1802)
 = *gracilis* (Nylander, 1856)

Subfamily Pseudomyrmecinae

Genus *Pseudomyrmex* Lund, 1831

- Pseudomyrmex boopis* (Roger, 1863)
 = *excavata* (Mayr, 1870)
 = *flaviventris* (Emery, 1896)
 = *guatemalensis* (Enzmann, 1944)
Pseudomyrmex brunneus (Smith, 1877)
 = *nigrita* (Enzmann, 1944)
Pseudomyrmex concolor (Smith, 1860)
 = *latinoda* (Mayr, 1878)
 = *penetrator* (Smith, 1877)
Pseudomyrmex dendroicus (Forel, 1904)
 = *emarginata* (Forel, 1904)
Pseudomyrmex distinctus (Smith, 1877)
 = *puchella* (Forel, 1899)
Pseudomyrmex ejectus (Smith, 1858)
 = *lincecumii* (Buckley, 1866)
Pseudomyrmex elongatus (Mayr, 1870)
 = *alliodorae* (Wheeler W.M., 1942)
 = *tandem* (Forel, 1906)
Pseudomyrmex euryblemma (Forel, 1899)
 = *kurokii* (Forel, 1906)
Pseudomyrmex faber (Smith, 1858)
 = *perforator* (Smith, 1860)
Pseudomyrmex ferrugineus (Smith, 1877)
 = *bequaerti* (Wheeler W.M., 1942)
 = *fulvescens* (Emery, 1890)
 = *hondurana* (Enzmann, 1944)
 = *saffordi* (Enzmann, 1944)
 = *vesana* (Wheeler W.M., 1942)
Pseudomyrmex filiformis (Fabricius, 1804)
 = *cephalotes* (Smith, 1855)
 = *longiceps* Forel, 1906
Pseudomyrmex godmani (Forel, 1899)
 = *semotus* Kempf, 1960
Pseudomyrmex gracilis (Fabricius, 1804)
 = *bicolor* (Guerin-Meneville, 1844)
 = *canescens* (Wasmann, 1915)
 = *dimita* (Roger, 1863)
 = *glabriventris* (Santschi, 1922)

- = *mexicana* (Roger, 1863)
- = *pilosula* (Smith, 1877)
- = *sericata* (Smith, 1858)
- = *velifera* (Stitz, 1933)
- = *volatilis* (Smith, 1877)
- Pseudomyrmex kuenckeli* (Emery, 1890)
 - = *bierigi* (Santschi, 1932)
 - = *crenulata* (Enzmann, 1944)
 - = *dichroa* (Forel, 1904)
- Pseudomyrmex maculatus* (Smith, 1855)
 - = *laticeps* (Forel, 1906)
- Pseudomyrmex nigrocinctus* (Emery, 1890)
 - = *alfari* (Forel, 1906)
 - = *bicina* (Santschi, 1922)
 - = *peltata* (Menozzi, 1927)
- Pseudomyrmex oculatus* (Smith, 1855)
 - = *dolichopsis* (Forel, 1899)
 - = *implicata* Forel, 1911
 - = *altinoda* (Mann, 1916)
 - = *tuberculata* (Enzmann, 1944)
 - = *wessoni* (Enzmann, 1944)
- Pseudomyrmex osurus* (Forel, 1911)
 - = *insularis* (Enzmann, 1944)
- Pseudomyrmex peperi* (Forel, 1913)
 - = *convarians* (Forel, 1913)
- Pseudomyrmex perboscii* (Guerin-Meneville, 1844)
 - = *icterica* (Wheeler W.M., 1922)
 - = *simoides* (Forel, 1911)
 - = *testacea* (Smith, 1852)
- Pseudomyrmex phyllophilus* (Smith, 1858)
 - = *mutica* (Mayr, 1887)
- Pseudomyrmex rufomedium* (Smith, 1877)
 - = *stolli* (Forel, 1912)
- Pseudomyrmex schuppi* (Forel, 1901)
 - = *confusior* (Forel, 1901)
 - = *geraensis* (Forel, 1912)
- Pseudomyrmex simplex* (Smith, 1877)
 - = *capperi* (Forel, 1899)
 - = *delicatula* (Forel, 1899)
 - = *panamensis* (Forel, 1899)
- Pseudomyrmex spinicola* (Emery, 1890)
 - = *atrox* (Forel, 1912)
 - = *gaigei* (Forel, 1914)
 - = *infernalis* (Wheeler W.M., 1942)
- Pseudomyrmex tachigaliae* (Forel, 1904)
 - = *bradleyi* (Wheeler W.M., 1942)
- Pseudomyrmex termitarius* (Smith, 1855)
 - = *elegans* (Smith, 1855)
 - = *infusca* (Forel, 1908)
- Pseudomyrmex triplaridis* (Forel, 1904)
 - = *boxi* (Wheeler W.M., 1942)
 - = *boxi* (Enzmann, 1944)
- Pseudomyrmex triplarinus* (Weddell, 1850)
 - = *arborissanctae* (Emery, 1894)
 - = *ecuadoriana* (Enzmann, 1944)

- Pseudomyrmex veneficus* (Wheeler W.M., 1942)
 - = *venifica* (Enzmann, 1944)
- Pseudomyrmex viduus* (Smith, 1858)
 - = *baileyi* (Wheeler W.M., 1942)
 - = *caroli* (Forel, 1899)
 - = *trigrina* (Wheeler W.M., 1942)
 - = *ulei* (Forel, 1904)
 - = *sapii* (Forel, 1904)

Subfamily Cerapachyinae

Tribe Acanthostichini

Genus *Acanthostichus* Mayr, 1887

- Acanthostichus brevicornis* Emery, 1894
 - = *ramosmexiae* Bruch, 1924
- Acanthostichus laticornis* Forel, 1908
 - = *obscuridens* Santschi, 1934
- Acanthostichus texanus* Forel, 1904
 - = *townsendi* Ashmead, 1906

Tribe Cylindromyrmecini

Genus *Cylindromyrmex* Mayr, 1870

- Cylindromyrmex striatus* Mayr, 1870
 - = *whymperi* (Cameron, 1891)
 - = *tibialis* Stitz, 1932

Subfamily Ecitoninae

Tribe Cheliomyrmecini

Genus *Cheliomyrmex* Mayr, 1870

- Cheliomyrmex andicola* Emery, 1894
 - = *ursinum* Emery, 1901
- Cheliomyrmex morosus* (Smith, 1859)
 - = *nortoni* Mayr, 1870
 - = *payarum* (Forel, 1899)

Tribe Ecitonini

Genus *Eciton* Latreille, 1804

- Eciton drepanophorum* (Smith, 1858)
 - = *bellicosum* Santschi, 1921
 - = *quitensis* Forel, 1921
 - = *ferox* Santschi, 1920
- Eciton dulcium* Forel, 1912
 - = *hirsutum* Santschi, 1912
- Eciton hamatum* (Fabricius, 1782)
 - = *curvidentata* (Latreille, 1802)
 - = *amazona* Santschi, 1923
 - = *funesta* Santschi, 1921
 - = *mattogrossensis* Luederwalt, 1920
 - = *pittieri* Forel, 1899

Eciton mexicanum Roger, 1863
 = *rogeri* Dalla Torre, 1892

Eciton vagans (Oliver, 1792)
 = *simillima* Smith, 1855

Genus *Labidus* Jurine, 1807

- Labidus coecus* (Latrelle, 1802)
 = *vasator* (Smith, 1860)
 = *jurunii* Shuckard, 1840
 = *erratica* Smith, 1860
 = *latreilli* Jurine, 1807
 = *rubra* Buckley, 1867
 = *saji* Haldeman, 1852
 = *atriceps* Smith, 1859
 = *incerta* (André, 1885)
 = *fulvescens* (Blanchard, 1846)
 = *pilosus* Smith, 1859
 = *smithi* (Dalla Torre, 1892)
 = *grassator* Forel, 1911
 = *panzeri* Smith, 1859
 = *bilboa* Emery, 1901
 = *elsbethae* Forel, 1922
 = *servillei* Westwood, 1842
 = *opacifrons* (Wheeler W.M., 1921)
 = *kulowi* (Forel, 1901)
 = *selysi* (Forel, 1904)
 = *hostilis* Santschi, 1920
- Labidus praedator* (Smith, 1858)
 = *westwoodi* (Emery, 1900)
 = *emiliae* (Mann, 1916)
 = *guianense* (Wheeler W.M., 1921)
 = *ferruginea* (Norton, 1868)

Genus *Neivamyrmex* Borgmeier, 1940

- Neivamyrmex adnepos* (Wheeler W.M., 1922)
 = *marginatum* (Borgmeier, 1939)
- Neivamyrmex andrei* (Emery, 1901)
 = *oslari* (Wheeler W.M., 1908)
- Neivamyrmex californicus* (Mayr, 1870)
 = *obscura* (Forel, 1914)
- Neivamyrmex dorbignii* (Shuckard, 1840)
 = *spiegazzinii* (Emery, 1888)
- Neivamyrmex emeryi* (Santschi, 1921)
 = *exagona* (Santschi, 1923)
- Neivamyrmex falcifer* (Emery, 1900)
 = *mesognathum* (Santschi, 1920)
- Neivamyrmex guyanensis* (Santschi, 1916)
 = *nocua* (Santschi, 1921)
- Neivamyrmex halidaii* (Shuckard, 1840)
 = *frontalis* (Menozzi, 1924)
 = *gravenhorstii* (Westwood, 1842)
 = *amplipennis* (Smith, 1859)
 = *colombi* (Santschi, 1921)
 = *lemoulti* (Santschi, 1912)

Neivamyrmex harrisi (Haldeman, 1852)

= *wheeleri* (Emery, 1901)

Neivamyrmex hetschkoi (Mayr, 1886)

= *taunayi* (Borgmeier, 1928)

= *oglobini* (Santschi, 1934)

= *silvestrii* (Emery, 1906)

Neivamyrmex jermannii (Forel, 1901)

= *luederwaldti* (Emery, 1911)

= *filicorne* (Santschi, 1916)

Neivamyrmex melanocephalus (Emery, 1895)

= *xipe* (Wheeler W.M., 1914)

Neivamyrmex opacithorax (Emery, 1894)

= *castaneum* (Borgmeier, 1939)

Neivamyrmex pertii (Shuckard, 1840)

= *illigeri* (Shuckard, 1840)

= *fonscolobbi* (Westwood, 1842)

Neivamyrmex pilosus (Smith, 1858)

= *angustius* (Forel, 1909)

= *porrectognathum* (Borgmeier, 1933)

Neivamyrmex planidorsus (Emery, 1906)

= *paranaense* (Forel, 1912)

Neivamyrmex pseudops (Forel, 1909)

= *grandipseudops* (Forel, 1915)

= *garbei* Forel, 1912

Neivamyrmex raptor (Forel, 1911)

= *raptans* (Forel, 1911)

Neivamyrmex romandii (Shuckard, 1840)

= *desartsii* (Forel, 1907)

Neivamyrmex sulcatus (Mayr, 1868)

= *richeri* (Forel, 1913)

Neivamyrmex swainsonii (Shuckard, 1840)

= *arizonenese* (Wheeler W.M., 1908)

Genus *Nomamyrmex* Borgmeier, 1936

Nomamyrmex esenbeckii Westwood, 1842

= *fimbriatum* (Santschi, 1920)

= *interrupta* (Santschi, 1923)

= *terrificum* (Santschi, 1930)

= *afflictus* (Gallardo, 1919)

= *crudele* (Santschi, 1929)

= *indomabile* (Santschi, 1930)

Nomamyrmex hartigii (Westwood, 1842)

= *apicifer* (Santschi, 1916)

= *hansi* (Forel, 1912)

= *schlechtendali* (Mayr, 1887)

Subfamily Amblyoponinae

Tribe Amblyoponini

Genus *Amblyopone* Erichson, 1842

Amblyopone elongata (Santschi, 1912)

= *barretoi* (Bruch, 1921)

= *minor* (Santschi, 1922)

= *paranense* (Santschi, 1925)

Genus *Prionopelta* Mayr, 1866

- Prionopelta punctulata* Mayr, 1866
 = *bruchi* Santschi, 1923
 = *mayri* Forel, 1909

Subfamily Ponerinae**Tribe Ponerini****Genus *Anochetus* Mayr, 1861**

- Anochetus altisquamis* Mayr, 1887
 = *fumata* Luederwalt, 1918
Anochetus diegensis Forel, 1912
 = *bierigi* Santschi, 1931
Anochetus emarginatus (Fabricius, 1804)
 = *quadrispinosus* (Smith, 1858)
 = *rugosus* Emery, 1890
Anochetus inermis André, 1889
 = *meinerti* Forel, 1905
Anochetus neglectus Emery, 1894
 = *australis* Emery, 1906

Genus *Centromyrmex* Mayr, 1866

- Centromyrmex brachycola* (Roger, 1861)
 = *bohemanni* Mayr, 1866
 = *paulina* Forel, 1911

Genus *Dinoponera* Roger, 1861

- Dinoponera australis* Emery, 1901
 = *brevis* Borgmeier, 1937
Dinoponera gigantea (Perty, 1833)
 = *grandis* (Guerin-Meneville, 1838)
Dinoponera quadriceps Kempf, 1971
 = *opaca* Kempf, 1971

Genus *Hypoponera* Santschi, 1938

- Hypoponera menozzii* (Santschi, 1932)
 = *longiceps* (Menozzi, 1931)
Hypoponera opaciceps (Mayr, 1887)
 = *perkinsi* (Forel, 1899)
Hypoponera opacior (Forel, 1893)
 = *chilensis* (Forel, 1914)
Hypoponera punctatissima (Roger, 1859)
 = *androgyna* (Roger, 1859)
 = *schauinslandi* (Emery, 1899)

Genus *Leptogenys* Roger, 1861

- Leptogenys crudelis* (Smith, 1858)
 = *rubicunda* Borgmeier, 1930
Leptogenys maxillosa (Smith, 1858)
 = *falcata* Roger, 1861
 = *cribata* Emery, 1895

Genus *Odontomachus* Latreille, 1804

- Odontomachus bauri* Emery, 1892
 = *microcephalus* Emery, 1890
 = *notata* Mann, 1920
 = *paucidens* Emery, 1893
 = *pubescens* Roger, 1861
 = *rugiscuama* Forel, 1908
 = *sericeus* Provancher, 1895
Odontomachus brunneus (Patton, 1894)
 = *fuscus* Stitz, 1916
Odontomachus caelatus Brown, 1976
Odontomachus chelifer (Latreille, 1802)
 = *emacerata* Santschi, 1922
 = *leptocephalus* Emery, 1890
Odontomachus clarus Roger, 1861
 = *clarionensis* Wheeler W.M., 1934
 = *coninodis* Wheeler W.M., 1915
 = *desertorum* Wheeler W.M., 1915
 = *texana* Buckley, 1867
Odontomachus haematodus (L., 1758)
 = *maxilosa* (De Geer, 1773)
 = *hirsutiusculus* Smith, 1858
 = *pallipes* Crawley, 1916
Odontomachus laticeps Roger, 1861
 = *striativentris* Emery, 1890
Odontomachus meinerti Forel, 1905
 = *minuta* Luederwaldt, 1918

Genus *Pachycondyla* Smith, 1858

- Pachycondyla constricta* (Mayr, 1884)
 = *josephi* (Forel, 1886)
Pachycondyla crenata (Roger, 1861)
 = *confusa* (Santschi, 1921)
 = *fiebrigi* (Forel, 1912)
 = *stipitum* (Forel, 1901)
 = *sulcatula* (Santschi, 1919)
Pachycondyla harpax (Fabricius, 1804)
 = *concina* Wheeler W.M., 1925
 = *dibullana* Forel, 1901
 = *irina* Wheeler W.M., 1925
 = *montezumia* Smith, 1858
Pachycondyla impressa (Roger, 1861)
 = *andicola* Santschi, 1913
 = *cearensis* Forel, 1901
 = *inca* Emery, 1901
 = *montana* Forel, 1912
 = *purpurascens* Forel, 1899
 = *transversa* Emery, 1890
Pachycondyla laevigata (Smith, 1858)
 = *gagatina* Emery, 1890
 = *whitleyi* (Wheeler W.M., 1922)
Pachycondyla stigma (Fabricius, 1804)
 = *americana* (Mayr, 1862)
 = *quadridentata* (Smith, 1859)

Pachycondyla striatinodis Emery, 1890
 = *bakeri* (Mann, 1916)

Pachycondyla unidentata Mayr, 1862
 = *eburneipes* (Wheeler W.M. & Bequaert, 1929)
 = *maya* (Wheeler W.M. & Bequaert, 1929)
 = *rugosula* (Emery, 1902)
 = *trinidadensis* (Wheeler W.M. & Bequaert, 1929)

Pachycondyla villosa (Fabricius, 1804)
 = *bicolor* (Guerin-Meneville, 1844)
 = *pedunculata* (Smith, 1858)
 = *pilosa* (Smith, 1858)

Tribe Thaumatomyrmecini

Genus *Thaumatomyrmex* Mayr, 1887

Thaumatomyrmex atrox Weber, 1939
 = *manni* Weber, 1939
 = *zeteki* Smith M.R., 1944

Thaumatomyrmex ferox Mann, 1922
 = *paludis* Weber, 1942

Tribe Platythyreini

Genus *Platythyrea* Roger, 1863

Platythyrea pilosula (Smith, 1858)
 = *incerta* Emery, 1890

Platythyrea punctata (Smith, 1858)
 = *cineracea* Forel, 1886
 = *pruinosa* Mayr, 1870

Platythyrea sinuata (Roger, 1860)
 = *boliviiana* Santschi, 1921
 = *meinerti* Forel, 1905

Subfamily Ectatomminae

Tribe Ectatommini

Genus *Ectatomma* Smith, 1858

Ectatomma edentatum Roger, 1863
 = *inversa* Santschi, 1912
 = *iris* Forel, 1909
 = *macdonaghi* Forel, 1915
 = *morgani* Forel, 1912

Ectatomma muticum Mayr, 1870
 = *lobulifera* Forel, 1909

Ectatomma opaciventre (Roger, 1861)
 = *concolor* Santschi, 1919

Ectatomma permagnum Forel, 1908
 = *strigosum* Emery, 1894

= *aerea* Forel, 1912
 = *confusa* Forel, 1909

Ectatomma ruidum (Roger, 1860)
 = *scabrosa* (Smith, 1862)
 = *aztecum* Emery, 1901

Ectatomma tuberculatum (Oliver, 1792)
 = *tridentata* (Fabricius, 1804)
 = *ferrugineus* Norton, 1868
 = *acrista* Forel, 1909

Genus *Gnampogenys* Roger, 1863

Gnampogenys caelata Kempf, 1967
 = *soror* Kempf & Brown, 1968

Gnampogenys concinna (Smith, 1858)
 = *conica* Borgmeier, 1929
 = *romani* (Wheeler W.M., 1923)
 = *semicircularis* Borgmeier, 1929

Gnampogenys continua Mayr, 1887
 = *panamensis* Santschi, 1931

Gnampogenys minuta (Emery, 1896)
 = *carinata* (Weber, 1940)
 = *emeryi* (Forel, 1901)
 = *mus* (Santschi, 1931)
 = *scabrosus* Mann, 1922

= *pneodonax* Kempf, 1968

Gnampogenys moelleri (Forel, 1912)
 = *splendens* (Santschi, 1929)

Gnampogenys mordax (Smith, 1858)
 = *nodosa* (Latrelle, 1802)
 = *purensis* Forel, 1912

= *sebastiani* Borgmeier, 1937

Gnampogenys pleurodon (Emery, 1896)

= *emeryi* (Santschi, 1929)

Gnampogenys regularis Mayr, 1870
 = *fiebrigii* Forel, 1909

= *splendidum* Pergande, 1896

Gnampogenys striatula Mayr, 1884
 = *antillana* (Santschi, 1929)
 = *angustiloba* (Forel, 1908)
 = *angustipleura* (Forel, 1908)
 = *brasiliensis* (Emery, 1902)

= *paulina* (Forel, 1908)

= *pernambucana* (Santschi, 1929)

Gnampogenys strigata (Norton, 1868)
 = *simplex* (Emery, 1896)

= *concentrica* Mayr, 1870

Gnampogenys sulcata (Smith, 1858)

= *bufonum* Weber, 1938
 = *cearensis* (Forel, 1912)
 = *nitens* Mann, 1916

= *ypirangensis* Borgmeier, 1928

Gnampogenys triangularis Mayr, 1887
 = *richteri* (Forel, 1913)

Tribe Typhlomyrmecini

Genus *Typhlomyrmex* Mayr, 1862

Typhlomyrmex clavicornis Emery, 1906
 = *divergens* Forel, 1906
 = *richardsi* Donisthorpe, 1939

- Typhlomyrmex pusillus* Emery, 1894
 = *schmidti* Menozzi, 1927
Typhlomyrmex rogenhoferi Mayr, 1862
 = *manco* Wheeler W.M., 1925
 = *robustus* Emery, 1890
 = *sculpturatus* Santschi, 1931

Subfamily Heteroponerinae

Tribe Heteroponerini

Genus *Acanthoponera* Mayr, 1862

- Acanthoponera minor* (Forel, 1899)
 = *schwarzi* Wheeler W.M., 1923 = *spininodis*
 Weber, 1939
 = *crassa* Brown, 1958
Acanthoponera mucronata (Roger, 1860)
 = *goyana* Borgmeier, 1939
 = *plaumannii* Donisthorpe, 1938
 = *wagneri* Santschi, 1921

Genus *Heteroponera* Mayr, 1887

- Heteroponera dolo* (Roger, 1860)
 = *aurea* (Forel, 1913)
Heteroponera inermis (Emery, 1894)
 = *schwebeli* Luederwalt, 1918

Subfamily Paraponerinae

Genus *Paraponera* Smith, 1858

- Paraponera clavata* (Fabricius, 1775)
 = *aculeata* (Oliver, 1792)
 = *tarsalis* (Perty, 1833)

Subfamily Proceratiinae

Tribe Proceratiini

Genus *Proceratium* Roger, 1863

- Proceratium micrommatum* (Roger, 1863)
 = *cavernicola* (Borgmeier, 1937)
 = *convexiceps* Borgmeier, 1957
Proceratium silaceum Roger, 1863
 = *crassicornis* Emery, 1895
 = *rugulosum* Wheeler W.M., 1915
 = *vestitum* Emery, 1895
 = *mancum* Mann, 1922

Tribe Probolomyrmecini

Genus *Probolomyrmex* Mayr, 1901

- Probolomyrmex boliviensis* Mann, 1923
 = *angusticeps* Smith M.R. 1949

Subfamily Myrmicinae

Tribe Attini

Genus *Acromyrmex* Mayr, 1865

- Acromyrmex ambiguus* Emery, 1888
 = *erectus* Santschi, 1925
Acromyrmex aspersus (Smith F., 1858)
 = *affinis* Santschi, 1925
 = *inquirens* (Forel, 1914)
 = *mesonotalis* (Emery, 1905c)
Acromyrmex coronatus (Fabricius, 1804)
 = *meinerti* (Fabricius, 1804)
 = *modesta* Forel, 1901
 = *moelleri* Forel, 1893
 = *ochraceolus* Santschi, 1925
 = *flavescens* Santschi, 1925
 = *medianus* Santschi, 1925
 = *obscurior* Santschi, 1925
 = *ornatus* Santschi, 1925
Acromyrmex crassispinus Forel, 1909
 = *atratus* Santschi, 1925
 = *diabolica* Santschi, 1922
 = *insularis* Santschi, 1925
 = *mediocris* Santschi, 1925
 = *rusticus* Santschi, 1925
 = *rufescens* Santschi, 1925
Acromyrmex fracticornis Forel, 1909
 = *jorgensenni* Forel, 1913
Acromyrmex heyeri Forel, 1899
 = *gaudens* Santschi, 1925
 = *lillensis* Santschi, 1925
Acromyrmex hystrix Latreille, 1802
 = *emili* Forel, 1904
Acromyrmex laticeps Emery, 1905
 = *hortulanus* Santschi, 1925
Acromyrmex landolti Forel, 1885
 = *senex* Santschi, 1925
 = *cloosae* (Forel, 1912)
 = *planorum* Weber 1937
Acromyrmex lundii (Guérin-Meneville, 1838)
 = *bonariensis* Emery, 1905
 = *dubia* Forel, 1908
 = *risii* Forel, 1908
Acromyrmex niger (Smith F., 1858)
 = *muticinoda* Forel, 1901
 = *homalops* Emery, 1905
Acromyrmex nobilis Santschi, 1939
 = *guentheri* Forel, 1893
 = *pallida* Crawley, 1921
Acromyrmex rugosus (Smith F., 1858)
 = *bigener* Santschi, 1925
 = *vestitus* Santschi, 1925
 = *pallida* (Smith F., 1858)

- Acromyrmex silvestrii* Emery, 1905
 = *mesopotamicus* Gallardo, 1916
Acromyrmex striatus (Roger, 1863)
 = *laeviventris* Santschi, 1925
Acromyrmex subterraneus Forel, 1893
 = *mixtus* Santschi, 1925
 = *purensis* Forel, 1912

Genus *Apterostigma* Mayr, 1865

- Apterostigma auriculatum* Lattke, 1997
 = *demerarae* Wheeler W.M., 1925
 = *immobile* Weber, 1940
Apterostigma dentigerum Wheeler W.M., 1925
 = *angulatum* Weber, 1938
 = *dubium* Weber, 1938
Apterostigma ierense Weber, 1937
 = *fitzgeraldi* Weber, 1937
Apterostigma robustum Emery, 1896
 = *amiae* Weber, 1937
 = *branneri* Mann, 1916
 = *jubatum* Wheeler W.M., 1925
 = *tic* Weber, 1938
Apterostigma urichii Forel, 1893
 = *guianense* Weber, 1937
 = *nitidum* Weber, 1937

Genus *Atta* Fabricius, 1804

- Atta cephalotes* (Linnaeus, 1758)
 = *migratoria* (De Geer, 1773)
 = *grossa* (Fabricius, 1787)
 = *fervens* (Drury, 1782)
 = *visitatrix* (Christ, 1791)
 = *integrior* Forel, 194
 = *isthmicola* Weber, 1941
 = *lutea* Forel, 1893
 = *oaxaquensis* Gonçalves, 1942
 = *opaca* Forel, 1904
 = *polita* Emery, 1905
Atta columbica Guérin-Méneville, 1845
 = *erecta* Santschi, 1929
 = *tonspipes* Santschi, 1929
 = *lebasi* Guérin-Meneville, 1844
Atta laevigata (Smith F., 1858)
 = *bolchevista* Gonçalves, 1942
 = *venezuelensis* Gonçalves, 1942
Atta sexdens (Linnaeus, 1758)
 = *sexdentata* (Latreille, 182)
 = *abdominalis* (Smith, F. 1858)
 = *coptophylla* Guérin-Meneville, 1844
 = *flavicornis* (Fabricius, 1789)
 = *autuorii* Borgmeier, 1950
 = *fuscata* Santschi, 1922
 = *piriventris* Santschi, 1919
 = *rubropilosa* Forel, 1908

- Atta vollenweideri* Forel, 1893
 = *tristis* Santschi, 1919

Genus *Cyphomyrmex* Mayr, 1862

- Cyphomyrmex costatus* Mann, 1922
 = *colombianus* Weber, 1940
Cyphomyrmex faunulus Wheeler W.M., 1925
 = *petiolatus* Weber, 1938
Cyphomyrmex flavidus Pergande, 1896
 = *dentatus* Forel, 1901
Cyphomyrmex hamulatus Weber, 1938
 = *trinitatis* Weber, 1938
Cyphomyrmex minutus Mayr, 1862
 = *steinheli* Forel, 1885
 = *comalensis* Wheeler W.M., 1907
 = *arnoldi* Aguayo, 1932
 = *atrata* Forel, 1912
 = *breviscapus* Weber, 1940
 = *flavesiensis* Weber, 1940
 = *venezuelensis* Weber, 1938
Cyphomyrmex olitor Forel, 1893
 = *quebradae* Kusnezov, 1949
Cyphomyrmex rimosus (Spinola, 1851)
 = *difformis* (Smith F., 1858)
 = *cochunae* Kusnezov, 1949
 = *fuscus* Emery, 1894
Cyphomyrmex salvini Forel, 1899
 = *acutus* Weber, 1940
 = *championi* Forel, 1899
Cyphomyrmex transversus Emery, 1894
 = *olindanus* Forel, 1901
 = *pencosensis* Forel, 1914

Genus *Mycetarotes* Emery, 1913

- Mycetarotes parallelus* Emery, 1906
 = *luederwaldti* (Forel, 1911)

Genus *Mycetophylax* Emery, 1913

- Mycetophylax conformis* Mayr, 1884
 = *brittoni* (Wheeler W.M., 1907)
 = *litoralis* Weber, 1937
Mycetophylax emeryi Forel, 1907
 = *hummelincki* Weber, 1948

Genus *Mycocepurus* Forel, 1893

- Mycocepurus goeldii* Forel, 1893
 = *gentilis* Santschi, 1925
 = *oglobini* Santschi, 1933
 = *oglobini* (Kusnezov, 1951)
 = *schuppi* Forel, 1901
Mycocepurus smithii Forel, 1893
 = *attaxenus* (Menozzi, 1936)
 = *bolivianus* Weber, 1938
 = *borinquenensis* Wheeler W.M., 1907

= *eucarnitae* Forel, 1913
 = *manni* Weber, 1938
 = *reconditus* Borgmeier, 1937
 = *tolteca* Wheeler W.M., 1907
 = *trinidadensis* Weber, 1937

Genus *Myrmicocrypta* Smith, 1860

Myrmicocrypta collaris Emery, 1913
 = *corniculata* Emery, 1913

Genus *Sericomyrmex* Mayr, 1865

Sericomyrmex amabilis Wheeler W.M., 1925
 = *bierigi* Santschi, 1931

Genus *Trachymyrmex* Forel, 1893

Trachymyrmex bugnioni Forel, 1912
 = *abortivus* Wheeler W.M., 1925
 = *cucumis* Mann, 1922
 = *humilis* Wheeler W.M., 1916
 = *morgani* Weber, 1940
 = *shomburgki* Weber, 1938
 = *tucuche* Weber, 1938
Trachymyrmex cornetzi Forel, 1912
 = *annulatus* Santschi, 1936
 = *bivittatus* Wheeler W.M. 1922
 = *brevispinosus* Weber, 1945
 = *gatun* Weber, 1940
 = *naranjo* Forel, 1912
 = *uncifer* Santschi, 1931
Trachymyrmex diversus Mann, 1916
 = *phippsi* Weber, 1937
Trachymyrmex farinosus Emery, 1894
 = *trifurcatus* Weber, 1938
Trachymyrmex uruchi (Forel 1893)
 = *fusca* (Emery, 1894)
 = *marthae* Forel, 1912
 = *panamensis* Wheeler W.M., 1925
 = *radicis* Weber, 1938

Tribe Basicerotini

Genus *Basiceros* Schulz, 1906

Basiceros convexiceps Mayr, 1887
 = *squamifer* Borgmeier, 1937

Genus *Eurhopalothrix* Brown & Kempf, 1961

Eurhopalothrix bolaui (Mayr, 1870)
 = *amoena* (Mann, 1922)
Eurhopalothrix gravis Mann, 1922
 = *reichenspergeri* Santschi, 1923
 = *schmidti* Menozzi, 1936

Genus *Octostruma* Forel, 1912

Octostruma balzani Emery, 1894
 = *barberi* (Mann, 1922)

= *equilatera* (Weber, 1934)
 = *lutzi* (Wheeler W.M., 1913)
Octostruma iheringi Emery, 1888
 = *godmani* (Forel, 1899)
 = *simoni* (Emery, 1890)
 = *spei* (Forel, 1912)
 = *wighti* (Wheeler W.M., 1908)

Tribe Blepharidattini

Genus *Wasemannia* Forel, 1893

Wasemannia auropunctata Roger, 1863
 = *atomun* (Santschi, 1914)
 = *glabra* Santschi, 1931
 = *panamana* (Enzmann, 1947)
Wasemannia sulcaticeps Emery, 1894
 = *bruchi* Santschi, 1919

Tribe Cephalotini

Genus *Cephalotes* Latreille, 1802

Cephalotes atratus (Linnaeus, 1758)
 = *quadridens* (De Geer, 1773)
 = *dibiatus* (Smith F., 1854)
 = *rufiventris* (Emery, 1894)
 = *nitidiventris* Santschi, 1920
 = *crassispina* Santschi, 1920
 = *erectus* Santschi, 1920
Cephalotes basalis (Smith, 1876)
 = *multispinus* (Smith, F. 1876)
Cephalotes bimaculatus (Smith F., 1860)
 = *aztecus* (Smith, 1860)
Cephalotes bohlsi (Emery, 1896)
 = *medusa* (Santschi, 1919)
Cephalotes bruchi (Forel, 1912)
 = *ridiculus* (Santschi, 1915)
 = *pampensis* (Santschi, 1915)
Cephalotes complanatus (Guérin-Méneville, 1844)
 = *angulatus* (Smith F., 1858)
Cephalotes conspersus (Smith F., 1867)
 = *variegata* (Forel, 1911)
Cephalotes cordatus (Smith F., 1853)
 = *boliviensis* (Santschi, 1921)
Cephalotes depressus (Klug, 1824)
 = *sorocabensis* (Forel, 192)
Cephalotes femoralis (Smith F., 1853)
 = *silvae* (Forel, 1906)
Cephalotes grandinosus (Smith F., 1860)
 = *nevadensis* (Forel, 1912)
Cephalotes hamulus (Roger, 1863)
 = *haytianus* (Forel, 1901)
 = *affinis* (Wheeler, 1936)
Cephalotes jheringi (Emery, 1894)
 = *peltatus* (Emery, 1896)
 = *ellenriederi* (Forel, 1911)

- Cephalotes liogaster* (Santschi, 1916)
 = *lobigaster* (Kempf, 1958)
- Cephalotes maculatus* (Smith F., 1876)
 = *magdalenensis* (Forel, 1899)
 = *nanus* (Forel, 1912)
- Cephalotes marginatus* (Fabricius, 1804)
 = *descenpinosus* (Santschi, 192)
- Cephalotes membranaceus* (Klug, 1824)
 = *fervidus* (Smith, 1876)
- Cephalotes minutus* (Fabricius, 1804)
 = *quadrimaculatus* (Klug, 1824)
 = *caustica* (Kollar, 1832)
 = *cognatus* (Smith, 1867)
 = *exiguus* (Smith, 1867)
 = *volexemi* (Emery, 1878)
- Cephalotes multispinosus* (Norton, 1868)
 = *gibbosus* (Smith, 1876)
- Cephalotes notatus* (Mayr, 1866)
 = *fleddermannii* (Kempf, 1958)
- Cephalotes oculatus* (Spinola, 1853)
 = *aethiops* (Smith, 1854)
- Cephalotes opacus* Santschi, 1920
 = *abdominalis* Santschi, 1929
- Cephalotes pallens* (Klug, 1824)
 = *discocephalus* (Smith, 1854)
 = *araneolus* (Smith, 1854)
- Cephalotes pallidicephalus* (Smith F., 1876)
 = *striativentris* (Emery, 1894)
- Cephalotes placidus* (Smith F., 1860)
 = *fenestralis* (Smith F., 1867)
- Cephalotes pusillus* (Klug, 1824)
 = *elongatus* (Klug, 1824)
 = *obtusus* (Smith, 1858)
 = *brevispinosa* (Santschi, 1921)
- Cephalotes quadratus* (Mayr, 1868)
 = *convexus*
- Cephalotes scutulatus* (Smith F., 1867)
 = *angulosus* (Smith, 1876)
 = *jucundus* (Smith, 1876)
- Cephalotes setulifer* (Emery, 1894)
 = *orbis* (Forel, 1906)
- Cephalotes spinosus* (Mayr, 1862)
 = *punctatus* (Mayr, 1862)
 = *peruvianus* (Forel, 1911)
- Cephalotes squamosus* (Vierbergen y Scheven, 1995)*
 = *pseudoaztecus* (Vierbergen y Scheven, 1995)*
- Cephalotes umbraculatus* (Fabricius, 1804)
 = *elegans* (Smith, 1854)
 = *flavomaculatus* (Mayr, 1862)
- Cephalotes unimaculatus* (Smith F., 1854)
 = *steinheili* (Forel, 198)
 = *signatus* (Wheeler W.M., 1936)

Genus *Procryptocerus* Emery, 1887

- Procryptocerus adlerzi* (Mayr, 1887)
 = *latians* Forel, 1912

- Procryptocerus attenuatus* (Smith F., 1876)
 = *puncticeps* (Smith F., 1876)
 = *guianensis* Weber, 1938
- Procryptocerus batesi* Forel, 1899
 = *laeviventris* Forel, 1899
- Procryptocerus mayri* Forel, 1899
 = *reichenspergeri* Santschi, 1921

Tribe Crematogastrini

Genus *Crematogaster* Lund, 1831

- Crematogaster abstinentis* Forel, 1899
 = *pygmeaea* Forel, 1904
 = *suturalis* Forel, 1912
 = *ornatipilis* Wheeler W.M., 1918
 = *erici* Santschi, 1929
 = *chacoana* Santschi, 1933
- Crematogaster acuta* (Fabricius, 1804)
 = *quadriceps* Smith, 1858
 = *centralis* Santschi, 1932
- Crematogaster arcuata* Forel, 1899
 = *aruga* Forel, 1913
- Crematogaster brasiliensis* Mayr, 1878
 = *armandi* Forel, 1921
 = *ludio* Forel, 1912
 = *inca* Wheeler W.M., 1925
 = *cocciphila* Borgmeier, 1934
- Crematogaster carinata* Mayr, 1862
 = *parabiotaica* Forel, 1904
- Crematogaster corticicola* Mayr, 1887
 = *rugiceps* Forel, 1911
- Crematogaster crinosa* Mayr, 1862
 = *minutior* Forel, 1893
 = *schuppi* Forel, 1901
 = *striatinoda* Forel, 1912
 = *sampaioi* Forel, 1912
 = *towsendi* Wheeler W.M., 1925
 = *chathamensis* Wheeler W.M., 1933
- Crematogaster cristata* Santschi, 1929
 = *terricola* Borgmeier, 1929
- Crematogaster curvispinosa* Mayr, 1862
 = *antillana* Forel, 1893
 = *accola* Wheeler W.M., 1934
 = *sculpturata* Pergande, 1896
 = *kemali* Santschi, 1923
 = *fuliginea* Santschi, 1925
 = *phytoeca* Wheeler W.M., 1934
 = *panamana* Wheeler W.M., 1942
 = *obscura* Santschi, 1929
- Crematogaster distans* Mayr, 1870
 = *paraensis* Forel, 1904
 = *parviceps* Forel, 1908
 = *pevsnerae* Forel, 1912
 = *vanda* Borgmeier, 1929

- = *descolei* Kusnezov, 1949
 = *ursus* Mayr, 1866
Crematogaster erecta Mayr, 1866
 = *virgula* Forel, 1899
 = *projecta* Santschi, 1925
Crematogaster evallans Forel, 1907
 = *carbonescens* Forel, 1913
Crematogaster limata Smith, 1858
 = *palans* Forel, 1912
 = *ascendens* Wheeler W.M., 1925
 = *dextella* Santschi, 1929
Crematogaster magnifica Santschi, 1925
 = *nociva* Borgmeier, 1937
Crematogaster montezumia Smith, 1858
 = *sulcata* Mayr, 1870
 = *ramulinida* Forel, 1899
 = *functa* Forel, 1911
 = *cristulata* Santschi, 1925
 = *proletaria* Santschi, 1933
Crematogaster obscurata Emery, 1895
 = *agnita* Wheeler W.M., 1934
Crematogaster stollii Forel, 1885
 = *amazonensis* Forel, 1905
 = *autrini* Mann, 1916
 = *guianensis* Crawley, 1916
 = *parvispina* Wheeler W.M., 1922
 = *hyperphyes* Kusnezov, 1953
Crematogaster sumichrasti Mayr, 1870
 = *surdior* Forel, 1885
 = *titlanica* Wheeler W.M., 1936
 = *maya* Wheeler W.M., 1936
Crematogaster torosa Mayr, 1870
 = *tumulifera* Forel, 1899
 = *arizonensis* Wheeler W.M., 1908
Crematogaster vicina André, 1893
 = *wighti* Wheeler W.M., 1908

Tribe Dacetini

Genus *Pyramica* Roger, 1862

- Pyramica alberti* (Forel, 1893)
 = *intermedia* (Wheeler W.M., 1913)
 = *guianensis* (Weber, 1934)
Pyramica brevicornis (Mann, 1922)
 = *mustelina* (Weber, 1934)
Pyramica emiliae (Forel, 1907)
 = *alegrensis* (Brown, 1953)
Pyramica epinotalis (Weber, 1934)
 = *skwarrae* (Wheeler W.M., 1934)
Pyramica gundlachi Roger, 1862
 = *banillensis* (Santschi, 1930)
 = *vincentensis* (Forel, 1893)
 = *bierigi* (Santschi, 1930)
 = *infuscata* (Weber, 1934)
 = *berlesei* (Weber, 1934)

- = *isthmica* (Santschi, 1931)
Pyramica nigrescens (Wheeler W.M., 1911)
 = *nana* (Santschi, 1930)
 = *longipilis* (Weber, 1934)
Pyramica schulzi (Emery, 1894)
 = *studiosi* (Weber, 1934)
Pyramica subdentata (Mayr, 1887)
 = *tristani* (Menozzi, 1931)
 = *clavata* (Weber, 1934)

Genus *Strumigenys* Smith, 1860

- Strumigenys biolleyi* Forel, 1908
 = *tridens* Weber, 1934
 = *luctuosa* Menozzi, 1936
Strumigenys cordovensis Mayr, 1887
 = *mokensis* Forel, 1905
Strumigenys elongata Roger, 1863
 = *imitator* Mayr, 1887
 = *nicaraguensis* Weber, 1934
Strumigenys emmae (Emery, 1890)
 = *clypeatus* Szabó, 1909
 = *malesiana* (Forel, 1913)
 = *wheeleri* (Donisthorpe, 1916)
Strumigenys godmani Forel, 1899
 = *ferox* Weber, 1934
Strumigenys louisianae Roger, 1863
 = *unidentata* Mayr, 1887
 = *longicornis* Emery, 1894
 = *fusca* Emery, 1894
 = *obscuriventris* Wheeler W.M., 1908
 = *cubaensis* Mann, 1920
 = *laticephala* Smith M.R., 1931
 = *guatemalensis* Weber, 1934
 = *costaricensis* Weber, 1934
 = *clasmosporgia* Brown, 1953
 = *producta* Brown, 1953
Strumigenys mandibularis Smith, 1860
 = *batesi* Forel, 1911
Strumigenys saliens Mayr, 1887
 = *procera* Emery, 1894
 = *angusticeps* Forel, 1912
Strumigenys smithii Forel, 1886
 = *inaequalis* Emery, 1890
 = *truncata* (Forel, 1912)

Tribe Formicoxenini

Genus *Cardiocondyla* Emery, 1869

- Cardiocondyla emeryi* Forel, 1881
 = *nereis* Wheeler W.M., 1927
 = *monilicornis* (Emery, 1917)

Genus *Nesomyrmex* Wheeler W.M., 1910

- Nesomyrmex asper* (Mayr, 1887)
 = *rufa* (Emery, 1896)
 = *sulfurea* Forel, 1912

- Nesomyrmex echinatinodis* Forel, 1886
 = *aculeatinodis* (Emery, 1896)
 = *cordincola* (Wheeler W.M., 1942)
 = *dalmasi* (Forel, 1899)
 = *pungentinodis* (Emery, 1896)
 = *schmidti* (Menozzi, 1927)
- Nesomyrmex pulcher* Emery, 1917
 = *formosus* (Emery, 1915) (Hom)
- Nesomyrmex sculptiventris* Mayr, 1887
 = *borgmeieri* (Wheeler W.M., 1931)
 = *distincta* (Crawley, 1922)
 = *major* (Forel, 1899)
- Nesomyrmex spininodis* Mayr, 1887
 = *genualia* (Santschi, 1922)
 = *umbratilis* (Wheeler W.M., 1921)

Genus *Temnothorax* Mayr, 1861

- Temnothorax rugatulus* Emery, 1895
 = *anectens* (Wheeler W.M., 1931)
 = *cockerelli* (Wheeler W.M., 1917)
 = *mediorufus* (Wheeler W.M., 1917)

Genus *Ochetomyrmex* Mayr, 1878

- Ochetomyrmex semipolitus* Mayr, 1878
 = *mayri* Forel, 198
 = *subpolita* (Wheeler W.M. 1916)
 = *argentinus* (Kusnezov, 1957)
 = *bolivianus* (Kusnezov, 1962)

Genus *Xenomyrmex* Forel, 1885

- Xenomyrmex floridanus* Emery, 1895
 = *rufescens* Wheeler W.M., 1931
 = *cubanus* Wheeler W.M., 1931
 = *lucayanus* Wheeler W.M., 1931
- Xenomyrmex stollii* Forel, 1885
 = *mexicanus* Wheeler W.M., 1931
 = *nodosus* Creighton, 1953

Tribe Myrmecinini

Genus *Myrmecina* Curtis, 1829

- Myrmecina americana* Emery, 1895
 = *brevispinosa* Creighton, 1950
 = *quadrispina* Enzmann, 1946
 = *texana* Wheeler W.M., 1908
 = *californica* Smith M.R., 1948

Tribe Myrmicini

Genus *Hylomyrma* Forel, 1912

- Hylomyrma balzani* (Emery, 1894)
 = *speciosa* (Borgmeier, 1937)
- Hylomyrma reitteri* (Mayr, 1887)
 = *goeldii* (Forel, 1912)

- Genus *Pogonomyrmex* Mayr, 1868
Pogonomyrmex barbatus Smith, 1858
 = *molefaciens* (Buckley, 1860)
 = *nigrescens* Wheeler W.M., 1902
- Pogonomyrmex brevibarbis* Emery, 1906
 = *ater* Donisthorpe, 1933
 = *carettei* Santschi, 1922
- Pogonomyrmex californicus* Buckley, 1867
 = *estebanius* Pergande, 1894
 = *longinodis* Emery, 1895
 = *nitritatus* Cook, 1953
- Pogonomyrmex coarctatus* Mayr, 1868
 = *striaticeps* Emery, 1906
- Pogonomyrmex cunicularius* Mayr, 1887
 = *brevispinosus* (Smith F., 1858)
 = *carnivora* Santschi, 1925
 = *pencosensis* Forel, 1914
 = *serpens* Santschi, 1922
- Pogonomyrmex desertorum* Wheeler W.M., 192
 = *ferrugineus* Olsen, 1934
- Pogonomyrmex imberbiculus* Wheeler W.M., 1902
 = *townsendi* Wheeler W.M., 1909
- Pogonomyrmex maricopa* Wheeler W.M., 1914
 = *barnesi* Smith M.R., 1929
 = *sinaloanus* Olsen, 1934
- Pogonomyrmex rastratus* Mayr, 1868
 = *neuquensis* Santschi, 1922
 = *weiseri* Santschi, 1922
- Pogonomyrmex rugosus* Emery, 1895
 = *similis* Olsen, 1934
 = *fuscatus* Emery, 1895
- Pogonomyrmex tenuispinus* Forel, 1914
 = *dentatus* Olsen, 1934
- Pogonomyrmex vermiculatus* Emery, 1906
 = *intermedia* Menozzi, 1935
 = *semistriata* Emery, 1906
 = *spinolae* Emery, 1906

Tribe Pheidolini

Genus *Aphaenogaster* Mayr, 1853

- Aphaenogaster ensifera* Forel, 1899
 = *manni* (Wheeler W.M. & Creighton, 1934)

Genus *Messor* Forel, 1890

- Messor andrei* Mayr, 1886
 = *castaneus* (Wheeler W.M. & Creighton, 1934)

Genus *Pheidole* Westwood, 1839

- Pheidole aberrans* Mayr, 1868
 = *fartilia* Forel, 1913
 = *mutica* Emery, 1906
- Pheidole alacris* Santschi, 1923
 = *mediorubra* Santschi, 1933

- Pheidole alfaroi* Emery, 1896
 = *innupta* Menozzi, 1931
- Pheidole angusta* Forel, 1908
 = *idulis* Forel, 1912
- Pheidole arhuaca* Forel, 1901
 = *bimons* Forel, 1912
- Pheidole bambusarum* Forel, 1908
 = *ogloblini* Kusnezov, 1952
- Pheidole biconstricta* Mayr, 1870
 = *lallemandi* Forel, 1901
 = *holmgreni* Wheeler W.M., 1925
 = *festata* Wheeler W.M., 1925
- Pheidole bilimeki* Mayr, 1870
 = *anastasii* Emery, 1896
 = *deplanata* Pergande, 1896
 = *antoniensis* Forel, 1901
 = *anectans* Wheeler W.M., 1905
 = *jamaicensis* Wheeler W.M., 1908
- Pheidole bolleyi* Forel, 1908
 = *tristani* Forel, 1908
- Pheidole brevicoma* Mayr, 1887
 = *kraepelini* Forel, 1901
- Pheidole cameroni* Mayr, 1887
 = *monticola* Cameron, 1892
- Pheidole fabricator* (Smith, 1858)
 = *nigriventris* (Smith, 1858)
- Pheidole gauthieri* Forel, 1901
 = *gauthieri oxymora* Forel, 1912
- Pheidole gertrudae* Forel, 1886
 = *gertrudae leonhardi* Forel, 1901
 = *gertrudae loretensis* Santschi, 1933
 = *humilis* Borgmeier, 1930
- Pheidole gibba* Mayr, 1887
 = *lopesi* Borgmeier, 1937
- Pheidole hecate* Wheeler W.M., 1911
 = *hecate malevola* Wheeler W.M., 1911
 = *hecate bruesi* Wheeler W.M., 1917
- Pheidole heyperi* Forel, 1899
 = *ultrix* Forel, 1912
- Pheidole hirtula* Forel, 1899
 = *acolhua* Wheeler W.M., 1914
- Pheidole megacephala* (Fabricius, 1793)
 = *trinodis* (Losana, 1834)
 = *testacea* (Smith, 1858)
- Pheidole moerens* Wheeler W.M., 1908
 = *dominicensis* Wheeler W.M., 1913
 = *creola* Wheeler W.M. & Mann, 1914
- Pheidole nigella* Emery, 1894
 = *dimidiata* Emery, 1894
- Pheidole nitidula* Emery, 1888
 = *strobeli* Emery, 1906
 = *perversa* Forel, 1908
 = *misera* Santschi, 1916
 = *silvicola* Borgmeier, 1927
- Pheidole obtusospinosa* Pergande, 1896
 = *arizonica* Santschi, 1911
 = *subdentata* Pergande, 1896
- Pheidole oxyops* Forel, 1908
 = *regia* Forel, 1908
- Pheidole peregrina* Wheeler W.M., 1916
 = *inpariceps* Santschi, 1923
 = *fusciclava* Santschi, 1923
- Pheidole perpusilla* Emery, 1894
 = *decem* Forel, 1901
- Pheidole polita* Emery, 1894
 = *wheeleri* Mann, 1916
- Pheidole pubiventris* Mayr, 1887
 = *cearensis* Forel, 1901
 = *nevadensis* Forel, 1901
 = *timmii* Forel, 1901
 = *foederalis* Borgmeier, 1928
- Pheidole punctatissima* Mayr, 1870
 = *napaea* Wheeler W.M., 1934
- Pheidole radoszkowskii* Mayr, 1884
 = *acuta* Emery, 1894
 = *luteola* Forel, 1893
 = *parvinoda* Forel, 1912
 = *paranana* Santschi, 1925
 = *reflexans* Santschi, 1933
 = *australis* Emery, 1890
- Pheidole rudigenis* Emery, 1906
 = *bucolica* Santschi, 1929
- Pheidole rufipilis* Forel, 1908
 = *divexa* Forel, 1908
 = *impia* Santschi, 1923
- Pheidole schmalzi* Emery, 1894
 = *rehi* Forel, 1900
- Pheidole sospes* Forel, 1908
 = *minutissima* Kusnezov, 1952
- Pheidole subarmata* Mayr, 1884
 = *borinquensis* Wheeler W.M., 1908
 = *cornuta* Emery, 1890
 = *dentimentum* Santschi, 1929
 = *elongatula* Forel, 1893
 = *hondurensis* Mann, 1922
 = *imbecillis* Emery, 1906
 = *nassavensis* Wheeler W.M., 1905
 = *nefasta* Santschi, 1929
- Pheidole tepicana* Pergande, 1896
 = *carbonaria* Pergande, 1896
 = *rugifrons* Pergande, 1896
- Pheidole terresi* Wheeler W.M. & Mann, 1914
 = *illota* Wheeler W.M. & Mann, 1914
- Pheidole transversostriata* Mayr, 1887
 = *nigridens* Forel, 1901
 = *lacerta* Wheeler W.M., 1922
- Pheidole triconstricta* Emery, 1906
 = *ambulans* Emery, 1906
 = *discursans* Forel, 1912

- = *rosarensis* Forel, 1913
- = *saviozae* Forel, 1911
- = *hebe* Santschi, 1923
- Pheidole tristis* (Smith, 1858)
 - = *emeryi* Mayr, 1887
 - = *fumipennis* (Smith, 1858)
 - = *rubra* (Smith, 1858)
- Pheidole ursus* Mayr, 1870
 - = *cressoni* (André, 1887)
 - = *gracilinoda* Forel, 1904
- Pheidole vafra* Santschi, 1923
 - = *idiota* Santschi, 1923
- Pheidole williamsi* Wheeler W.M., 1919
 - = *seymourensis* Wheeler W.M., 1924

Tribe Solenopsidini

Genus *Megalomyrmex* Forel, 1885

- Megalomyrmex balzani* Emery, 1894
 - = *duckei* Forel, 1912
- Megalomyrmex silvestrii* Wheeler W.M., 1909
 - = *brasiliensis* Borgmeier, 1930
 - = *silvestrii* Wheeler W.M., 1925
 - = *sjostedti* Wheeler W.M., 1925
 - = *wheeleri* Weber, 1940
- Megalomyrmex staudingeri* Emery, 1890
 - = *nasutus* Forel, 1912

Genus *Monomorium* Mayr, 1855

- Monomorium cyaneum* Wheeler W.M., 1914
 - = *emersoni* Gregg, 1945
- Monomorium floricola* (Jerdon, 1851)
 - = *poeicum* Roger, 1863
 - = *floreanum* Stitz, 1932
- Monomorium minimum* (Buckley, 1867)
 - = *atra* (Buckley, 1867)
 - = *metoecus* Brown & Wilson, 1957
- Monomorium monomorium* Bolton, 1987
 - = *minutum* Mayr, 1855

Genus *Oxyepoecus* Santschi, 1926

- Oxyepoecus bruchi* Santschi, 1926
 - = *minuta* (Kusnezov, 1952)
- Oxyepoecus inquilinus* (Kusnezov, 1952)
 - = *turgidus* Kempf, 1969
- Oxyepoecus rastratus* (Mayr, 1887)
 - = *luederwaldti* (Forel, 1913)

Genus *Tranopelta* Mayr, 1866

- Tranopelta gilva* Mayr, 1866
 - = *amblyops* (Emery, 1894)
 - = *heyperi* (Forel, 1901)
 - = *gilva brunnea* Forel, 1909
 - = *heyperi columbica* Forel, 1912
 - = *gilva albida* Mann, 1916

Genus *Solenopsis* Westwood, 1840

- Solenopsis aurea* Wheeler W.M., 1906
 - = *huachucana* Wheeler W.M., 1915
- Solenopsis bruchiella* Emery, 1922
 - = *bruchi* Forel, 1918
- Solenopsis daguerrei* (Santschi, 1930)
 - = *acuminata* (Borgmeier, 1949)
- Solenopsis gayi* (Spinola, 1851)
 - = *fazi* Santschi, 1923
- Solenopsis geminata* (Fabricius, 1804)
 - = *clypeata* (Smith, 1858)
 - = *mandibularis* Westwood, 1840
 - = *paleata* (Lund, 1831)
 - = *cephalotes* Smith, 1859
 - = *glaber* (Smith, 1862)
 - = *laboriosus* (Smith, 1860)
 - = *polita* (Smith, 1862)
 - = *drewseni* (Mayr, 1861)
 - = *lincecumii* (Buckley, 1867)
 - = *saxicola* (Buckley, 1867)
 - = *nigra* Forel, 1908
- Solenopsis germaini* Emery, 1895
 - = *schedingi* Forel, 1907
- Solenopsis latastei* Emery, 1895
 - = *hoffmanni* Forel, 1912
- Solenopsis molesta* (Say, 1836)
 - = *debilis* Mayr, 1886
 - = *exigua* (Buckley, 1867)
 - = *minuta* (Say, 1836)
- Solenopsis patagonica* Emery, 1906
 - = *thoracica* Santschi, 1923
- Solenopsis pygmaea* Forel, 1901
 - = *exigua* Forel, 1893
- Solenopsis quinquecuspis* Forel, 1913
 - = *blumi* Buren, 1972
- Solenopsis richteri* Forel, 1909
 - = *tricuspidis* Forel, 1912
 - = *oblongiceps* Santschi, 1936
- Solenopsis succinea* Emery, 1890
 - = *inermiceps* Wheeler W.M. & Mann, 1914
- Solenopsis xyloni* McCook, 1879
 - = *maniosa* Wheeler W.M., 1915
 - = *pylades* Forel, 1904

Tribe Stenammini

Genus *Rogeria* Emery, 1894

- Rogeria blanda* (Smith, 1858)
 - = *foveata* Kempf, 1964
- Rogeria procera* Emery, 1896
 - = *brasiliensis* (Borgmeier, 1955)

Genus *Stenamma* Westwood, 1839

- Stenamma diecki* Emery, 1895
 - = *impressum* Buren, 1944

Tribe Tetramoriini**Genus *Tetramorium* Mayr, 1855***Tetramorium caespitum* (L., 1758)= *immigrans* Santschi, 1927*Tetramorium spinosum* (Pergande, 1896)= *insons* (Wheeler W.M., 1915)= *wheeleri* (Forel, 1901)**Appendix 3 / Anexo 3**

Subfamilies, tribes and genera of ants of the World, following the classification proposed by Bolton, 2003. (*): Extinct taxons.

Subfamilias, tribus y géneros de hormigas del mundo, basada en la clasificación propuesta por Bolton, 2003. ()*: taxones extintos.**Subfamily Aneuretinae****Tribe Aneuretini Emery, 1913***Aneuretellus** Dlussky, 1988*Aneuretus* Emery, 1893*Mianeuretus** Carpenter, 1930*Paraneuretus** Wheeler W.M., 1915*Protaneuretus** Wheeler W.M., 1915*Aneuretinae Incertae sedis:**Burmomyrma** Dlussky, 1996**Subfamily Dolichoderinae****Tribe Dolichoderini Forel, 1878***Alloiomma** Zhang, 1989*Amymex* Kusnezov, 1953*Anillidris* Santschi, 1936*Anonychomyrma* Donisthorpe, 1947*Asymphylomyrmex** Wheeler W.M., 1915*Axinidris* Weber, 1941*Azteca* Forel, 1878*Bothriomyrmex* Emery, 1869*Ctenobethylus** Brues, 1939*Doleromyrma* Forel, 1907*Dolichoderus* Lund, 1831*Dorymyrmex* Mayr, 1866*Ectophyllum* Forel, 1909*Elaeomyrmex** Carpenter, 1930*Elaphroditus** Zhang, 1989*Emplastus** Donisthorpe, 1920*Eotapinoma** Dlussky, 1988*Eurymyrmex** Zhang, Sun & Zhang, 1994*Forelius* Emery, 1888*Froggattella* Forel, 1902*Iridomyrmex* Mayr, 1862*Kotshkorkia** Dlussky, 1981*Leptomyrmex* Mayr, 1862*Leptomyrmula** Emery, 1913*Linepithema* Mayr, 1866*Liometopum* Mayr, 1861*Loweriella* Shattuck, 1992*Miomyrmex* Carpenter, 1930*Ochetellus* Shattuck, 1992*Papyrius* Shattuck, 1992*Petraeomyrmex** Carpenter, 1930*Philidris* Shattuck, 1992*Pityomyrmex** Wheeler W.M., 1915*Protazteca* Carpenter, 1930*Tapinoma* Foerster, 1850*Technomyrmex* Mayr, 1872*Turneria* Forel, 1895*Zerichinius** Dlussky, 1988**Subfamily Formicinae****Tribe Lasiini Ashmead, 1905***Acanthomyops* Mayr, 1862*Acropyga* Roger, 1862*Anoplolepis* Santschi, 1914*Cladomyrma* Wheeler W.M., 1920*Lasiophanes* Emery, 1895*Lasius* Fabricius, 1804*Myrmecocystus* Wesmael, 1838*Prolasius* Forel, 1892*Stigmacros* Forel, 1905*Teratomyrmex* McAreavey, 1957**Tribe Plagiolepidini Forel, 1886***Agraulomyrmex* Prins, 1932*Aphomomyrmex* Emery, 1899*Brachymyrmex* Mayr, 1868

- Bregmatomyrma* Wheeler W.M., 1929
Euprenolepis Emery, 1906
Lepisiota Sanstchi, 1926
Myrmelachista Roger, 1863
Paratrechina Motschoulsky, 1863
Petalomyrmex Snelling, 1979
Plagiolepis Mayr, 1861
Prenolepis Mayr, 1861
Pseudaphomomyrmex Wheeler W.M., 1920
Pseudolasius Emery, 1887
Tapinolepis Emery, 1925

Tribe Myrmoteratini Emery, 1895

- Myrmoteras* Forel, 1893

Tribe Gesomyrmecini Ashmead, 1905

- Gesomyrmex* Mayr, 1968
Santschiella Forel, 1916
*Sicilomyrmex** Wheeler W.M., 1915

Gesomyrmecini Incertae sedis:

- Prodinmorphomyrmex** Wheeler W.M., 1915

Tribe Myrmecorhynchini Wheeler W.M., 1917

- Myrmecorhynchus* André, 1896
Notoncus Emery, 1895
Pseudonotoncus Clark, 1934

Tribe Oecophyllini Emery, 1895

- Oecophylla* Smith F., 1860

Tribe Gigantiopini Ashmead, 1905

- Gigantiops* Roger, 1863

Tribe Camponotini Forel, 1878

- Calomyrmex* Emery, 1895
Camponotus Mayr, 1861
*Chimaeromyrma** Dlussky, 1988
Echinopla Smith F., 1857
Forelophilus Kutter, 1931
Opisthopis Dalla Torre, 1896
Overbeckia Viehmeyer, 1916
Phasmomyrmex Stitz, 1910
Polyrhachis Smith F., 1857
*Pseudocamponotus** Carpenter, 1930

Tribe Notostigmatini Bolton, 2003

- Notostigma* Emery, 1920

Tribe Formicini Latreille, 1809

- Alloformica* Dlussky, 1969
Bajcaridris Agosti, 1994
Cataglyphis Foester, 1850
Formica Linnaeus, 1758
*Glaphyromyrmex** Wheeler W.M., 1915
Polyergus Latreille, 1804
Proformica Ruzsky, 1902
*Protoformica** Dlussky, 1967
Rossomyrmex Arnol'di, 1928

Tribe Melophorini Forel, 1912

- Melophorus* Lubbock, 1883

Formicinae Incertae sedis:

- Imhoffia** Heer, 1850
*Kyromyrma** Grimaldi & Agosti, 2000
*Leucotaphus** Donisthorpe, 1920
*Protrechina** Wilson, 1985
*Tylolasius** Zhang, 1989

Subfamily Myrmeciinae

Tribe Myrmeciini Emery, 1877

- Myrmecia* Fabricius, 1804

Tribe Prionomyrmecini Wheeler W.M. 1915

- Nothomyrmecia* Clark, 1934
*Prionomyrmex** Mayr, 1868

Myrmeciinae Incertae sedis:

- Archymyrmex** Cockerell, 1923
*Ameghinoia** Viana & Haedo Rossi 1957
*Polanskiella** Rossi de García, 1983

Subfamily Pseudomyrmecinae

Tribe Pseudomyrmecini M.R. Smith, 1952

- Myrcidris* Ward, 1990
Pseudomyrmex Lund, 1831
Tetraponera Smith F., 1852

Subfamiliy Cerapachyinae

Tribe Acanthostichini Emery, 1901

- Acanthostichus* Mayr, 1887

Tribe Cerapachyini Forel, 1893

Cerapachys Smith F., 1857
Simopone Forel, 1891
Sphinctomyrmex Mayr, 1866

Tribu Cylindromyrmecini Emery, 1901

Cylindromyrmex Mayr, 1870

Subfamily Ecitoninae**Tribe Cheliomyrmecini Wheeler, 1921**

Cheliomyrmex Mayr, 1870

Tribe Ecitonini Forel, 1893

Ecton Latreille, 1804
Labidus Jurine, 1807
Neivamyrmex Borgmeier, 1940
Nomamyrmex Borgmeier, 1936

Subfamily Leptanilloidinae**Tribe Leptanilloidini Bolton, 1992**

Asphinctanilloides Brandão, Diniz, Agosti &
 Delabie, 1999
Leptanilloides Mann, 1923

Subfamily Aenictinae**Tribe Aenictini Emery, 1901**

Aenictus Shuckard, 1840

Subfamily Dorylinae**Tribe Dorylini Leach, 1815**

Dorylus Fabricius, 1793

Subfamily Aenictogitoninae**Tribe Aenictogitonini Ashmead, 1905**

Aenictogiton Emery, 1901

Subfamily Leptanillinae**Tribe Anomalomyrmini Taylor, 1990**

Anomalomyrma Taylor, 1990
Protanilla Bolton, 1990

Tribe Leptanillini Emery, 1910

Leptanilla Emery, 1870
Phaulomyrma Wheeler G.C.
 & Wheeler E.W., 1930
Yavnella Kugler J., 1987

Subfamily Amblyoponinae**Tribe Amblyoponini Forel, 1893**

Adetomyrma Ward, 1994
Amblyopone Erichson, 1842
Bannapone Xu, 2000
*Casaleia** Pagliano & Scaramozzino, 1990
Concoctio Brown, 1974
Myopopone Roger, 1861
Mystrum Roger, 1862
Onychomyrmex Emery, 1895
Prionopelta Mayr, 1866

Tribe Apomyrmini Dlussky & Fedoseeva, 1988

Apomyrma Brown, Gotwald & Lévieux, 1971

Amblyoponinae *Incertae sedis*:

Paraprisionopelta Kusnezov, 1955

Subfamily Ponerinae**Tribe Ponerini Lepeletier, 1835**

Anochetus Mayr, 1861
Asphinctopone Santschi, 1914
Belonopelta Mayr, 1870
Centromyrmex Mayr, 1866
Cryptopone Emery, 1893
Diacamma Mayr, 1862
Dinoponera Roger, 1861
Dolioponera Brown, 1974
Emeryopone Forel, 1912
Harpegnathos Jerdon, 1851
Hypoponera Santschi, 1938
Leptogenys Roger, 1861
Loboponera, Bolton & Brown, 2002
Myopias Roger, 1861
Odontomachus Latreille, 1804
Odontoponera Mayr, 1862
Pachycondyla Smith F., 1858
Phrynoponera Wheeler W.M., 1920
Plectroctena Smith F., 1920
Ponera Latreille, 1804
Psalidomyrmex André, 1890

Simopelta Mann, 1922
Strebognathus Mayr, 1862

Tribe Thaumatomymecini Emery, 1901

Thaumatomyrmex Mayr, 1887

Tribe Platythyreini Emery, 1901

Platythyrea Roger, 1893

Ponerinae *Incertae sedis*:

*Archiponera** Carpenter, 1930
*Poneropsis** Heer, 1867
*Protopone** Dlussky, 1988

Subfamily Ectatomminae

Tribe Ectatommini Emery, 1895

Ectatommma Smith F., 1858
Gnampogenys Roger, 1863
Rhytidoponera Mayr, 1862

Ectatommini *Incertae sedis*:

*Electroponera** Wheeler W.M., 1915

Tribe Typhlomyrmecini Emery, 1911

Typhlomyrmex Mayr, 1862

Ectatomminae *Incertae sedis*:

*Canapone** Dlussky, 1999

Subfamily Heteroponerinae

Tribe Heteroponerini Bolton, 2004

Acanthoponera Mayr, 1862
Heteroponera Mayr, 1887

Heteroponerini *Incertae sedis*:

Aulacopone Arnol'di, 1930

Subfamily Paraponerinae

Tribe Paraponerini Emery, 1901

Paraponera F. Smith, 1858

Subfamily Proceratiinae

Tribe Proceratiini Emery, 1895

*Bradoponera** Mayr, 1868
Discothyrea Roger, 1863
Proceratium Roger, 1863

Tribe Probolomyrmecini Perrault, 2000

Probolomyrmex Mayr, 1901

Subfamily Agroecomyrmecinae

Tribe Agroecomyrmecini Carpenter, 1930

*Agroecomyrmex** Wheeler W.M., 1910
*Eulithomyrmex** Wheeler W.M. 1910
Tatuidris Brown & Kempf, 1968

Subfamily Myrmicinae

Tribe Adelomyrmecini Fernández, 2004

Adelomyrmex Emery, 1897
Cryptomyrmex Fernández, 2004
Baracidirs Bolton, 1981

Tribe Basicerotini Brown, 1949

Basiceros Schulz, 1906
Eurhopalothrix Brown & Kempf, 1961
Octostruma Forel, 1912
Protalaridris Brown, 1980
Rhopalothrix Mayr, 1870
Talaridris Weber, 1941

Tribe Dacetini Forel, 1892

Acanthognathus Mayr, 1887
Colobostruma Wheeler W.M., 1927
Dacetum Perty, 1833
Epopostruma Forel, 1895
Mesostruma Brown, 1948
Microdacetum Santschi, 1913
Orectognathus Smith F., 1853
Pyramica Roger, 1862
Strumigenys F. Smith, 1860

Tribe Phalacromyrmecini Dlussky & Fedoseeva, 1988

Ishakidris Bolton, 1984
Phalacromyrmex Kempf, 1960
Pilotrochus Brown, 1978

Tribe Cataulacini Emery, 1895*Cataulacus* Smith F., 1853**Tribe Cephalotini M.R. Smith, 1949***Cephalotes* Latreille, 1802*Procryptocerus* Emery, 1887**Tribe Attini F. Smith, 1858***Acromyrmex* Mayr, 1865*Apterostigma* Mayr, 1865*Atta* Fabricius, 1804*Cyphomyrmex* Mayr, 1862*Mycetagoicus* Brandão & Mayhé-Nunes, 2001*Mycetarotes* Emery, 1913*Mycetophylax* Emery, 1913*Mycetosoritis* Wheeler, 1907*Mycocepurus* Forel, 1893*Myrmicocrypta* Smith F., 1860*Pseudoatta* Gallardo, 1916*Sericomyrmex* Mayr, 1865*Trachymyrmex* Forel, 1893

Attini ichnotaxon:

Attaichnus Laza, 1982**Tribe Blepharidattini Wheeler & Wheeler, 1991***Blepharidatta* Wheeler W.M., 1915*Wasmannia* Forel, 1893**Tribe Stenammini Ashmead, 1905***Ancyridris* Wheeler W.M., 1935*Bariamyrmra* Lattke, 1990*Calyptomyrmex* Emery, 1887*Cyphoidris* Weber, 1952*Dacatria* Rigato, 1994*Dacetinops* Brown & Wilson, 1957*Docroaspis* Emery, 1908*Indomyrma* Brown, 1986*Lachnomyrmex* Wheeler W.M., 1910*Lasiomyrma* Terayama & Yamane, 2000*Lordomyrma* Emery, 1897*Proatta* Forel, 1912*Rogeria* Emery, 1894*Rostromyrmex* Rosciszewski, 1994*Stenamma* Westwood, 1839*Tetramorium* Bolton, 1991*Vollenhovia* Mayr, 1865Stenammini *incertae sedis*:*Ilemomyrmex** Wilson, 1985**Tribe Solenopsidini Forel, 1893***Alderzia* Forel, 1902*Allomerus* Mayr, 1878*Anillomyrma* Emery, 1913*Bondroitia* Forel, 1911*Carebara* Westwood, 1840*Diplomorium* Mayr, 1901*Machomyrma* Forel, 1895*Mayriella* Forel, 1902*Megalomyrmex* Forel, 1885*Monomorium* Mayr, 1855*Oxyepoecus* Santschi, 1926*Phacota* Roger, 1862*Pheidologeton* Mayr, 1862*Solenopsis* Westwood, 1840*Tranopelta* Mayr, 1866Solenopsidini *incertae sedis*:*Hypopomyrmex** Emery, 1891*Oxyidris** Wilson, 1985**Tribe Myrmicini Lepeletier, 1835***Eutetramorium* Emery, 1899*Huberia* Forel, 1890*Hyloomyrma* Forel, 1912*Manica* Jurine, 1807*Myrmica* latreille, 1804*Pogonomyrmex* Mayr, 1868*Secostruma* Bolton, 1988Myrmicini *incertae sedis*:*Nothomyrmica** Wheeler W.M., 1915**Tribe Tetramoriini, Emery 1895***Anergates* Forel, 1874*Decamorium* Forel, 1913*Rhoptromyrmex* Mayr, 1901*Strongylognathus* Mayr, 1853*Teleutomyrmex* Kutter, 1950*Tetramorium*, Mayr 1855**Tribe Pheidolini Emery, 1877***Anisopheidole* Forel, 1914*Aphaenogaster* Mayr, 1853

Chimaeridirs Wilson, 1989
Goniomma Emery, 1895
Kartidris Boton, 1991
Lophomyrmex Emery, 1892
Messor Forel, 1890
Ocymyrmex Emery, 1886
Oxyopomyrmex André, 1881
Pheidole Westwood, 1839

Pheidolini incertae sedis:

*Lonchomyrmex** Mayr, 1867
*Paraphaenogaster** Dlussky, 1981

Tribe Lenomyrmicini Bolton, 2003

Lenomyrmex Fernández & Palacio, 1999

Tribe Paratopulini Bolton 2003

Paratopula Wheeler W.M., 1919

Tribe Crematogastrini Forel, 1893

Crematogaster Lund, 1831
Recurvidris Bolton, 1992

Tribe Ankylomyrmini Bolton, 2003

Ankylomyrma Bolton, 1973

Tribe Liomyrmecini Bolton, 2003

Liomyrmex Mayr, 1865

Tribe Meranoplini Emery, 1914

Meranoplus Smith F., 1853
*Parameramoplus** Wheeler W.M., 1915

Tribe Myrmicariini Forel, 1893

Myrmicaria Saunders W.W., 1842

Tribe Formicoxenini Forel, 1893

Atopomyrmex André, 1889
Cardiocondyla Emery, 1869
Chalepoxenus Menozzi, 1923
Dilobocondyla Santschi, 1910
Formicoxenus Mayr, 1855
Gauromyrmex Menozzi, 1933
Harpagoxenus Forel, 1893
Leptothorax Mayr, 1855

Myrmoxenus Ruzsky, 1902
Nesomyrmex Wheeler W.M., 1910
Ochetomyrmex Mayr, 1878
Peronymrmex Viehmeyer, 1922
Podomyrma Smith F., 1859
Poecilomyrma Mann, 1921
Protomognathus Wheeler W.M., 1905
Romblonella Wheeler W.M., 1935
Rotastruma Bolton, 1991
Stereomyrmex Emery, 1901
Temnothorax Mayr, 1861
Terataner Emery, 1912
Vombisidris Bolton, 1991
Xenomyrmex Forel, 1885

Formicoxenini incertae sedis:

*Stigmomyrmex** Mayr, 1868
Tricytarus Donisthorpe, 1947

Tribe Stegomyrmecini Wheeler W.M., 1922

Stegomyrmex Emery, 1912

Tribe Myrmecinini Ashmead, 1905

Acanthomyrmex Emery, 1893
Myrmecina Curtis, 1829
Perissomyrmex Smith M.R., 1947
Pristomyrmex Mayr, 1866

Myrmecinini incertae sedis:

*Enneamerus** Mayr, 1868
*Stiphromyrmex** Wheeler W.M., 1915

Tribe Metaponini Forel, 1911

Metapone Forel, 1911

Tribe Melissotarsini Emery, 1901

Melissotarsus Emery, 1877
Rhopalomastix Forel, 1900

Myrmicinae incertae sedis:

*Attopsis** Heer, 1850
*Cephalomyrmex** Carpenter, 1930
*Electromyrmex** Wheeler W.M., 1910
*Eocenidris** Wilson, 1985
*Eoformica** Cockerell, 1921
*Eomyrmex** Hong, 1974
*Miosolenopsis** Zhang, 1989
*Zhangidris** Bolton, 2003

Subfamily Armaniinae***Tribe Armaniini* Dlussky, 1983**

- Archaeopone** Dlussky, 1975
*Armania** Dlussky, 1983
*Dolichomyrma** Dlussky, 1975
*Kethania** Dlussky, 1999
*Poneropterus** Dlussky, 1983
*Pseudarmania** Dlussky, 1983

Subfamily Sphecomyrminae***Tribe Haidomyrmecini* Bolton, 2003**

- Haidomyrmex** Dlussky, 1996

Tribe Sphecomyrmini* Wilson & Brown, 1967

- Baikuris** Dlussky, 1987
*Cretomyrma** Dlussky, 1975
*Dlusskyidris** Bolton, 1994
*Sphecomyrma** Wilson & Brown, 1967

Subfamily Brownimeciinae***Tribe Brownimeciini* Bolton, 2003**

- Brownimecia** Grimaldi, Agosti
& Carpenter, 1997

Subfamily Formiciinae***Tribe Formiciini* Lutz, 1986**

- Formicium** Westwood, 1854

Genera *incertae sedis* in Formicidae

*Cretopone** Dlussky, 1975 (Bolton 2003 put this name as *incertae sedis* in their poneromorphs subfamilies)

*Petropone** Dlussky, 1975 (Bolton 2003 put this name as *incertae sedis* in their poneromorphs subfamilies)

*Calyptites** Scudder, 1877

*Cariridris** Brandao & Martins-Neto, 1990

Condylodon Lund, 1831

Hypochyra Buckley, 1866

Noonilla Petersen, 1968

Syntaphus Donisthorpe, 1920

Subfamily *incertae sedis* in Formicidae:

Subfamily Paleosminthurinae*


Appendix 4 / Anexo 4

Total count of the species and genera for each subfamily (a) and country (b). The total numbers don't include species which description is under publication process.


Conteo total de las especies y géneros para cada una de las Subfamilias (a) y cada uno de los países (b). Los totales no incluyen las especies cuya descripción se encuentra en proceso de publicación.

Countries	br	co	cr	ar	me	pn	gi	pe	bo	vn	su	gf	ec	tt	pr	am	gu	us	ho	an	ni	ur	be	ch	ad	bh	es	ber	az	Total
Subfamilies																											Total			
Dolichoderinae	92	38	28	50	17	29	31	18	30	20	29	29	6	20	14	14	18	11	6	7	2	6	7	14	5	1	1	0	0	228
Formicinae	159	62	47	66	83	36	50	41	46	26	41	42	18	31	26	39	20	26	10	21	9	20	8	20	0	9	0	5	0	409
Myrmeciinae	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
Pseudomyrmecinae	64	32	35	12	31	31	23	32	16	17	21	21	10	19	13	14	23	10	8	3	9	3	7	2	10	3	2	0	0	144
Cerapachyinae	17	6	5	6	5	2	2	6	5	5	2	2	5	0	3	2	2	1	1	0	0	1	1	1	4	0	0	0	0	39
Ectitoninae	68	18	26	30	34	21	22	21	27	7	22	23	20	5	22	0	11	10	5	2	7	7	5	1	1	0	1	0	0	141
Leptanilloidinae	3	2	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8
Amblyoponinae	9	0	2	5	2	1	1	0	1	2	2	1	0	1	0	2	3	0	1	1	1	0	2	0	0	0	0	0	0	20
Ponerinae	105	44	45	20	26	30	41	25	24	26	38	37	29	16	15	17	18	4	22	14	4	7	5	1	8	8	0	0	0	220
Ectatomminae	47	24	20	12	8	19	18	13	21	25	18	18	10	6	11	5	3	1	13	2	2	0	0	5	0	0	0	0	102	
Heteroponerinae	10	4	1	2	1	1	0	1	2	2	0	1	1	0	1	0	1	0	0	0	0	1	0	1	0	0	0	0	0	17
Paraponerinae	1	1	1	0	0	1	1	1	0	1	1	1	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	2	
proceratiinae	2	5	3	2	5	7	1	1	2	1	1	1	1	0	0	0	2	1	3	0	0	0	0	5	0	0	0	0	33	
Agroecomyrmecinae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	
Myrmicinae	581	320	306	252	231	195	148	173	149	154	108	106	133	109	95	83	74	79	70	56	56	42	41	18	21	23	13	0	1	1553
Paleosminthurinae	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	
Total Countries	1159	556	520	458	443	373	338	332	324	286	283	282	235	208	201	176	173	144	137	109	91	90	74	60	60	44	18	5	1	

Countries	br	co	me	ar	pn	gi	cr	bo	su	gf	vn	pe	pr	ec	gu	ho	tt	am	an	ur	us	ni	be	ch	bh	ad	es	beraz	Total	
Subfamilies																											Total			
Dolichoderinae	6	5	6	6	6	5	5	5	5	5	4	4	4	3	5	3	3	6	5	4	3	1	4	3	1	4	1	0	0	9
Formicinae	6	6	12	6	5	6	5	5	6	6	6	5	5	3	4	2	4	5	5	4	6	2	3	6	3	0	0	3	0	15
Myrmeciinae	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
Pseudomyrmecinae	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0	0	0	2
Cerapachyinae	9	5	1	1	2	2	5	3	2	2	4	4	2	5	2	1	0	2	0	1	1	0	1	1	0	4	0	0	0	4
Ectitoninae	5	5	4	4	4	5	3	3	5	5	4	4	4	4	3	4	3	0	1	3	2	3	3	1	0	1	1	0	5	
Leptanilloidinae	1	1	0	0	0	0	1	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
Amblyoponinae	2	0	2	3	1	1	1	2	1	2	0	0	0	2	1	1	1	1	1	0	1	0	1	0	0	0	0	0	0	3
Ponerinae	10	7	7	7	6	7	7	8	6	5	6	6	5	5	6	8	5	7	6	5	4	4	3	1	5	1	0	0	0	12
Ectatomminae	3	3	3	3	3	3	3	3	3	3	3	3	3	2	3	3	2	2	2	2	2	2	2	2	2	0	0	0	3	
Heteroponerinae	4	4	1	2	3	2	3	4	2	3	2	3	3	3	1	0	1	0	0	1	0	2	0	1	0	0	0	0	0	2
Paraponerinae	1	1	0	0	1	1	1	1	0	1	1	1	0	1	1	0	0	0	0	1	0	0	0	1	0	0	0	0	1	
proceratiinae	2	2	1	2	3	1	3	1	1	1	2	1	0	1	0	1	0	0	1	0	2	0	0	0	1	0	0	0	3	
Agroecomyrmecinae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	
Myrmicinae	43	33	33	32	33	32	28	29	30	30	24	23	25	23	24	24	27	20	17	14	15	17	12	7	12	6	8	0	1	57
Paleosminthurinae	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Total Countries	95	73	71	68	68	66	66	65	64	63	58	55	53	52	51	48	47	44	38	36	36	34	29	24	24	23	14	5	1	



Hormiga obrera típica (*Paratrechina longicornis*). Tomado de Smith (1947)



Obrera de *Sphecomyrmex freyi* (Redibujado de Wilson et al. 1967b)

Guidelines for authors

(see also: <http://www.humboldt.org.co/download/guia.pdf>)

Manuscript Preparation

Authors interested in contributing to Biota Colombiana may submit manuscripts by email, or by sending a diskette with the file by regular mail. Any word-processing program may be used for text (Word is recommended). Taxonomic lists and any other table are better to be sent in spread format (Excel is recommended). Lists for taxonomic groups inferior to genera, such as subgenera or species groups, will not be accepted.

Submitted manuscripts must be accompanied by a cover letter which clearly indicates:

1. Author's (s) complete name(s) and mailing address (es). (It is important to include email address (es) for direct communication.)
2. The complete title of the article.
3. Names, sizes, and types of files submitted.
4. A list of the names and addresses of at least three (3) reviewers³ who are qualified to evaluate the manuscript.

Lists Structure

Biota Colombiana is a completely bilingual publication, with all contents in both Spanish and English. Authors may present manuscripts in either language, or in both. Manuscripts that are submitted in only one language will be translated by the editorial office, but must subsequently be reviewed and approved by the author(s). Biota Colombiana is divided into six sections: 1) Neotropical Lists; 2) National Lists; 3) Regional Lists; 4) Notes and Comments 5) Bibliography Reviews and 6) Bibliography Newness. The organizational structure for articles in the first three sections is: a) Title and Authors, b) Key words, c) Introduction, d) Synopsis or Conspectus, e) Taxonomic List⁴, f) Literature Cited, g) Appendices.

Evaluation

The evaluation's final answer could be: a) Accepted (In this case we assume that not any change, omission or addition to the article is required and the presented form is recommended to be published); b) Conditional acceptance (The article is accepted and recommended to be published but it must have the corrections indicated by the evaluator); y c) Rejected (in this case the evaluator considers that the article presentation contents and/or form are not adjusted to the publication scheme of Biota Colombiana).

INSTRUCTIONS

- The manuscript should be configured for letter size paper, with margins of 2.5 cm on all sides, and should be single-spaced and left-aligned (including title and bibliography).
- The tables in which the lists are presented do not require any specific page configuration. All the columns must fit across a single page (columns on separate pages are not acceptable). Avoid the use of borders of any type in editing the tables.
- In setting up the tables, use column headings ONLY for the first page. Always start additional pages with a new record; NEVER cut the information in one record so that it continues on a subsequent page.
- Use Times New Roman or Arial font, size 12, for all texts. Use size 10 text in tables. Avoid the use of bold or underlining (italics are recommended where considered necessary for emphasis or clarity).
- All pages of text (with the exception of the title page) should be numbered. Page numbers should be located at the lower right of the page.
- Each reference in the bibliography should be cited in the text, and vice versa. Only cite published material or that which is in press. To cite a single author use the format: Gómez (1995) or (Gómez 1995); to cite two authors: Otero & Blum (1970) or (Otero & Blum 1970); for more than two authors: Silva et al. (1998) or (Silva et al. 1998). To cite various references at the end of a sentence, order them chronologically, from oldest to most recent, and then alphabetically: (Otero & Blum 1970; Gómez 1995; MacArthur 1995; Silva et al. 1998). Use the same format for citing the same author more than once: Santos (1995, 1997), or (Santos 1995, 1997), or among several citations: (Otero & Blum 1970; Santos 1995, 1997; Silva et al. 1998).
- In the bibliography, write the complete reference, according to the following pattern:
 - **Journals:** Agosti D., C. R. F. Brandao, S. Diniz 1999 The New World species of the subfamily Leptanilloidinae (Hymenoptera: Formicidae) Systematic Entomology 24:14-20
 - **Books:** Watkins W. F. Jr. 1976 The identification and distribution of New World Army Ants Waco, Texas, 102pp.
 - **Chapters:** Fernández F., E. E. Palacio, W. P. MacKay 1996 Introducción al estudio de las hormigas (Hymenoptera: Formicidae) de Colombia pp.349-412 In: G. D. Amat, G. Andrade, F. Fernández (eds.) Insectos de Colombia, Estudios Escogidos Academia Colombiana de Ciencias Exactas, Fisicas y Naturales & Centro Editorial Javeriano, Bogotá D.C.
- Use a space and DO NOT USE PERIODS to separate author from year, year from title, etc. In the case of several authors, separate them by commas; do not use &, y, or and before the last author. The volume, issue, and page numbers should be presented in continuous format, without spacing. Check all citations for names, accent marks, and spelling. In the case of journal articles, write the full name of the journal, without abbreviations. Do not use hanging indents or underlining in the bibliography.
- To refer to numbers within the text, write only the numbers from one to ten in words (e.g., five species), unless combining with other numbers or measures (e.g., 3 genera and 7 species; 8 km²). Use commas to separate thousands (with the exception of 1000), and commas for decimal points. For percentages, use the symbol % followed by the value, without spaces.
- Within the text, refer to personal communication or observations in the following way: E. Palacio (pers. comm.); (E. Palacio, pers. comm.); C. Sarmiento (pers. obs.); (C. Sarmiento, pers. obs.).
- In neotropical or national lists, use the standardized abbreviations at the end of this issue. If it's possible, use different columns for geopolitical and biogeographical subdivisions. When using several abbreviations for the same record, order them alphabetically and separate them by spaces. DO NOT USE COMMAS (e.g., am cq gu vu).
- If you use a biogeographical referent in the Distribution column of national lists that is different from region, include a key to the abbreviations. If the biogeographical system you use has been published previously in earlier issues of the journal, it is recommended that you use the same key for the abbreviations.
- Read these guidelines completely before submitting a manuscript for evaluation. If you have questions, consult the editor of Biota Colombiana.

³ The editor and members of the editorial committee reserve the right to select the reviewers for each manuscript and to choose anonymous reviewers different from those initially suggested by the author(s).

⁴ This is the most important component of the manuscript. All Neotropical, national, and/or regional lists must be presented in table form. Neotropical and national lists should include, at minimum, the following information: a) Taxa, b) Distribution, c) Altitudinal Distribution, d) Reference, e) Observations.

Guía para autores

(ver también: <http://www.humboldt.org.co/download/guia.pdf>)

Preparación del manuscrito

Los autores que deseen hacer sus contribuciones a *Biota Colombiana* pueden someter sus documentos a través de correo electrónico, o enviando por correo una copia magnética del documento. Para la elaboración de los textos del manuscrito se puede usar un procesador de palabra cualquiera (preferiblemente MS Word); los listados y cualquier otra tabla que si incluya en el manuscrito deben ser hechos en una hoja de cálculo (en cuyo caso se recomienda MS Excel). Los listados taxonómicos pueden provenir de investigación original, consulta de literatura, revisión de colecciones biológicas, o cualquier otra fuente susceptible de verificación.

Para someter un manuscrito es necesario anexar una carta de intención en la que se indique claramente:

1. Nombre(s) completo(s) del(os) autor(es), y direcciones para envío de correspondencia (es indispensable suministrar una dirección de correo electrónico para comunicación directa)
2. Título completo del Manuscrito
3. Nombres, tamaños y tipos de archivos suministrados
4. Lista de mínimo 2 o 3 revisores¹ que puedan evaluar el manuscrito, con sus respectivas direcciones

Estructura de los listados:

Biota Colombiana es una publicación bilingüe, con sus contenidos en español e inglés. Los autores pueden presentar sus manuscritos en cualquiera de los dos idiomas, o en ambos si existe dicha facilidad. *Biota Colombiana* está organizada en seis secciones: 1) Listados Neotropicales; 2) Listados Nacionales; 3) Listados Regionales; 4) Notas y Comentarios; 5) Reseñas Bibliográficas y 6) Novedades Bibliográficas. La estructura para las tres primeras secciones es: a) Título y Autores, b) Palabras clave, c) Introducción, d) Sinopsis o Conspectus, e) Listado Taxonómico², f) Bibliografía, g) Anexos (Para mayores detalles por favor visitar la siguiente página web: <http://www.humboldt.org.co/download/guia.pdf>).

Evaluación del manuscrito

Los manuscritos sometidos serán revisados por pares científicos calificados cuya respuesta final de evaluación puede ser: a) Aceptado (en cuyo caso se asume que no existe ningún cambio, omisión u adición al artículo, y que se recomienda su publicación en la forma actualmente presentada); b) Aceptación Condicional (se acepta y recomienda el artículo para su publicación solo si se realizan los cambios indicados por el evaluador); y c) Rechazo (en el cual el evaluador considera que los contenidos y/o forma de presentación del artículo no se ajustan al esquema de publicación de *Biota Colombiana*).

LISTA DE INSTRUCCIONES

- Para la presentación del manuscrito configure las páginas de los textos en tamaño carta, márgenes de 2,5cm en todos los lados, espacio sencillo y alineación hacia la izquierda (incluyendo título y bibliografía).
- Las tablas donde se presenta el listado, no necesitan de una configuración de página específica. Se requiere que todas las columnas estén dentro de una misma página (no se aceptan columnas en páginas separadas). Evite el uso de bordes de cualquier tipo en la edición de las tablas.
- En la construcción de las tablas, use encabezados para las columnas UNICAMENTE en la primera hoja. Continúe siempre las tablas en hojas nuevas con registros nuevos; NUNCA corte la información de un registro para continuarlo en una nueva página.
- Utilice como fuente Times New Roman o Arial, tamaño 12, en todos los textos. Para tablas cambie el tamaño de la fuente a 10. Evite el uso de negritas o subrayados; géneros y especies en itálica.
- Todas las páginas de texto (a excepción de la primera, correspondiente al título), deben numerarse en la parte inferior - derecha de la hoja.
- Cada referencia de la bibliografía debe estar citada en el texto, y viceversa. Para citar un autor use: Gómez (1995) o (Gómez 1995); para dos autores: Otero & Blum (1970) o (Otero & Blum 1970); para más de dos autores: Silva *et al.* (1998) o (Silva *et al.* 1998). Para citar varias referencias al final de una frase ordénelas cronológicamente, de la más antigua a la más reciente, y luego ordénelas alfabéticamente: (Otero & Blum 1970; Gómez 1995; MacArthur 1995; Silva *et al.* 1998); utilice este mismo esquema para citar varias veces un mismo autor: Santos (1995, 1997), o (Santos 1995, 1997), o en medio de varias citas: (Otero & Blum 1970; Santos 1995, 1997; Silva *et al.* 1998).
- En la bibliografía escriba las referencias completas, de acuerdo al siguiente patrón:
 - **Revistas:** Agosti D., C. R. F. Brandao, S. Diniz (1999) The New World species of the subfamily Leptanilloidinae (Hymenoptera: Formicidae) *Systematic Entomology* 24:14-20
 - **Libros:** Watkins W. F. Jr. (1976) The identification and distribution of New World Army Ants Waco, Texas, 102pp.
 - **Capítulos:** Fernández F., E. E. Palacio, W. P. MacKay (1996) Introducción al estudio de las hormigas (Hymenoptera: Formicidae) de Colombia pp.349-412 En: G. D. Amat, G. Andrade, F. Fernández (eds.) Insectos de Colombia, Estudios Escogidos Academia Colombiana de Ciencias Exactas, Físicas y Naturales & Centro Editorial Javeriano, Bogotá D.C.
- Utilice espacios sencillos y NO PUNTOS SEGUIDOS para separar autores de año, año de título, etc. En caso de varios autores sepárelos por medio de comas; no utilice &, y, o and para referirse al último autor. Los volúmenes, números y páginas deben presentarse de manera continua, sin espacios entre sí. Revise todas las citas en cuanto a nombres, acentos y ortografía. Para el caso de artículos de revistas, escriba el nombre completo de la revista, no use abreviaturas. No use sangrías ni subrayados en la presentación de la bibliografía.
- Para referirse a números dentro del texto, escriba en palabras únicamente los números del uno al diez (p.e., cinco especies) a menos que esté combinado con otros números o se trate de medidas (p.e., 3 géneros y 7 especies; 8 km²). Para porcentajes utilice el símbolo % seguido del valor, sin espacios.
- Dentro del texto puede referirse a comunicaciones y/o observaciones personales, de la siguiente forma: E. Palacio (com. pers.); (E. Palacio, com. pers.); C. Sarmiento (obs. pers.); (C. Sarmiento, obs. pers.).
- Para la elaboración de listados neotropicales o nacionales, utilice las abreviaturas suministradas en las páginas finales de cada número de la revista. Utilice ambos tipos de subdivisiones en columnas separadas (geopolítica y biogeográfica). Cuando utilice varias abreviaturas para un mismo registro, ordénelas alfabéticamente y sepárelas entre sí por espacios, NO USE COMAS (por ejemplo: am cq gn vu).
- Si para la columna de Distribución en listados nacionales utiliza un referente biogeográfico distinto a región, debe suministrar de forma separada una clave de las abreviaturas empleadas. Si el esquema biogeográfico utilizado por usted ya ha sido presentado en artículos de números anteriores de la revista, se recomienda usar la misma clave de abreviaturas allí expuesta.
- No se olvide de leer la presente guía en su totalidad antes de someter un manuscrito para evaluación. Cualquier duda o inquietud consultela inmediatamente al editor en jefe de *Biota Colombiana*.

¹ Tanto el Editor en Jefe como los miembros del comité editorial se reservan el derecho de seleccionar los revisores para cada manuscrito; igualmente, los editores podrán buscar revisores anónimos distintos a los sugeridos inicialmente por el(es) autor(es).

² Es el componente más importante del manuscrito. Todos los listados (neotropicales, nacionales y/o regionales) deben ser presentados en forma de tabla. Los listados neotropicales y nacionales deben incluir, como mínimo, la siguiente información: a) Taxa, b) Distribución, c) Rango de distribución altitudinal, d) Referencia, e) Observaciones.

Abreviaturas de Países / Countries Abbreviations




Países / Countries

Antillas Mayores / Greater Antilles
 Antillas Menores / Lesser Antillas
 Argentina / Argentina
 Bahamas / Bahamas
 Belice / Belize
 Bolivia / Bolivia
 Brasil / Brazil
 Colombia / Colombia
 Costa Rica / Costa Rica
 Cuba / Cuba
 Chile / Chile
 Ecuador / Ecuador
 El Salvador / El Salvador
 Estados Unidos / United States

am	Guyana Francesa / French Guiana	gf
an	Guyana / Guyana	gi
ar	Guatemala / Guatemala	gu
bh	Honduras / Honduras	ho
be	Jamaica / Jamaica	ja
bo	México / Mexico	me
br	Nicaragua / Nicaragua	ni
co	Perú / Peru	pe
cr	Panamá / Panama	pn
cu	Paraguay / Paraguay	pr
ch	Surinam / Suriname	su
ec	Trinidad y Tobago / Trinidad and Tobago	tt
es	Uruguay / Uruguay	ur
eu	Venezuela / Venezuela	vn

Departamentos y Regiones Biogeográficas Continentales de Colombia Geopolitical Distribution and Continental Biogeographic Regions of Colombia



Departamentos / Departments


Amazonas
Antioquia
Arauca
Atlántico
Bolívar
Boyacá
Cauca
Cesar
Caldas
Córdoba
Caquetá
Casanare
Cundinamarca
Chocó
Guainía
Guaviare

ama
ant
ara
at
bl
by
cau
ce
cl
cor
cq
cs
cun
cho
gn
gv

Huila
La Guajira
Magdalena
Meta
Nariño
Norte de Santander
Putumayo
Quindío
Risaralda
Santander
San Andrés y Providencia
Sucre
Tolima
Vaupés
Valle del Cauca
Vichada

hu
lg
ma
met
na
ns
pu
qu
ri
snt
sp
suc
to
va
vc
vch

Unidades Biogeográficas de Colombia / Biogeographic units of Colombia



Unidades Biogeográficas / Biogeographic Units

Territorios Insulares Oceánicos Caribeños / Caribbean Oceanic Insular Territories

Territorios Insulares Oceánicos del Pacífico / Pacific Oceanic Insular Territories

Cinturón Arido Pericaribeño / Arid Peri-Caribbean Belt

Macizo de la Sierra Nevada de Santa Marta / Massif of the Sierra Nevada de Santa Marta

Provincia del Chocó-Magdalena / Choco-Magdalena Province

Provincia de la Orinoquia / Orinoquia Province

Provincia de la Guyana / Guyana Province

Provincia de la Amazonia / Amazonian Province

Provincia Norandina / North-Andean Province

I

II

III

IV

V

VI

VII

VIII

IX

Tomado de: Hernández J., A. Hurtado, R. Ortiz, T. Walschburger 1991 Unidades Biogeográficas de Colombia En: Hernández J., R. Ortiz, T. Walshburger, A. Hurtado (Eds.) Estado de la Biodiversidad en Colombia Informe Final Santafé de Bogotá, Instituto Colombiano para el Desarrollo de la Ciencia y la Tecnología “Francisco José de Caldas” – Colciencias

Biota Colombiana Vol. 5 (1), 2004

Una publicación del / A publication of: Instituto Alexander von Humboldt

En asocio con / In collaboration with:

Instituto de Ciencias Naturales de la Universidad Nacional de Colombia
Instituto de Investigaciones Marinas y Costeras - Invemar
Missouri Botanical Garden

Listados Neotropicales / Neotropical Lists

List of Neotropical ants (Hymenoptera: Formicidae) / <i>Lista de las hormigas Neotropicales (Hymenoptera: Formicidae)</i> – F. Fernández & S..Sendoya	3
-------------------------------------------------------------------------------------------------------------------------------------------------------------	---

